

Out of Equilibrium Relativistic Quantum Field Theory — Perturbation Theory and Progress of Phase Transition

A. Niégawa

Department of Physics, Osaka City University, Sumiyoshi-ku, Osaka 558, JAPAN

E-Mail: niegawa@sci.osaka-cu.ac.jp

Abstract

This paper describes perturbative framework, on the basis of closed-time-path formalism, for studying quasiuniform relativistic quantum field systems near equilibrium and nonequilibrium quasistationary systems. At the first part, starting from first principles, we construct perturbative schemes for relativistic complex-scalar-field theory. We clarify what assumption is involved in arriving at a standard perturbative framework and to what extent the $n (\geq 4)$ -point initial correlation functions that are usually discarded in the standard framework can in fact be discarded. Two calculational schemes are introduced, the one is formulated on the basis of the initial particle distribution function and the one is formulated on the basis of the “physical” particle distribution function. Both schemes are equivalent and lead to a generalized relativistic kinetic or Boltzmann equation. At the second part, using the perturbative loop-expansion scheme for an $O(N)$ linear σ model, we analyze how the chiral phase transition proceeds through disoriented chiral condensates. The system of coupled equations that governs the spacetime evolution of the condensate or order-parameter fields is derived. The region where the curvature of the “potential” is negative is dealt with by introducing the random-force fields. Application to simple situations is made.

I. INTRODUCTION

Since mid-fifties, efforts have been made to incorporate quantum field theory with nonequilibrium statistical mechanics [1–4]. Necessity of this incorporation originated from the field of solid-state physics. Since then, rapid progress of the studies of the early Universe and the quark-gluon plasma have further activated this field of research (cf., e.g., [5,6]). The quark-gluon plasma is a system of (anti)quarks and gluons, which is governed by quantum chromodynamics (QCD). According to QCD, there are two types of phase transitions; a chiral phase transition and a confined-deconfined phase transition. According to lattice analyses, critical points for these two transitions are very close. The quark-gluon plasma is the system that is in the deconfined and chiral-symmetric phases and is expected to be produced in relativistic heavy-ion collisions and to have existed in the early Universe. Such quark-gluon plasmas evolve into a confined (or hadronic) and chiral-symmetry breaking phases through the phase transitions. The study of this evolution on the basis of QCD is very involved and some tractable effective theories have been proposed. Among those is a linear σ model, which was proposed long ago [7] and is revived as a toy model that mimic the aspect of the chiral phase transition of QCD¹. A numerous work was reported so far on the “static” properties of this model or its variants. (See, e.g., [9,10]. Earlier work is quoted therein.) Recently, analyses of dynamical evolution of the system from a symmetric phase to a broken symmetric phase have begun (see, e.g., [11]). Such an analysis is still involved and different approximations have been employed by various authors.

The purpose of this paper is twofold. In the first part, we aim at laying perturbative framework for dealing with out-of-equilibrium relativistic quantum complex-scalar-field systems. In the second part, taking up an fermionless $O(N)$ linear σ model, we lay down the framework, based on the perturbative loop expansion of the effective action, for analyzing how the phase transition proceeds through disoriented chiral condensates (see, e.g., [12]). We use the standard framework of nonequilibrium statistical quantum-field theory that is formulated [1–3,13] by employing the closed-time path, $-\infty \rightarrow +\infty \rightarrow -\infty$, in a complex-time plane, which is referred to as the closed-time-path (CTP) formalism. Specializing the density matrix, which characterizes the initial states of the systems, to the one for the Gibbs distribution, the framework reduces to the standard framework of equilibrium thermal quantum field theories [3,14]. Throughout this paper, we are interested in quasiuniform systems near equilibrium or nonequilibrium quasistationary systems, which we simply refer to as out-of-equilibrium systems. This means, in particular, that the phase transition dealt with in the second part proceeds slowly.

The out-of equilibrium systems are characterized by two different spacetime scales: microscopic or quantum-field-theoretical and macroscopic or statistical. The first scale, the microscopic-correlation scale, characterizes the range of radiative correction to reactions taking place in the system while the second scale measures the range of interaction of particles. For a weak-coupling theory, in which we are interested in this paper, the former scale is much smaller than the latter scale. A well-known intuitive picture (cf., e.g., [13]) for dealing with such systems is to separate spacetime into many “cells” whose characteristic size, L^μ ($\mu = 0, \dots, 3$), is in between the microscopic and macroscopic scales. It is assumed that the correlation between different cells is negligible in the sense that microscopic or elementary reactions can be regarded as taking place in a single cell. On the other hand, in a single cell, relaxation phenomena are negligible. Thus, in dealing with a reaction, all the vertices in a Feynman diagram for the reaction are in a single cell. In more precise, propagators are Fourier transformed over a cell and carry an “index” that labels the cell.

Above intuitive picture may be implemented as follows. Let $\Delta(x, y)$ be a generic propagator or two-point function. An inverse Fourier transformation with respect to the relative coordinates $x - y$ yields

$$\Delta(X; P) \equiv \int d^4(x - y) e^{iP \cdot (x - y)} \Delta(x, y), \quad (1.1)$$

where $X = (x + y)/2$ is the midpoint, which is chosen as a label for the cells. Sometimes, X is called the macroscopic spacetime coordinates. It is obvious from the above argument that (1.1) is meaningful for $|P^\mu| \gtrsim 1/L^\mu$. The self-energy part $\Sigma(x, y)$ enjoys similar property.

Development of the phase transition is described by an evolution equation for a set of condensate or order-parameter fields $\vec{\varphi}(x)$. As mentioned above, we assume that the phase transition proceeds slowly. By this we mean that the x dependence of $\vec{\varphi}(x)$ is weak, or more precisely, x of $\vec{\varphi}(x)$ may be treated as a macroscopic spacetime coordinates. It should be noted here that, as the system approaches the critical point of the phase transition, the microscopic correlation scale diverges. This means that the formalism applies to the systems away from the critical region. Recalling that the phase transition occurs when the squared effective mass, $M^2(x)$, of the quasiparticles becomes

¹Lattice simulations of finite-temperature QCD indicates that the chiral phase transition is of second order (see, e.g., [8]). The phase transition described by the linear σ model is of second order.

zero, we can say that the formalism may be used in the region, where $|M^2(x)|$ is not small as compared to the typical scale parameter(s) of the system.

Microscopic reactions discussed above cause change in the density matrix, through which the number densities of quasiparticles change with macroscopic spacetime X^μ . Dealing with this is the subject of the “next stage,” where (weak) X^μ -dependence of $\Delta(X; P)$, $\Sigma(X; P)$, and $\bar{\varphi}(X)$ are explicitly taken into account.

As mentioned above, much work has already been devoted to the issues settled above. However, most work devoted to the first issue (to be dealt with in the first part of this paper) assume (explicitly or implicitly) special forms for the density matrix. Also as mentioned above, the second issue has so far been studied under various approximations. In this paper, starting from first principles, we proceed with discussing matters in a context which is as general as possible.

The plan of the paper is as follows: In Section II, a brief introduction is given to the CTP formalism for out-of-equilibrium quantum field theories. In Sections III - V, taking up a self-interacting relativistic complex-scalar quantum field theory, perturbative calculational schemes are constructed. In Section III, standard representation of the CTP Green functions in the interaction picture is derived from first principles. In the course of derivation, we clarify what assumption is necessary in arriving at the representation. In a standard perturbative scheme, the n (≥ 4)-point initial correlation functions are also usually discarded. We discuss to what extent they can really be discarded. Whenever necessary, they can be incorporated into the scheme (cf. [2]). In Section IV, we deal with the bare propagator (two-point function), which is a building block of perturbation theory. The quasiparticle fields are introduced through a sort of Bogoliubov transformation, which is a generalization of the so-called thermal Bogoliubov transformation in equilibrium thermal field theory [3,4,14]. Quasiparticle fields describe, in a sense, stable modes in the system. With the aid of the quasiparticle picture, we obtain a 2×2 matrix representation for the bare propagator. One of the advantages of this picture is that the elementary-reaction part and the relaxation part, mentioned above, are clearly identified. At the final part of Section IV, we propose two perturbative schemes, which are equivalent to each other. In Section V, through analyzing the structure of 2×2 self-energy matrix, we find the form for the self-energy-part resummed or full propagator. Then, after discussing how the two perturbative schemes work, we show that both schemes are equivalent and lead to a generalized relativistic kinetic or Boltzmann equation. Sections VI - VIII are devoted to the analysis of a fermionless $O(N)$ linear σ model, with the aid of the effective action. [The case of $N = 4$ is of practical interest.] Here, the condensate or order-parameter fields and the quantum fields come in. The latter describe quantum fluctuations around the former. The condensate fields, which define the internal reference frames for the quantum fields, change in spacetime. Taking this fact into account, in Section VI, we construct consistent perturbative loop-expansion scheme for studying how the chiral phase transition proceeds through disoriented chiral condensates. Besides the assumption that the transition proceed slowly, no further assumption is introduced. In Section VII, we compute the effective action to the lowest one-loop order and derive the system of coupled equations that governs the evolution of the condensate fields, the equations which consist of the generalized relativistic Boltzmann equation, the self-consistent gap equation, and the equation of motion for the condensate fields. At the region where the curvature of the “potential” is negative, the effective action Γ develops imaginary part. It is shown that $Im \Gamma$ can be dealt with by introducing the random-force fields that act on the system. The random-force fields cause negative correlation between the condensate fields. In Section VIII, we set up simple situations, for which approximate analytic calculations may be carried out. Section IX is devoted to concluding remarks.

Remarks on notations and the gradient approximation

Let $G(x, y)$ be (a component of) a generic two-point function. We introduce two kinds of spacetime coordinates:

$$\begin{aligned} x - y &: \text{relative or microscopic spacetime coordinates,} \\ X \equiv \frac{x + y}{2} &: \text{center of mass or macroscopic spacetime coordinates.} \end{aligned}$$

As mentioned above, X -dependence of $G(x, y)$ is assumed to be weak. Fourier transform of $G(x, y)$ with respect to $x - y$ reads (cf. (1.1))

$$G(x, y) = \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} G(X; P).$$

Throughout this paper, we use upper-case letters, P , etc., to denote a four-momentum vectors, $P = (P^\mu) = (p_0, \mathbf{p})$, etc. The magnitude of three vector $\mathbf{p}$ will be written as $p = |\mathbf{p}|$. As to the X -dependence of $G(X; P)$, throughout this paper, we keep only up to a first-order-derivative term (gradient approximation),

$$G(X; P) \simeq G(Y; P) + (X - Y) \cdot \partial_Y G(Y; P). \quad (1.2)$$

Functions of the type, $B(\mathbf{x}, \mathbf{y}; z_0)$ also enter in Sections IV and VI. Here $\mathbf{X} = (\mathbf{x} + \mathbf{y})/2$ and z_0 are macroscopic spacetime coordinates, i.e., $B(\mathbf{x}, \mathbf{y}; z_0)$ depends on $\mathbf{X}$ and on z_0 only weakly. Fourier transform of $B(\mathbf{x}, \mathbf{y}; z_0)$ on $\mathbf{x} - \mathbf{y}$ reads

$$B(\mathbf{x}, \mathbf{y}; z_0) = \int \frac{d^3 p}{(2\pi)^3} e^{i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} B(\mathbf{X}, z_0; \mathbf{p}).$$

Whenever necessary, we expand B within the gradient approximation:

$$B(\mathbf{X}, z_0; \mathbf{p}) \simeq B(\mathbf{Y}, x_0; \mathbf{p}) + (z_0 - x_0) \frac{\partial B(\mathbf{Y}, x_0; \mathbf{p})}{\partial x_0} + (\mathbf{X} - \mathbf{Y}) \cdot \nabla_{\mathbf{Y}} B(\mathbf{Y}, x_0; \mathbf{p}). \quad (1.3)$$

For the system of our concern, $|\partial B / \partial x_0| \lesssim |B| / L_0$, $|\partial B / \partial x^i| \lesssim |B| / L^i$ with L^μ as in Section I.

In Sections VI - VIII, there comes in the condensate fields $\vec{\varphi}$. Although the argument of $\vec{\varphi}$ is always macroscopic spacetime coordinates, we write $\vec{\varphi}(x)$ or $\vec{\varphi}(X)$, depending on the contexts.

II. CLOSED-TIME-PATH FORMALISM

Let us start with a generic Lagrangian density

$$\mathcal{L} \left(\phi^{(\alpha)}(x), \phi^{(\beta)}(x), \dots \right)$$

that governs the quantum-field systems. Here the superscripts label collectively the field type and internal degrees of freedom. Following standard procedure, we introduce an oriented closed-time path $C (= C_1 \oplus C_2)$ in a complex time plane, which goes from $-\infty$ to $+\infty$ (C_1) and then returns back from $+\infty$ to $-\infty$ (C_2). The time argument x_0 of the fields, $\phi^{(\alpha)}(x)$'s, is on the time path C . Along this closed time path, the classical action is defined as

$$\begin{aligned} \mathcal{A} &= \int_C dx_0 \int d^3 x \mathcal{L} \left(\phi^{(\alpha)}(x), \phi^{(\beta)}(x), \dots \right) \\ &\equiv \int_C \mathcal{L} \left(\phi^{(\alpha)}(x), \phi^{(\beta)}(x), \dots \right). \end{aligned} \quad (2.1)$$

Define a path-ordered product of field operators along C :

$$T_C \left(\phi^{(\alpha_1)}(x_1), \dots, \phi^{(\alpha_n)}(x_n) \right).$$

Here T_C means to rearrange the field operators as follows: When x_{j0} of $\phi^{(\alpha_j)}(x_j)$ is nearer to the end point of C , $-\infty \in C_2$, than x_{i0} of $\phi^{(\alpha_i)}(x_i)$, $\phi^{(\alpha_j)}(x_j)$ is placed on the left of $\phi^{(\alpha_i)}(x_i)$ with the sign due to the statistics of the relevant fields taken into account. Path-ordered Green function is defined as an average over ensemble prepared at an initial time $t = -\infty$, which is characterized by a density matrix ρ :

$$\begin{aligned} G &\equiv (-i)^{n-1} \text{Tr} \left[T_C \left(\phi^{(\alpha_1)}(x_1) \dots \phi^{(\alpha_n)}(x_n) \right) \rho \right] \\ &\equiv (-i)^{n-1} \langle T_C \left(\phi^{(\alpha_1)}(x_1) \dots \phi^{(\alpha_n)}(x_n) \right) \rangle. \end{aligned} \quad (2.2)$$

In particular, the two-point function is defined as

$$G^{(\alpha\beta)}(x, y) = -i \langle T_C \left(\phi^{(\alpha)}(x) \phi^{(\beta)}(y) \right) \rangle.$$

Let us introduce [2] matrix notation $\hat{G}^{(\alpha\beta)}(x, y)$, whose (i, j) component $G_{ij}^{(\alpha\beta)}(x, y)$ stands for $G^{(\alpha\beta)}(x, y)$ with $x_0 \in C_i$ and $y_0 \in C_j$:

$$\begin{aligned} G_{11}^{(\alpha\beta)}(x, y) &= -i \langle T \left(\phi^{(\alpha)}(x) \phi^{(\beta)}(y) \right) \rangle, \\ G_{12}^{(\alpha\beta)}(x, y) &= -i \sigma \langle \phi^{(\beta)}(y) \phi^{(\alpha)}(x) \rangle, \\ G_{21}^{(\alpha\beta)}(x, y) &= -i \langle \phi^{(\alpha)}(x) \phi^{(\beta)}(y) \rangle, \\ G_{22}^{(\alpha\beta)}(x, y) &= -i \langle \bar{T} \left(\phi^{(\alpha)}(x) \phi^{(\beta)}(y) \right) \rangle, \end{aligned} \quad (2.3)$$

where T ($\bar{T}$) is a time- (an anti-time-) ordering symbol and $\sigma = -1$ when both $\phi^{(\alpha)}$ and $\phi^{(\beta)}$ are fermionic field and $\sigma = +1$ otherwise. In writing down (2.3), the “limit” $\phi^{(\alpha)}(x_0 \in C_1, \mathbf{x}) = \phi^{(\alpha)}(x_0 \in C_2, \mathbf{x})$ has been taken. Throughout this paper, we freely take this limit.²

Let us break up the time path C into two parts:

$$\int_C dx_0 = \int_{C_1} dx_0 + \int_{C_2} dx_0 = \int_{-\infty}^{\infty} dx_0 \Big|_{x_0 \in C_1} - \int_{-\infty}^{\infty} dx_0 \Big|_{x_0 \in C_2}. \quad (2.4)$$

Then, in conformity with the matrix notation, we may rewrite the classical action in (2.1) as

$$\begin{aligned} \mathcal{A} &= \int_{-\infty}^{+\infty} dx_0 \int d^3x \hat{\mathcal{L}}, \\ \hat{\mathcal{L}} &= \mathcal{L}(\phi_1^{(\alpha)}(x), \phi_1^{(\beta)}(x), \dots) - \mathcal{L}(\phi_2^{(\alpha)}(x), \phi_2^{(\beta)}(x), \dots), \end{aligned} \quad (2.5)$$

where $\phi_i^{(\alpha)}(x)$ stands for the field whose time argument x_0 is on C_i ($i = 1, 2$). The field $\phi_1^{(\alpha)}$ is called a type-1 or physical field while $\phi_2^{(\alpha)}$ is called a type-2 field. We call $\hat{\mathcal{L}}$ in (2.5) the hat-Lagrangian density [4,15]. Let us write (2.3) collectively

$$\hat{G}^{(\alpha\beta)}(x, y) = -i \langle \hat{T}(\hat{\phi}^{(\alpha)}(x) {}^t\hat{\phi}^{(\beta)}(y)) \rangle, \quad (2.6)$$

where

$$\hat{\phi}^{(\alpha)}(x) = \begin{pmatrix} \phi_1^{(\alpha)}(x) \\ \phi_2^{(\alpha)}(x) \end{pmatrix}, \quad {}^t\hat{\phi}^{(\beta)}(y) = (\phi_1^{(\beta)}(y), \phi_2^{(\beta)}(y)).$$

Four $G_{ij}^{(\alpha\beta)}$ ($i, j = 1, 2$) in (2.3) are not mutually independent. As a matter of fact, it can readily be found from (2.3) that

$$\sum_{i, j=1}^2 (-)^{i+j} G_{ij}^{(\alpha\beta)} = 0.$$

Retarded, advanced, and correlation functions are related, in respective order, to $G_{ij}^{(\alpha,\beta)}$ through

$$\begin{aligned} G_R^{(\alpha\beta)}(x, y) &= -i\theta(x_0 - y_0) \langle [\phi^{(\alpha)}(x), \phi^{(\beta)}(y)]_{-\sigma} \rangle \\ &= G_{11}^{(\alpha\beta)}(x, y) - G_{12}^{(\alpha\beta)}(x, y), \\ G_A^{(\alpha\beta)}(x, y) &= i\theta(y_0 - x_0) \langle [\phi^{(\alpha)}(x), \phi^{(\beta)}(y)]_{-\sigma} \rangle \\ &= G_{11}^{(\alpha\beta)}(x, y) - G_{21}^{(\alpha\beta)}(x, y), \\ G_c^{(\alpha\beta)}(x, y) &= -i \langle [\phi^{(\alpha)}(x), \phi^{(\beta)}(y)]_{\sigma} \rangle \\ &= G_{11}^{(\alpha\beta)}(x, y) + G_{22}^{(\alpha\beta)}(x, y), \end{aligned} \quad (2.7)$$

where $[A, B]_{\sigma} \equiv AB + \sigma BA$ ($\sigma = \pm$).

III. COMPLEX SCALAR FIELD

We take up self-interacting relativistic complex-scalar field theory with a conserved charge, whose Lagrangian density is

²In this respect, cf. the argument given below in conjunction with (7.49).

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_0 + \mathcal{L}_{int}, \\
\mathcal{L}_0 &= \partial_\mu \phi_B^\dagger \partial^\mu \phi_B - m_B^2 \phi_B^\dagger \phi_B, \\
\mathcal{L}_{int} &= -\frac{\lambda_B}{4} (\phi_B^\dagger \phi_B)^2,
\end{aligned} \tag{3.1}$$

where the suffices “ B ” stand for bare quantity. Aiming at constructing perturbation theories, we go to an interaction picture. One can choose any value for the time $t = t_i$, when the interaction picture and the Heisenberg picture coincide. As in, e.g., [2,3,16], we choose $t_i = -\infty$. For dealing with ultraviolet (UV) divergences, throughout this paper, we adopt the modified minimal-subtraction ($\overline{\text{MS}}$) scheme [17], which is a variant of mass-independent renormalization schemes. We introduce UV-renormalized field ϕ , mass m , and coupling constant λ through

$$\phi_B = \sqrt{Z} \phi, \quad m_B^2 = Z_m m^2, \quad \lambda_B = Z_\lambda \lambda. \tag{3.2}$$

Note that, although the same letter ϕ is used, ϕ here is the UV-renormalized field in the interaction picture, while ϕ 's in Section II are the field in the Heisenberg picture. $\mathcal{L}$ in (3.1) may now be written as

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_0(\phi^\dagger, \phi) + \mathcal{L}_{int}(\phi^\dagger, \phi) + \mathcal{L}_{rc}(\phi^\dagger, \phi), \\
\mathcal{L}_0(\phi^\dagger, \phi) &= \partial_\mu \phi^\dagger \partial^\mu \phi - m^2 \phi^\dagger \phi, \\
\mathcal{L}_{int}(\phi^\dagger, \phi) &= -\frac{\lambda}{4} (\phi^\dagger \phi)^2, \\
\mathcal{L}_{rc}(\phi^\dagger, \phi) &= (Z-1) \partial_\mu \phi^\dagger \partial^\mu \phi - (Z_m Z - 1) m^2 \phi^\dagger \phi - (Z_\lambda Z^2 - 1) \frac{\lambda}{4} (\phi^\dagger \phi)^2.
\end{aligned} \tag{3.3}$$

We shall treat $\mathcal{L}_0$ as the nonperturbative part and $\mathcal{L}_{int} + \mathcal{L}_{rc}$ as the perturbative part. The UV renormalizability of the theory is demonstrated at the end of this section.

The hat-Lagrangian density (cf. (2.5)) reads

$$\hat{\mathcal{L}} = \mathcal{L}(\phi_1^\dagger, \phi_1) - \mathcal{L}(\phi_2^\dagger, \phi_2).$$

From this with (3.3) we can read off the vertex matrices: $\mathcal{L}_{int}$ yields the $(\phi^\dagger \phi)^2$ vertex matrix,

$$i\hat{V}_4 = -i\lambda\hat{\tau}_3 \tag{3.4}$$

with $\hat{\tau}_3$ the third Pauli matrix and $\mathcal{L}_{rc}$ yields

$$\begin{aligned}
i\hat{V}_2 &= -i[(Z-1)\partial^2 + (Z_m Z - 1)m^2] \hat{\tau}_3 \quad (\phi^\dagger \phi \text{ vertex}), \\
i\hat{V}_4' &= -i(Z_\lambda Z^2 - 1)\lambda\hat{\tau}_3 \quad ((\phi^\dagger \phi)^2 \text{ vertex}).
\end{aligned} \tag{3.5}$$

A. 2n-point Green function

We assume that the density matrix ρ commutes with the charge operator,

$$\begin{aligned}
[\rho, Q] &= 0, \\
Q &= i \int d^3x (\phi^\dagger \dot{\phi} - \dot{\phi}^\dagger \phi),
\end{aligned} \tag{3.6}$$

where $\dot{\phi} \equiv \partial\phi(x)/\partial x_0$, etc. (cf. below for undoing this assumption). Then, with obvious notation, nonvanishing path-ordered Green functions (cf. (2.2)) are

$$G(1, \dots, n; 1', \dots, n') = i(-)^n \langle T_C \left\{ \phi(1) \dots \phi(n) \phi^\dagger(1') \dots \phi^\dagger(n') e^{i \int_C \mathcal{L}'} \right\} \rangle,$$

where $\mathcal{L}' \equiv \mathcal{L}_{int} + \mathcal{L}_{rc}$. Here, it is convenient to introduce a generating functional:

$$\begin{aligned}
Z_C[J] &= \langle T_C \exp \left(i \int_C [\mathcal{L}'(\phi^\dagger(x), \phi(x)) + J^*(x) \phi(x) + \phi^\dagger(x) J(x)] \right) \rangle \\
&= e^{i \int_C \mathcal{L}'(-i\delta/\delta J(x), -i\delta/\delta J^*(x))} \langle T_C \exp \left(i \int_C [J^*(x) \phi(x) + \phi^\dagger(x) J(x)] \right) \rangle,
\end{aligned} \tag{3.7}$$

where $J(x)$ and $J^*(x)$ are c-number external-source functions. G is obtained from (3.7) through

$$G(1, \dots, n; 1', \dots, n') = i \frac{\delta^n Z_C[J]}{\delta J^*(1) \dots \delta J^*(n) \delta J(1') \dots \delta J(n')} \Big|_{J^* = J = 0}. \quad (3.8)$$

Hereafter we refer $Z_C[J]$ to as the CTP generating functional. It should be understood [2] that at the end of calculation we set $J_1(x) = J_2(x)$, where $J_i(x) = J(x)$ with $x_0 \in C_i$ ($i = 1, 2$).

As a generalization of the Wick theorem in vacuum theory, we obtain [2,14,18]

$$\begin{aligned} T_C \exp \left(i \int_C [J^*(x)\phi(x) + \phi^\dagger(x)J(x)] \right) \\ = e^{-i \int_C \int_C J^*(x) \Delta_{0C}(x-y) J(y)} : T_C \exp \left(i \int_C [J^*(x)\phi(x) + \phi^\dagger(x)J(x)] \right) :, \end{aligned} \quad (3.9)$$

where the symbol $: \dots :$ indicates to take the normal ordering with respect to the creation and annihilation operators in vacuum theory. In (3.9), $\Delta_{0C}(x-y)$ takes the form, in matrix notation,

$$\begin{aligned} \hat{\Delta}_0(x-y) &= \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \hat{\Delta}_0(P) \\ \hat{\Delta}_0(P) &= \begin{pmatrix} \Delta_F(P) & \theta(-p_0)[\Delta_F(P) - \Delta_F^*(P)] \\ \theta(p_0)[\Delta_F(P) - \Delta_F^*(P)] & -\Delta_F^*(P) \end{pmatrix}. \end{aligned} \quad (3.10)$$

where

$$\Delta_F(P) = \frac{1}{P^2 - m^2 + i\eta} \quad (\eta = 0^+)$$

is the Feynman propagator in vacuum theory. Simple manipulation yields

$$\langle : T_C e^{i \int_C [J^*(x)\phi(x) + \phi^\dagger(x)J(x)]} : \rangle = e^{i\mathcal{W}_C}$$

with

$$\mathcal{W}_C = \sum_{n=1}^{\infty} \frac{i^{n-1}}{n!} \langle : T_C \left\{ \int_C [J^*(x)\phi(x) + \phi^\dagger(x)J(x)] \right\}^n : \rangle_c \quad (3.11)$$

$$= - \sum_{n=1}^{\infty} \frac{1}{(n!)^2} \int_C \dots \int_C \mathcal{W}_{2n}(x_1, \dots, x_n; y_1, \dots, y_n) J^*(x_1) \dots J^*(x_n) J(y_1) \dots J(y_n). \quad (3.12)$$

Here the subscript 'c' denotes connected part and

$$\mathcal{W}_{2n}(x_1, \dots, x_n; y_1, \dots, y_n) = i(-)^n \langle : \phi(x_1) \dots \phi(x_n) \phi^\dagger(y_1) \dots \phi^\dagger(y_n) : \rangle_c \quad (3.13)$$

represents initial correlation. In obtaining (3.12) from (3.11), we have used (3.6), which leads to

$$\langle : \phi(x_1) \dots \phi(x_m) \phi^\dagger(y_1) \dots \phi^\dagger(y_n) : \rangle_c = 0 \quad \text{if } m \neq n.$$

If the condition (3.6) does not hold, the following argument in this paper should be generalized accordingly. Each of the $2n$ time arguments in (3.13) is on either C_1 or C_2 . Then (3.13) consists of 2^{2n} components, all of which are identical. In the interaction representation adopting here, $(\partial^2 + m^2)\phi(x) = 0$, from which we obtain

$$\begin{aligned} (\partial_{x_j}^2 + m^2) \mathcal{W}_{2n}(x_1, \dots, x_n; y_1, \dots, y_n) &= (\partial_{y_j}^2 + m^2) \mathcal{W}_{2n}(x_1, \dots, x_n; y_1, \dots, y_n) = 0 \\ (j &= 1, \dots, n). \end{aligned} \quad (3.14)$$

The bare two-point function or propagator, $\hat{\Delta}(x, y)$, is obtained from (3.7) with $\mathcal{L}' = 0$ and (3.8) - (3.12):

$$\hat{\Delta}(x, y) = \hat{G}(x, y) \Big|_{\mathcal{L}' = 0} = \hat{\Delta}_0(x-y) + \mathcal{W}_2(x; y) \hat{A}_+, \quad (3.15)$$

$$\hat{A}_\pm \equiv \begin{pmatrix} 1 & \pm 1 \\ \pm 1 & 1 \end{pmatrix}. \quad (3.16)$$

Thus we have learned that, in perturbation theory, the nonperturbative part consists of the bare two-point function $i\hat{\Delta}(x, y)$ and the multi-point initial-correlation functions $i(-)^{n-1}\mathcal{W}_{2n}$ ($n \geq 2$). From (3.14), we see that $\mathcal{W}_{2n}$ does not contribute to the on-shell amplitudes. Throughout this paper, we shall assume that $\mathcal{W}_{2n}$'s ($n \geq 2$) can be treated as (nonamputated) “perturbative parts” and, whenever necessary, we include “ $2n$ -point propagator”

$$-i(-)^n\mathcal{W}_{2n}(x_1, \dots, x_n; y_1, \dots, y_n) \quad (n \geq 2) \quad (3.17)$$

into the Feynman rules. We shall discuss, in the following subsection C, to what extent can one discard $\mathcal{W}_{2n}$ ($n \geq 2$).

B. Bare two-point function

We analyze in detail the bare two-point function (3.15), which is a building block of the perturbation theory:

$$\begin{aligned} \hat{\Delta}(x, y) &= -i\langle \hat{T}(\hat{\phi}(x)\hat{\phi}^\dagger(y)) \rangle \\ &= \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x-y)} \hat{\Delta}_0(P) + \mathcal{W}(x; y) \hat{A}_+, \end{aligned} \quad (3.18)$$

$$(\partial_x^2 + m^2)\hat{\Delta}(x, y) = (\partial_y^2 + m^2)\hat{\Delta}(x, y) = -\hat{\tau}_3 \delta^4(x - y). \quad (3.19)$$

Here $\hat{\Delta}_0$ is as in (3.10) and

$$\begin{aligned} \hat{\phi} &= {}^t(\phi_1, \phi_2), \quad \hat{\phi}^\dagger = (\phi_1^\dagger, \phi_2^\dagger), \\ \mathcal{W}(x; y) &\equiv \mathcal{W}_2(x; y) = -i\langle : \phi(x)\phi^\dagger(y) : \rangle_c, \end{aligned} \quad (3.20)$$

$$(\partial_x^2 + m^2)\mathcal{W}(x; y) = (\partial_y^2 + m^2)\mathcal{W}(x; y) = 0. \quad (3.21)$$

Let us find an explicit form for $\mathcal{W}(x; y)$ in terms of the density matrix ρ . For the purpose of later use, we consider a system enclosed in a rectangular parallelepiped. It is natural to take the sizes L^j ($j = 1, 2, 3$), introduced in Section I, as the lengths of edges. As discussed above, when we deal with microscopic reactions, L^j is regarded as large. For simplicity of presentation, we assume that $L^1 = L^2 = L^3 (\equiv L)$. For a set of single-particle wave functions, we employ the periodic boundary condition, on the basis of which $\phi(x)$ may be expanded as

$$\phi(x) = \sum_{\mathbf{p}} \frac{1}{\sqrt{2E_p V}} [a_{\mathbf{p}} e^{-iP \cdot x} + b_{\mathbf{p}}^\dagger e^{iP \cdot x}], \quad (3.22)$$

where $V = L^3$, $E_p = \sqrt{p^2 + m^2}$ is the energy of ϕ with momentum $\mathbf{p}$,

$$\mathbf{p} = \frac{2\pi}{L} \mathbf{n}, \quad n^i = 0, \pm 1, \pm 2, \dots \quad (i = 1, 2, 3),$$

and $a_{\mathbf{p}}$ ($b_{\mathbf{p}}^\dagger$) is an annihilation (a creation) operator of a particle (an antiparticle) of momentum $\mathbf{p}$:

$$\begin{aligned} [a_{\mathbf{p}}, a_{\mathbf{p}'}^\dagger] &= [b_{\mathbf{p}}, b_{\mathbf{p}'}^\dagger] = \delta_{\mathbf{p}\mathbf{p}'}, \\ a_{\mathbf{p}}|0\rangle &= b_{\mathbf{p}}|0\rangle = 0. \end{aligned} \quad (3.23)$$

Substituting (3.22) and its hermitian conjugate $\phi^\dagger$ to (3.20), we obtain,

$$\mathcal{W}(x; y) = -\frac{i}{2V} \sum_{\mathbf{p}, \mathbf{q}} \frac{1}{\sqrt{E_p E_q}} \left[\langle a_{\mathbf{q}}^\dagger a_{\mathbf{p}} \rangle e^{-i(P \cdot x - Q \cdot y)} + \langle b_{\mathbf{q}} b_{\mathbf{p}}^\dagger \rangle e^{i(P \cdot x - Q \cdot y)} \right],$$

where $p^0 = E_p$ and $q^0 = E_q$. Taking the limit $V (\equiv L^3) \rightarrow \infty$ and Fourier transforming on $x - y$, we obtain, after some manipulations,

$$\begin{aligned} \mathcal{W}(X; P) &= \int d^4(x - y) e^{iP \cdot (x - y)} \mathcal{W}(x; y) \\ &= -2\pi i \int d^4Q e^{-iQ \cdot X} \mathcal{F}(P, Q), \end{aligned} \quad (3.24)$$

$$\begin{aligned} \mathcal{F}(P, Q) &\equiv \delta(2P \cdot Q) \delta(P^2 - m^2 + Q^2/4) \\ &\times \sum_{\tau=\pm} \theta(\tau p_0 + \tau q_0/2) \theta(\tau p_0 - \tau q_0/2) F_\tau(\tau \mathbf{p} - \mathbf{q}/2, \tau \mathbf{p} + \mathbf{q}/2), \end{aligned} \quad (3.25)$$

where

$$F_{\pm}(\mathbf{p}, \mathbf{q}) \equiv \langle a_{\pm}^{\dagger}(\mathbf{p}) a_{\pm}(\mathbf{q}) \rangle. \quad (3.26)$$

with

$$a_{+}(\mathbf{p}) = \lim_{V \rightarrow \infty} \sqrt{\frac{2E_p V}{(2\pi)^3}} a_{\mathbf{p}}, \quad a_{-}(\mathbf{p}) = \lim_{V \rightarrow \infty} \sqrt{\frac{2E_p V}{(2\pi)^3}} b_{\mathbf{p}}. \quad (3.27)$$

In the case where the density matrix ρ is diagonal in Fock space, $F_{\pm}(\mathbf{p}, \mathbf{q}) \propto \delta(\mathbf{p} - \mathbf{q})$. Substitution of this into (3.25) yields $\delta(\mathbf{q})$, with which we have $\delta(2P \cdot Q) \rightarrow \delta(q_0)/(2|p_0|)$, so that $\mathcal{F}(P, Q) \propto \delta^4(Q)$. Thus, we see from (3.24) that $\mathcal{W}(X; P)$ is X independent or $\mathcal{W}(x; y)$ ($= \mathcal{W}(x - y)$) is translation invariant. The density matrix dealt with in, e.g., [13] is of this type. The case of equilibrium system is also of this type:

$$F_{\pm}(\mathbf{p}, \mathbf{q}) = 2E_p \delta(\mathbf{p} - \mathbf{q}) N_{\pm}(E_p), \quad (3.28)$$

where $N_{+(-)}$ is a particle (an antiparticle) number density. Equation (3.28) leads to

$$\begin{aligned} \mathcal{W}_{equ}(X; P) &= \mathcal{W}_{equ}(P) \\ &= -2\pi i \delta(P^2 - m^2) [\theta(p_0) N_{+}(E_p) + \theta(-p_0) N_{-}(E_p)], \end{aligned} \quad (3.29)$$

from which we have for the bare two-point function,

$$\hat{\Delta}_{equ}(P) = \hat{\Delta}_0(P) - 2\pi i \delta(P^2 - m^2) [\theta(p_0) N_{+}(E_p) + \theta(-p_0) N_{-}(E_p)], \quad (3.30)$$

a well-known form in equilibrium thermal field theory [3,14].

Let us turn back to the present out-of-equilibrium case. As discussed in Section I, $\mathcal{W}(X; P)$ does not change appreciably in a single spacetime cell. Then, we see from (3.24) that, for $|Q^{\mu}| \gtrsim 1/L^{\mu}$, $\mathcal{F}(P, Q) \simeq 0$, which means

$$F_{\pm}(\pm \mathbf{p} - \mathbf{q}/2, \pm \mathbf{p} + \mathbf{q}/2) \simeq 0 \quad \text{for } q \gtrsim 1/L. \quad (3.31)$$

Here let us make following observation:

- 1) P is the momentum that is conjugate to the relative spacetime coordinates $x - y$, where x and y are in a single spacetime cell.
- 2) In computing an amplitude in perturbation theory, $\mathcal{W}(X; P)$ is to be multiplied by some function $\mathcal{F}(P)$ and is to be integrated out over P^{μ} .

The point 1) means that $\mathcal{W}(X; P)$ is meaningful only for $|p^0| \gtrsim 1/L^0$ and $p \gtrsim 1/L$. Then, the computation of the amplitude is reliable only when

$$\int_0^{\infty} d^4 P \mathcal{W}(X; P) \mathcal{F}(P) \simeq \int_{1/L^0}^{\infty} dp^0 \int_{1/L}^{\infty} d^3 p \mathcal{W}(X; P) \mathcal{F}(P).$$

Let us start from analyzing the “hard- p region,” $p > O(1/L)$. Substituting (3.31) into (3.25), we see that P^{μ} in (3.25) satisfies

$$p_0^2 = p^2 + m^2 + O(1/L^2).$$

To take care of the $O(1/L^2)$ term, we expand $\delta(P^2 - m^2 + Q^2/4)$ in (3.25) in powers of Q^2 ,

$$\delta(P^2 - m^2 + Q^2/4) = \delta(P^2 - m^2) + \frac{Q^2}{4} \frac{d\delta(P^2 - m^2)}{dP^2} + \dots$$

Substituting this back into (3.25), we obtain for $\mathcal{W}(X; P)$ in (3.24), with obvious notation,

$$\begin{aligned} \mathcal{W}(X; P) &= -2\pi i \left[\delta(P^2 - m^2) - \frac{1}{4} \frac{d\delta(P^2 - m^2)}{dP^2} \partial_X^2 + \dots \right] \int \frac{d^3 q}{2|p_0|} e^{-i\mathbf{p} \cdot \mathbf{q} X_0/p_0 + i\mathbf{q} \cdot \mathbf{X}} \\ &\quad \times \sum_{\tau=\pm} [\theta(\tau p_0 + \tau|\mathbf{p} \cdot \mathbf{q}|/2p_0) F_{\tau}(\tau \mathbf{p} - \mathbf{q}/2; \tau \mathbf{p} + \mathbf{q}/2)]. \end{aligned} \quad (3.32)$$

In conformity with the gradient approximation we are adopting (cf. (1.2)), we neglect the term with ∂_X^2 and with “...” in (3.32). Using $\theta(\pm p_0 \pm q_0/2) \simeq \theta(\pm p_0)$ in (3.25), we finally obtain

$$\mathcal{W}(X; P) \simeq -2\pi i \delta(P^2 - m^2) [\theta(p_0) N_+(X; E_p, \hat{\mathbf{p}}) + \theta(-p_0) N_-(X; E_p, -\hat{\mathbf{p}})], \quad (3.33)$$

where $\hat{\mathbf{p}} \equiv \mathbf{p}/p$ and

$$N_{\pm}(X; E_p, \pm \hat{\mathbf{p}}) = \int \frac{d^3 q}{2|p_0|} e^{-i\mathbf{p} \cdot \mathbf{q} X_0/p_0 + i\mathbf{q} \cdot \mathbf{X}} F_{\pm}(\pm \mathbf{p} - \mathbf{q}/2; \pm \mathbf{p} + \mathbf{q}/2).$$

Equation (3.33) is of the standard form, which is used in the literature.

Now we turn to the analysis of the region $p \sim 1/L$. It can readily be seen that the condition for $\mathcal{W}(X; P)$ to take the standard form (3.33) is

$$F_{\pm}(\pm \mathbf{p} - \mathbf{p}/2; \pm \mathbf{p} + \mathbf{p}/2) \simeq 0 \quad \text{for } q \gtrsim |\vec{\delta}_1| < O(1/L). \quad (3.34)$$

This condition is much stronger than (3.31). However, in most practical cases, requiring the weaker condition (3.31) is sufficient. This can be seen as follows. Let $\mathcal{P}$ be a typical scale(s) of the system under consideration. In the case of thermal-equilibrium system, $\mathcal{P}$ is the temperature of the system. As will be seen later, there emerges an effective mass $M_{ind}(X)$. In the case of $m \gg \sqrt{\lambda}\mathcal{P}$, $M_{ind}(X)$ is not much different from m and, for $m \lesssim \sqrt{\lambda}\mathcal{P}$, $M_{ind}(X)$ is of $O(\sqrt{\lambda}\mathcal{P})$:

$$M_{ind}(X) = \text{Max}(m, \sqrt{\lambda}\mathcal{P}). \quad (3.35)$$

Most amplitudes, when computed in perturbation theory to be constructed below, are insensitive to the region $|P^\mu| \ll \text{Max}(m, \sqrt{\lambda}\mathcal{P})$ of $\mathcal{W}(X; P)$, and then the precise form of $\mathcal{W}(X; P)$ in this region is irrelevant. Here, it should be noted that m and $\sqrt{\lambda}\mathcal{P}$ are the scales that are characteristic to microscopic correlations. Then, from the setup in Section I, we assume that $\text{Max}(m, \sqrt{\lambda}\mathcal{P}) > O(1/L)$. Thus, for dealing with the amplitudes of the above-mentioned type, requiring the condition (3.31) is sufficient.

Throughout in the sequel, for the piece $\mathcal{W}(X; P)$ of the propagator (cf. (3.18) and (3.24)), we use (3.33) for all P^μ regions $-\infty < P^\mu < +\infty$. As pointed out above, in computing some quantity, if infrared divergence arises at $p = 0$, we should reanalyze the quantity using the fundamental form, (3.24) with (3.25).

After all this, we find for the Fourier transform of $\hat{\Delta}(x, y)$:

$$\hat{\Delta}(X; P) \simeq \hat{\Delta}_0(P) + \mathcal{W}(X; P) \hat{A}_+, \quad (3.36)$$

with $\hat{\Delta}_0(P)$ as in (3.10) and $\mathcal{W}(X; P)$ as in (3.33). We refer the representation (3.36) to as the $F\bar{F}$ representation, where F ($\bar{F}$) stands for $\Delta_F = (\Delta_0)_{11}$ ($\Delta_F^* = -(\Delta_0)_{22}$) in $\hat{\Delta}_0(P)$, Eq. (3.10).

Infrared region in massless complex-scalar field theory, $\mathcal{R}_{IR}$, is defined in momentum space as $\mathcal{R}_{IR} = \{P^\mu; |P^\mu| \leq O(\sqrt{\lambda}\mathcal{P})\}$. In quantum electrodynamics (QED) and QCD, $\mathcal{R}_{IR} = \{P^\mu; O(g^2\mathcal{P}) < |P^\mu| \leq O(g\mathcal{P})\}$ with g the gauge coupling constant. It is well known that, in such theories as QED and QCD, the infrared region is full of rich structures and contains interesting physics [14,19,20]. Some quantities in QED or QCD is sensitive [14] to a yet “deeper” infrared region, $\mathcal{R}'_{IR} = \{P^\mu; O(g^3\mathcal{P}) < |P^\mu| \leq O(g^2\mathcal{P})\}$. If the inverse size of a spacetime cell, $\mathcal{R}_{cell} = \{P^\mu; |P^\mu| \leq 1/L^\mu\}$, is much smaller than the region $\mathcal{R}_{IR}$ or $\mathcal{R}'_{IR}$, then the above-mentioned interesting physics may be treated within our perturbative schemes. In the opposite case, the interesting physics is hidden outside of our schemes. In more precise, the region $\mathcal{R}_{IR}$ or $\mathcal{R}'_{IR}$ is outside of the region, where the representation (3.36) is meaningful.

C. Multi-point initial correlations

In this subsection, we discuss to what extent $\mathcal{W}_{2n}$ ($n \geq 2$), Eq. (3.13), may be discarded. First of all, we recall that all the 2^{2n} components of $\mathcal{W}_{2n}$ are identical. Then, from (2.7) with (3.18), we see that $\mathcal{W}_2 (= \mathcal{W})$ does not appear in the retarded and advanced Green functions. This means that $\mathcal{W}_2$ does not appear in the two-point correlation in the linear response theory. With the aid of the standard formulae (cf. Sections 2 and 5 of [2]), one can also show that $\mathcal{W}_{2n}$ does not appear in multi-point correlations in the nonlinear as well as the linear response theory.

We restrict our concern to $\mathcal{W}_4$. The argument below may be generalized to the case of general $\mathcal{W}_{2n}$ ($n \geq 3$). Straightforward computation using (3.22) yields

$$\begin{aligned} \mathcal{W}_4(x_1, x_2; y_1, y_2) = & i \sum_{\mathbf{p}_1, \mathbf{q}_1} \sum_{\mathbf{p}_2, \mathbf{q}_2} \left[\prod_i \frac{1}{\sqrt{2E_{p_i} V}} \frac{1}{\sqrt{2E_{q_i} V}} \right] \mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} \\ & \times \left[e^{-i \sum_{j=1}^2 (P_j \cdot x_j - Q_j \cdot y_j)} + e^{i \sum_{j=1}^2 (Q_j \cdot x_j - P_j \cdot y_j)} \right], \end{aligned} \quad (3.37)$$

where $p_1^0 = E_{p_1}$, etc., and

$$\mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} \equiv \langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle - \langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{p}_1} \rangle \langle a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_2} \rangle - \langle a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} \rangle \langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{p}_2} \rangle. \quad (3.38)$$

Here we have assumed for simplicity that $\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle = \langle b_{\mathbf{q}_1}^\dagger b_{\mathbf{q}_2}^\dagger b_{\mathbf{p}_1} b_{\mathbf{p}_2} \rangle$, etc. The first exponential function in (3.37) may be written as, with obvious notation,

$$e^{-i \sum_{j=1}^2 (P_j \cdot x_j - Q_j \cdot y_j)} = e^{-iX \cdot \sum_{j=1}^2 (P_j - Q_j)} e^{-i \sum_{j=1}^2 (P_j \cdot x'_j - Q_j \cdot y'_j)}$$

where $X = (x_1 + x_2 + x_3 + x_4)/4$ is the center-of-mass or the macroscopic spacetime coordinates and x'_j and y'_j are the microscopic spacetime coordinates. Similar decomposition can be made for the second exponential factor in (3.37). As setted up in Section I, $\mathcal{W}_4$ in (3.37) depends weakly on X^μ , which means that, for $|p_1^0 + p_2^0 - q_1^0 - q_2^0| \gtrsim 1/L^0$ and/or $|\mathbf{p}_1 + \mathbf{p}_2 - \mathbf{q}_1 - \mathbf{q}_2| \gtrsim 1/L$,

$$\mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} \simeq 0. \quad (3.39)$$

Thus, the momentum conservation holds approximately in a single spacetime cell. Note that

$$\mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} = O(1) \quad \text{for} \quad \left| \sum_{j=1}^2 (p_j^0 - q_j^0) \right| \lesssim 1/L^0 \quad \text{and} \quad \left| \sum_{j=1}^2 (\mathbf{p}_j - \mathbf{q}_j) \right| \lesssim 1/L. \quad (3.40)$$

Now we take the limit $V \rightarrow \infty$ and go to the momentum space:

$$\begin{aligned} G_4 \equiv & -i \int dx_1 dx_2 dy_1 dy_2 e^{i \sum_{j=1}^2 (P_j \cdot x_j - Q_j \cdot y_j)} \mathcal{W}_4(x_1, x_2; y_1, y_2) \\ = & \frac{(2\pi)^4}{4} \frac{V^2}{\sqrt{E_{p_1} E_{p_2} E_{q_1} E_{q_2}}} \left[\mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} \prod_{j=1}^2 (\delta(p_{j0} - E_{p_j}) \delta(q_{j0} - E_{q_j})) \right. \\ & \left. + \mathcal{C}_{-\mathbf{p}_1 - \mathbf{p}_2; -\mathbf{q}_1 - \mathbf{q}_2} \prod_{j=1}^2 (\delta(p_{j0} + E_{p_j}) \delta(q_{j0} + E_{q_j})) \right]. \end{aligned} \quad (3.41)$$

A transition probability or a rate of a microscopic reaction is related to an on-shell amplitude. (In the case of equilibrium thermal field theories, this relation is settled in [21].) Let us now analyze the on-shell ($p_0^2 = E_p^2$) amplitudes. As mentioned above and can be seen from (3.41), G_4 *per se* does not contribute to the on-shell amplitudes. Then, $\mathcal{W}_4$ or G_4 appears as a part(s) of an on-shell amplitude. Let us see the structure of a connected amplitude S_4 , which includes a single G_4 . We may write, with obvious notation,

$$S_4 = (2\pi)^4 \int \left[\prod_{j=1}^2 \frac{d^4 P_j}{(2\pi)^4} \frac{d^4 Q_j}{(2\pi)^4} \right] G_4 H \delta^4 \left(\sum_l P_l + \sum_{j=1}^2 (Q_j - P_j) \right). \quad (3.42)$$

Here $\sum_l P_l + \sum_{j=1}^2 (Q_j - P_j)$ is the sum of the momenta that flow in H . As mentioned above $P_1 + P_2 \simeq Q_1 + Q_2$, we may replace the delta function in (3.42) with $\delta^4(\sum_l P_l)$:

$$\begin{aligned} S_4 \simeq & (2\pi)^4 \delta^4(\sum_l P_l) \tilde{S}_4, \\ \tilde{S}_4 = & \int \left[\prod_{j=1}^2 \frac{d^4 P_j}{(2\pi)^4} \frac{d^4 Q_j}{(2\pi)^4} \right] G_4 H. \end{aligned} \quad (3.43)$$

Let G'_4 be the Fourier-transformed $2n$ -point Green function computed in perturbative scheme constructed below, which is the counterpart of G_4 . Note that G'_4 , being of $O(\lambda)$ with λ the coupling constant, takes the form

$$G'_4 = (2\pi)^4 \delta^4(P_1 + P_2 - Q_1 - Q_2) \tilde{G}'_4.$$

The contribution (3.42) with G'_4 for G_4 reads

$$\begin{aligned} S'_4 &= (2\pi)^4 \delta^4 \left(\sum_l P_l \right) \tilde{S}'_4, \\ \tilde{S}'_4 &= \int \frac{d^4 P_1}{(2\pi)^4} \frac{d^4 P_2}{(2\pi)^4} \frac{d^4 Q_1}{(2\pi)^4} \tilde{G}'_4 H. \end{aligned} \quad (3.44)$$

Here we recall that the volume of the box, $V = L^3$, can be regarded as large. Let us see the dependence of (3.43) and (3.44) on (the large) V . $\tilde{G}'_4$ and then $\tilde{S}'_4$ are independent of V . Owing to the properties (3.39) and (3.40), integration over $\mathbf{q}_2$ in (3.43) yields a factor of $1/V$. Then, at this stage, $\tilde{S}_4$ is $O(V)$ larger than $\tilde{S}'_4$. Let us consider possible properties of $\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle$.

P1) *Strong factorizability*. There exist scale parameters $\vec{\delta}_2$ ($|\vec{\delta}_2| > 1/L$), so that, for $|\mathbf{p}_1 - \mathbf{p}_2| \gtrsim |\vec{\delta}_2|$ or $|\mathbf{q}_1 - \mathbf{q}_2| \gtrsim |\vec{\delta}_2|$,

$$\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle \simeq \langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{p}_1} \rangle \langle a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_2} \rangle + \langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{p}_2} \rangle \langle a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} \rangle, \quad (3.45)$$

which leads to $\mathcal{C}_{\mathbf{q}_1 \mathbf{q}_2; \mathbf{p}_1 \mathbf{p}_2} \simeq 0$.

P1') *Weak factorizability*. There exist scale parameters $\vec{\delta}_2$ ($|\vec{\delta}_2| > 1/L$), so that, for $|\mathbf{p}_1 - \mathbf{p}_2| \gtrsim |\vec{\delta}_2|$ and $|\mathbf{q}_1 - \mathbf{q}_2| \gtrsim |\vec{\delta}_2|$, (3.45) holds.

P2) *Strong "short-range correlation"*. There exist scale parameters $\vec{\delta}_3$ ($|\vec{\delta}_3| > 1/L$), so that, for $|\mathbf{p}_1 - \mathbf{q}_1| \gtrsim |\vec{\delta}_3|$ or $|\mathbf{p}_1 - \mathbf{q}_2| \gtrsim |\vec{\delta}_3|$,

$$\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle \simeq 0. \quad (3.46)$$

P2') *Weak "short-range correlation"*. There exist scale parameters $\vec{\delta}_3$ ($|\vec{\delta}_3| > 1/L$), so that, for $|\mathbf{p}_1 - \mathbf{q}_1| \gtrsim |\vec{\delta}_3|$ and $|\mathbf{p}_1 - \mathbf{q}_2| \gtrsim |\vec{\delta}_3|$, (3.46) holds. From (3.39) and (3.40), this condition implies $|\mathbf{p}_2 - \mathbf{q}_1| \gtrsim |\vec{\delta}_3|$ and $|\mathbf{p}_2 - \mathbf{q}_2| \gtrsim |\vec{\delta}_3|$.

Incidentally, P1) and P2) are not compatible with each other. In fact, when $\mathbf{p}_1 \simeq \mathbf{q}_1$ and $|\mathbf{p}_1 - \mathbf{q}_2| \gg \text{Max}(|\vec{\delta}_2|, |\vec{\delta}_3|)$, P2) yields $\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle \simeq 0$. On the other hand, P1) or P1') leads to (3.45). Owing to (3.34) with (3.26) and (3.27), the second term on the right-hand side (RHS) of (3.45) vanishes but the first term does not, which contradicts $\langle a_{\mathbf{q}_1}^\dagger a_{\mathbf{q}_2}^\dagger a_{\mathbf{p}_1} a_{\mathbf{p}_2} \rangle \simeq 0$ above.

Let us enumerate the cases, where $\tilde{S}_4$ and then $\mathcal{W}_4$ may possibly be ignored.

Property P1):

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O \left(\frac{|\vec{\delta}_2|^6 V}{\lambda \mathcal{P}^3} \right). \quad (3.47)$$

Property P2):

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O \left(\frac{|\vec{\delta}_3|^6 V}{\lambda \mathcal{P}^3} \right). \quad (3.48)$$

Properties P1') and P2):

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O \left(\frac{|\vec{\delta}_3|^3 \text{Min}(|\vec{\delta}_2|^3, |\vec{\delta}_3|^3) V}{\lambda \mathcal{P}^3} \right). \quad (3.49)$$

Properties P1) and P2'):

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O\left(\frac{|\vec{\delta}_2|^3 \text{Min}(|\vec{\delta}_2|^3, |\vec{\delta}_3|^3) V}{\lambda \mathcal{P}^3}\right). \quad (3.50)$$

Properties P1') and P2'):

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O\left(\frac{|\vec{\delta}_2|^3 |\vec{\delta}_3|^3 V}{\lambda \mathcal{P}^3}\right). \quad (3.51)$$

Let us inspect (3.47). If $|\vec{\delta}_2|^6 < O(\lambda \mathcal{P}^3/V)$ or $|\vec{\delta}_2| < O(\lambda^{1/6} \sqrt{\mathcal{P}/L})$, G_4 or $\mathcal{W}_4$ may be discarded. An exceptional case is where H in (3.42) is as large as $\propto \lambda/(|\vec{\delta}_2|^6 V)$ at $P_1 \simeq P_2 \simeq Q_1 \simeq Q_2$ with $P_1^2 = P_2^2 = Q_1^2 = Q_2^2 = m^2$. If $|\vec{\delta}_2| = O(1/L)$, (3.47) becomes

$$\frac{\tilde{S}_4}{\tilde{S}'_4} = O\left(\frac{1}{\lambda V \mathcal{P}^3}\right), \quad (3.52)$$

and then $\mathcal{W}_4$ may generally be discarded. Similar inspection may be made for (3.48) - (3.51).

Let us analyze the special case where the density matrix ρ is diagonal in Fock space:

$$\begin{aligned} \langle n_{\mathbf{k}}^{(a)} | \rho | n_{\mathbf{k}'}^{(a)} \rangle &= \delta_{\mathbf{k}\mathbf{k}'} \rho_{n_{\mathbf{k}}^{(a)}}, & \langle n_{\mathbf{k}}^{(b)} | \rho | n_{\mathbf{k}'}^{(b)} \rangle &= \delta_{\mathbf{k}\mathbf{k}'} \rho_{n_{\mathbf{k}}^{(b)}}, \\ a_{\mathbf{k}}^\dagger a_{\mathbf{k}} | n_{\mathbf{k}}^{(a)} \rangle &= n_{\mathbf{k}}^{(a)} | n_{\mathbf{k}}^{(a)} \rangle, & b_{\mathbf{k}}^\dagger b_{\mathbf{k}} | n_{\mathbf{k}}^{(b)} \rangle &= n_{\mathbf{k}}^{(b)} | n_{\mathbf{k}}^{(b)} \rangle, \end{aligned}$$

where $n_{\mathbf{k}}^{(a)/(b)} = 0, 1, 2, \dots$. In this case, Green functions are translation invariant and independent of the center-of-mass or macroscopic spacetime coordinates (cf. above after (3.27)). Equation (3.37) becomes

$$\begin{aligned} \mathcal{W}_4 &= \sum_{\mathbf{p}} \frac{i}{(2E_p V)^2} [\langle a_{\mathbf{p}}^{\dagger 2} a_{\mathbf{p}}^2 \rangle - 2\langle a_{\mathbf{p}}^\dagger a_{\mathbf{p}} \rangle^2] \\ &\times \left[e^{-iP \cdot (x_1 + x_2 - y_1 - y_2)} + \text{c.c.} \right], \end{aligned}$$

where 'c.c.' stands for the complex conjugate and (3.41) becomes

$$G_4 = -i \int dx_1 dx_2 dy_1 dy_2 e^{i \sum_{j=1}^2 (P_j \cdot x_j - Q_j \cdot y_j)} \mathcal{W}_4 \quad (3.53)$$

$$\begin{aligned} &= (2\pi)^{12} \delta^4 \left(\sum_{j=1}^2 (P_j - Q_j) \right) \frac{1}{V} \frac{2\pi \delta(P_1^2 - m^2)}{2E_{p_1}} [\langle a_{\mathbf{p}_1}^{\dagger 2} a_{\mathbf{p}_1}^2 \rangle - 2\langle a_{\mathbf{p}_1}^\dagger a_{\mathbf{p}_1} \rangle^2] \\ &\times \delta^4(P_2 - P_1) \delta^4(Q_1 - P_1), \end{aligned} \quad (3.54)$$

where, for simplicity, we have assumed $N_{\mathbf{p}} = N_{-\mathbf{p}}$. This leads to

$$\frac{S_4}{\tilde{S}_4} = O\left(\frac{1}{\lambda V \mathcal{P}^3}\right), \quad (3.55)$$

which is of the same order of magnitude as (3.52). Thus, G_4 or $\mathcal{W}_4$ may generally be ignored. Incidentally, for $\mathcal{W}_{2n}$, we obtain, in place of (3.55), $S_{2n}/\tilde{S}_{2n} = O(1/(\lambda V \mathcal{P}^3)^{n-1})$.

In the sequel of this paper, we ignore throughout the multi-point correlations, $\mathcal{W}_{2n}$ ($n \geq 2$).

D. UV renormalizability

Here, a comment is made on the UV renormalizability. We have completed the construction of the building blocks of Feynman rules. The vertices are given by (3.4) and (3.5) while the propagator is given by (3.36). As discussed at

the end of Section IIIA, it should be stressed that, if the “ $2n$ -point propagator” ($n \geq 2$) $\mathcal{W}_{2n}$ cannot be discarded, we should include it into the rules. We assume that, as in equilibrium thermal field theories, $\mathcal{W}(X; P)$ in (3.36) and $\mathcal{W}_{2n}$ ($n \geq 2$) in (3.17) are “soft.” Here, by “soft” we mean that, in the UV limit $|P^\mu| \rightarrow \infty$, $\mathcal{W}(X; P)$ ($\in \hat{\Delta}(X; P)$) tends to zero more quickly than the vacuum part $\hat{\Delta}^{(0)}(P)$ and $\mathcal{W}_{2n}$ tends to zero more quickly than the (leading contribution to the) $2n$ -point function. Then, genuine UV divergences emerge only from the vacuum part $\hat{\Delta}^{(0)}(P)$ of $\hat{\Delta}(X; P)$. This means that Green functions are UV finite if the vacuum theory is renormalizable and has been renormalized, which is in fact the case for vacuum theory of complex-scalar fields. For more elaborate argument, we refer the reader to [3,22].

IV. QUASIPARTICLE REPRESENTATION

A. Introduction of quasiparticle

It is convenient to rewrite $\hat{\Delta}(x, y)$, Eq. (3.18) or (3.36), using the so-called “Retarded/Advanced basis” [23]:

$$\hat{\Delta}(x, y) = \hat{\Delta}_{RA}(x - y) - iF(x, y)\hat{A}_+, \quad (4.1)$$

$$\hat{\Delta}_{RA}(x - y) = \hat{\Delta}_{diag}(x - y) + (\Delta_R(x - y) - \Delta_A(x - y)) \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (4.2)$$

$$F(x, y) = i\mathcal{W}(x; y) + \int \frac{d^4 P}{(2\pi)^3} e^{-iP \cdot (x-y)} \theta(-p_0) \delta(P^2 - m^2). \quad (4.3)$$

Here

$$\hat{\Delta}_{diag}(x - y) = \text{diag}(\Delta_R(x - y), -\Delta_A(x - y)), \quad (4.4)$$

$$\Delta_{R(A)}(x - y) = \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \Delta_{R(A)}(P) \quad (4.5)$$

with

$$\Delta_{R(A)}(P) = \frac{1}{P^2 - m^2 \pm i\epsilon(p_0)\eta} \quad (4.6)$$

the retarded (advanced) Green function. We refer (4.1) to as the R/A representation. When compared to the $F\bar{F}$ representation (3.36), the R/A representation makes the subsequent expressions compact and convenient for handling. A cost to be paid is that the R/A representation generates less intuitive expressions.

In the case of equilibrium system (cf. (3.29) and (3.30)), we have

$$\begin{aligned} \hat{\Delta}_{equ}(x - y) &= \hat{\Delta}_{RA}(x - y) \\ &+ \int \frac{d^4 P}{(2\pi)^4} e^{-P \cdot (x-y)} \epsilon(p_0) [\theta(-p_0) + N(E_p)] [\Delta_R(P) - \Delta_A(P)] \hat{A}_+, \end{aligned}$$

where a chemical potential (being conjugate to the charge) has been assumed to vanish, so that $N_+(E_p) = N_-(E_p)$ ($\equiv N(E_p)$). We note that, in momentum space, $\hat{\Delta}_{equ}$ may be diagonalized as

$$\hat{\Delta}_{equ}(P) = \begin{pmatrix} 1 & f_{equ}(p_0, E_p) \\ 1 & 1 + f_{equ}(p_0, E_p) \end{pmatrix} \hat{\Delta}_{diag}(P) \begin{pmatrix} 1 + f_{equ}(p_0, E_p) & f_{equ}(p_0, E_p) \\ 1 & 1 \end{pmatrix}, \quad (4.7)$$

where

$$f_{equ}(p_0, E_p) = \epsilon(p_0) [\theta(-p_0) + N(E_p)],$$

In configuration space, (4.7) reads

$$\hat{\Delta}_{equ}(x - y) = \int d^3 u d^3 v \sum_{\tau=\pm} \hat{B}_L^{(\tau)}(\mathbf{x} - \mathbf{u}) \hat{\Delta}_{diag}^{(\tau)}(\mathbf{u} - \mathbf{v}; x_0 - y_0) \hat{B}_R^{(\tau)}(\mathbf{v} - \mathbf{y}), \quad (4.8)$$

where $\hat{\Delta}_{diag}^{(+)}$ [$\hat{\Delta}_{diag}^{(-)}$] stands for a positive [negative] frequency part of $\hat{\Delta}_{diag}$ and

$$\hat{B}_L^{(\tau)}(\mathbf{x}) = \begin{pmatrix} \delta(\mathbf{x}) & f_{equ}^{(\tau)}(\mathbf{x}) \\ \delta(\mathbf{x}) & \delta(\mathbf{x}) + f_{equ}^{(\tau)}(\mathbf{x}) \end{pmatrix}, \quad \hat{B}_R^{(\tau)}(\mathbf{x}) = \begin{pmatrix} \delta(\mathbf{x}) + f_{equ}^{(\tau)}(\mathbf{x}) & f_{equ}^{(\tau)}(\mathbf{x}) \\ \delta(\mathbf{x}) & \delta(\mathbf{x}) \end{pmatrix} \quad (4.9)$$

with

$$f_{equ}^{(\tau)}(\mathbf{x}) = \epsilon(\tau) \int \frac{d^3p}{(2\pi)^3} e^{i\mathbf{p}\cdot\mathbf{x}} [\theta(-\tau) + N(E_p)].$$

It is to be noted that the form for the matrices $\hat{B}_{L(R)}^{(\tau)}$, which leads to (4.7) with diagonal $\hat{\Delta}_{diag}$, is not unique and the choice here is the simplest one [4,15].

The representation (4.8) may be interpreted in terms of quasiparticle picture. In fact, (4.8) naturally leads us to introduce quasiparticle fields, $\hat{\phi}_{RA}$ and $\hat{\phi}_{RA}^\dagger$, through

$$\begin{aligned} \hat{\phi}(x) &= \int d^3u \sum_{\tau=\pm} \hat{B}_L^{(\tau)}(\mathbf{x} - \mathbf{u}) \hat{\phi}_{RA}^{(\tau)}(\mathbf{u}, x_0) \\ \hat{\phi}^\dagger(y) &= \int d^3v \sum_{\tau=\pm} \hat{\phi}_{RA}^{(\tau)\dagger}(\mathbf{v}, y_0) \hat{B}_L^{(\tau)}(\mathbf{v} - \mathbf{y}), \end{aligned} \quad (4.10)$$

where $\hat{\phi}_{RA}^{(+)}$ [$\hat{\phi}_{RA}^{(-)\dagger}$] denotes positive frequency part of $\hat{\phi}_{RA}$ [$\hat{\phi}_{RA}^\dagger$] and $\hat{\phi}_{RA}^{(-)}$ [$\hat{\phi}_{RA}^{(+)\dagger}$] denotes negative frequency part. The transformation (4.10) is called thermal Bogoliubov transformation. The statistical average of $\hat{T}\hat{\phi}_{RA}\hat{\phi}_{RA}^\dagger$ (cf. (2.6)) assumes the form

$$\langle \hat{T}\hat{\phi}_{RA}(x) \hat{\phi}_{RA}^\dagger(y) \rangle = i\hat{\Delta}_{diag}(x - y). \quad (4.11)$$

Thus the fields $\hat{\phi}_{RA}$ and $\hat{\phi}_{RA}^\dagger$ describe well-defined propagating modes in the system, which we call retarded/advanced quasiparticles. It is to be noted that $\hat{\phi}_{RA}^\dagger$ is not a hermitian conjugate of $\hat{\phi}_{RA}$, which is a characteristic feature of the theory of this type [4,15].

With (4.10) and (4.11), we can readily show that $\hat{\Delta}_{equ}(x - y)$ turns out to (4.8). From (4.9), we see that

$$\int d^3u \hat{B}_L^{(\tau)}(\mathbf{x} - \mathbf{u}) \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u} - \mathbf{y}) = \hat{\tau}_3 \delta(\mathbf{x} - \mathbf{y}), \quad (4.12)$$

which shows that, under the transformation (4.10), the canonical commutation relation is preserved:

$$\begin{aligned} \left[\hat{\phi}(x), \dot{\hat{\phi}}^\dagger(y) \right] \delta(x_0 - y_0) &= \left[\hat{\phi}_{RA}(x), \dot{\hat{\phi}}_{RA}^\dagger(y) \right] \delta(x_0 - y_0) \\ &= i\hat{\tau}_3 \delta^4(x - y), \end{aligned} \quad (4.13)$$

where $\dot{\hat{\phi}}^\dagger(y) \equiv \partial \hat{\phi}^\dagger(y) / \partial y_0$. Furthermore, the nonperturbative or bilinear part of the hat-Lagrangian density is form invariant,

$$\begin{aligned} \hat{\mathcal{L}}_0 &= -\phi_1^\dagger (\partial^2 + m^2) \phi_1 + \phi_2^\dagger (\partial^2 + m^2) \phi_2 \\ &= -\dot{\hat{\phi}}^\dagger \hat{\tau}_3 (\partial^2 + m^2) \hat{\phi} \\ &= -\dot{\hat{\phi}}_{RA}^\dagger \hat{\tau}_3 (\partial^2 + m^2) \hat{\phi}_{RA}. \end{aligned} \quad (4.14)$$

Let us turn back to the out-of-equilibrium case. As a generalization of (4.10), we assume that the fields, $\hat{\phi}$ and $\hat{\phi}^\dagger$, are related to the retarded/advanced quasiparticle fields, $\hat{\phi}_{RA}$ and $\hat{\phi}_{RA}^\dagger$, through the transformation,

$$\begin{aligned} \hat{\phi}(x) &= \int d^3u \sum_{\tau=\pm} \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \hat{\phi}_{RA}^{(\tau)}(\mathbf{u}, x_0) \\ \hat{\phi}^\dagger(y) &= \int d^3v \sum_{\tau=\pm} \hat{\phi}_{RA}^{(\tau)\dagger}(\mathbf{v}, y_0) \hat{B}_L^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0), \end{aligned} \quad (4.15)$$

where

$$\begin{aligned}\hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) &= \begin{pmatrix} \delta(\mathbf{x} - \mathbf{u}) & f^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \\ \delta(\mathbf{x} - \mathbf{u}) & \delta(\mathbf{x} - \mathbf{u}) + f^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \end{pmatrix}, \\ \hat{B}_R^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0) &= \begin{pmatrix} \delta(\mathbf{v} - \mathbf{y}) + f^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0) & f^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0) \\ \delta(\mathbf{v} - \mathbf{y}) & \delta(\mathbf{v} - \mathbf{y}) \end{pmatrix}.\end{aligned}\quad (4.16)$$

We assume that $\hat{B}_{L(R)}^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0)$ depends on $(\mathbf{x} + \mathbf{u})/2$ and on x_0 only weakly. Corresponding to (4.12), we have

$$\int d^3u \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u}, \mathbf{y}; x_0) = \hat{\tau}_3 \delta(\mathbf{x} - \mathbf{y}), \quad (4.17)$$

so that, as in (4.13), the transformation (4.15) preserves the canonical commutation relation.

Incidentally, the so-called nonequilibrium thermo field dynamics [4,15] is formulated by taking (4.15) as a starting hypothesis.

B. Propagator

Using (4.15) and (4.11), we obtain for the propagator $\hat{\Delta}$,

$$\begin{aligned}\hat{\Delta}(x, y) &= -i \langle \hat{T} \hat{\phi}(x) \hat{\phi}^\dagger(y) \rangle \\ &= -i \int d^3u d^3v \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \langle \hat{T} \hat{\phi}_{RA}^{(\tau)}(\mathbf{u}, x_0) \hat{\phi}_{RA}^{(\tau)\dagger}(\mathbf{v}; y_0) \rangle \hat{B}_R^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0) \\ &= \int d^3u d^3v \sum_{\tau=\pm} \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \hat{\Delta}_{diag}^{(\tau)}(\mathbf{u} - \mathbf{v}; x_0 - y_0) \hat{B}_R^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0),\end{aligned}\quad (4.18)$$

which leads to (4.1) with

$$\begin{aligned}F(x, y) &= \sum_{\tau=\pm} F^{(\tau)}(x, y) \\ &= i \int d^3u \sum_{\tau=\pm} \left[\Delta_R^{(\tau)}(\mathbf{x} - \mathbf{u}; x_0 - y_0) f^{(\tau)}(\mathbf{u}, \mathbf{y}; y_0) \right. \\ &\quad \left. - f^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \Delta_A^{(\tau)}(\mathbf{u} - \mathbf{y}; x_0 - y_0) \right] \\ &= i \mathcal{W}(x; y) + \int \frac{d^4P}{(2\pi)^3} e^{-iP \cdot (x-y)} \theta(-p_0) \delta(P^2 - m^2).\end{aligned}\quad (4.19)$$

The last line is the definition of $F(x, y)$, Eq. (4.3). Equation (4.19) is a determining equation for $f^{(\tau)}$.

Straightforward manipulation using (1.3) in (4.18) yields

$$\begin{aligned}\hat{\Delta}(X; P) &\equiv \int d^4(x - y) e^{iP \cdot (x-y)} \hat{\Delta}(x, y) \\ &\simeq \hat{\Delta}^{(0)}(X; P) + \hat{\Delta}^{(1)}(X; P) \quad (X \equiv (x + y)/2), \\ \hat{\Delta}^{(0)}(X; P) &= \hat{\Delta}_{RA}(P) + \hat{\Delta}^{(0)'}(X; P),\end{aligned}\quad (4.20)$$

where

$$\hat{\Delta}_{RA}(P) = \begin{pmatrix} \Delta_R(P) & 0 \\ \Delta_R(P) - \Delta_A(P) & -\Delta_A(P) \end{pmatrix} \quad (4.21)$$

$$\begin{aligned}\hat{\Delta}^{(0)'}(X; P) &= -2\pi i \epsilon(p_0) f^{(\epsilon(p_0))}(X; \mathbf{p}) \delta(P^2 - m^2) \hat{A}_+ \\ \hat{\Delta}^{(1)}(X; P) &= \frac{i}{2} \frac{\partial f^{(\epsilon(p_0))}(X; \mathbf{p})}{\partial X^\mu} \frac{\partial}{\partial P_\mu} (\Delta_R(P) + \Delta_A(P)) \hat{A}_+ \\ &= -iP \cdot \partial_X f^{(\epsilon(p_0))}(X; \mathbf{p}) [\Delta_R^2(P) + \Delta_A^2(P)] \hat{A}_+.\end{aligned}\quad (4.22)$$

$$= -iP \cdot \partial_X f^{(\epsilon(p_0))}(X; \mathbf{p}) [\Delta_R^2(P) + \Delta_A^2(P)] \hat{A}_+.\quad (4.23)$$

This is not the end of the story. We should impose the condition (3.19), which leads to a constraint on $f^{(\tau)}(X; P)$. The most convenient way is to start from (3.21) with (4.19),

$$(\partial_x^2 + m^2)F^{(\tau)}(x, y) = (\partial_y^2 + m^2)F^{(\tau)}(x, y) = 0. \quad (4.24)$$

However, for the purpose of later use, we analyze (3.19) directly. From (4.18), we obtain

$$\begin{aligned} (\partial_x^2 + m^2)\hat{\Delta}(x, y) &= (\partial_x^2 + m^2) \int d^3u d^3v \sum_{\tau=\pm} \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \\ &\quad \times \hat{\Delta}_{diag}^{(\tau)}(\mathbf{u} - \mathbf{v}; x_0 - y_0) \hat{B}_R^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0) \\ &\simeq -\hat{\tau}_3 \delta^4(x - y) + \int d^3u d^3v \sum_{\tau=\pm} \left[2 \frac{\partial \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0)}{\partial x_0} \frac{\partial}{\partial x_0} \right. \\ &\quad \left. - \left\{ (\nabla_{\mathbf{x}}^2 - \nabla_{\mathbf{u}}^2) \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \right\} \right] \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u}, \mathbf{v}; x_0) \hat{\tau}_3 \hat{\Delta}^{(\tau)}(\mathbf{v}, \mathbf{y}; x_0, y_0), \end{aligned} \quad (4.25)$$

where $\hat{\Delta}^{(+)}$ [$\hat{\Delta}^{(-)}$] is a positive [negative] frequency part of $\hat{\Delta}$. Comparing this result with (3.19), we see that the second term should vanish, which reads, in the momentum space,

$$-2i \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x-y)} \sum_{\tau=\pm} \theta(\tau p_0) \left[P \cdot \partial_X f^{(\tau)}(X; \mathbf{p}) \right] \hat{\Delta}^{(\tau)}(X; P) = 0, \quad (4.26)$$

where $X = (x + y)/2$. Since p_0 and p are mutually independent, we obtain

$$\partial_{X_0} f^{(\tau)}(X; \mathbf{p}) = \mathbf{p} \cdot \nabla_{\mathbf{x}} f^{(\tau)}(X; \mathbf{p}) = 0 \quad (\tau = \pm), \quad (4.27)$$

as it should be, since, interaction is absent here. Thus $f^{(\tau)}(X; \mathbf{p})$ is X_0 independent and has vanishing gradient along the “flow lines.” It can easily be shown that (4.27) guarantees also $(\partial_y^2 + m^2)F^{(\tau)} = 0$, Eq. (4.24). Thanks to the relation (4.27), $\hat{\Delta}^{(1)}$ in (4.23) vanishes, and then

$$\begin{aligned} \hat{\Delta}(X; P) &\simeq \hat{\Delta}_{RA}(P) - 2\pi i \epsilon(p_0) f^{(\epsilon(p_0))}(X; \mathbf{p}) \delta(P^2 - m^2) \hat{A}_+ \\ &\simeq \hat{\Delta}_{RA}(P) + [\mathcal{W}(P; X) - 2\pi i \theta(-p_0) \delta(P^2 - m^2)] \hat{A}_+. \end{aligned} \quad (4.28)$$

$|p_0|$ prescription

Comparison between (4.28) and (3.33) yields

$$\begin{aligned} f^{(+)}(X; \mathbf{p}) &= N_+(X; E_p, \hat{\mathbf{p}}) \\ f^{(-)}(X; \mathbf{p}) &= -1 - N_-(X; E_p, -\hat{\mathbf{p}}), \end{aligned} \quad (4.29)$$

where N_+ (N_-) is the particle- (antiparticle)-number density function.

In the case of equilibrium system with vanishing chemical potential (cf. (3.30)),

$$\begin{aligned} f_{equ}^{(+)}(E_p) &= -1 - f_{equ}^{(-)}(E_p) \\ &= \frac{1}{e^{E_p/T} - 1}. \end{aligned}$$

$f_{equ}^{(\tau)}(E_p)$ always appears in combination with $\delta(P^2 - m^2) = \delta(p_0^2 - E_p^2)$. Then, at first sight, it seems that no difference arises between $f_{equ}^{(\tau)}(|p_0|)$ and $f_{equ}^{(\tau)}(E_p)$. This is, however, not the case, since, in general, $\hat{\Delta}(P)$ is to be multiplied by the functions that are singular at $|p_0| = E_p$. The correct choice has been known [24] for some time now to be $f_{equ}^{(\tau)}(|p_0|)$ — the “ $|p_0|$ prescription.”

Let us turn back to the present out-of-equilibrium case. On the basis of the above observation, we assume that the $|p_0|$ prescription should be adopted:

$$\begin{aligned}
f^{(+)}(X; \mathbf{p}) &= N_+(X; E_p, \hat{\mathbf{p}}) \rightarrow N_+(X; p_0, \hat{\mathbf{p}}) \equiv f^{(+)}(X; P) \\
f^{(-)}(X; \mathbf{p}) &= -1 - N_-(X; E_p, -\hat{\mathbf{p}}) \rightarrow -1 - N_-(X; -p_0, -\hat{\mathbf{p}}) \equiv f^{(-)}(X; P), \\
f(X; P) &= \theta(p_0) f^{(+)}(X; P) + \theta(-p_0) f^{(-)}(X; P).
\end{aligned} \tag{4.30}$$

In what follows, we simply refer f to as *the number density (function)*. An argument for the necessity of adopting the $|p_0|$ prescription is given in Appendix A.

It is to be noted that the translation into the $|p_0|$ prescription is *formally* achieved by replacing (4.18) and (4.16) with, in respective order,

$$\begin{aligned}
\hat{\Delta}(x, y) &= \int d^4u d^4v \hat{B}_L(x, u) \hat{\Delta}_{diag}(u - v) \hat{B}_R(v, y) \\
&\equiv [\hat{B}_L \cdot \hat{\Delta}_{diag} \cdot \hat{B}_R](x, y),
\end{aligned} \tag{4.31}$$

and

$$\begin{aligned}
\hat{B}_L(x, u) &= \begin{pmatrix} \delta^4(x - u) & f(x, u) \\ \delta^4(x - u) & \delta^4(x - u) + f(x, u) \end{pmatrix}, \\
\hat{B}_R(v, y) &= \begin{pmatrix} \delta^4(v - y) + f(v, y) & f(v, y) \\ \delta^4(v - y) & \delta^4(v - y) \end{pmatrix}.
\end{aligned} \tag{4.32}$$

The function $f(x, y)$ here is an inverse Fourier transform of the number-density function $f(X; P)$ in (4.30):

$$f(X; P) = \int d^4(x - y) e^{iP \cdot (x - y)} f(x, y) \quad (X = (x + y)/2).$$

Now (4.26) turns out to

$$-2i \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x - y)} [P \cdot \partial_X f(X; P)] \hat{\Delta}(X; P) = 0,$$

and then, in place of (4.27), we have

$$P \cdot \partial_X f(X; P) = 0. \tag{4.33}$$

This is a continuity equation for f along the “flow line” in a four-dimensional space. Equation (4.33) may be solved as

$$f(X; P) = \mathcal{F}(\mathbf{X} - (X_0 - T_{in})\mathbf{p}/p_0; P), \tag{4.34}$$

where $\mathcal{F}$ is an arbitrary function and T_{in} is an (arbitrary) “initial” time. Given the initial data $f(X_0 = T_{in}, \mathbf{X}; P)$ that characterizes the ensemble of the systems at $X_0 = T_{in}$, (4.34) fixes the form of $\mathcal{F}$ and we obtain

$$f(X; P) = f(X_0 = T_{in}, \mathbf{X} - (X_0 - T_{in})\mathbf{p}/p_0; P).$$

The propagator $\hat{\Delta}(X; P)$ with this $f(X; P)$ still takes the form (4.28):

$$\hat{\Delta}(X; P) = \hat{\Delta}_{RA}(P) + f(X; P) (\Delta_R(P) - \Delta_A(P)) \hat{A}_+. \tag{4.35}$$

In order to see the physical meaning of f to be the “number density,” let us compute a statistical average of the current density:

$$\begin{aligned}
\langle j^\mu(x) \rangle &\equiv \frac{i}{2} \left[\langle \phi^\dagger(x) \overleftrightarrow{\partial}^\mu \phi(x) \rangle - \langle \phi(x) \overleftrightarrow{\partial}^\mu \phi^\dagger(x) \rangle \right] \\
&= \frac{1}{2} (\partial_{x_\mu} - \partial_{y_\mu}) [\Delta_{12}(y, x) + \Delta_{21}(y, x)]_{y=x} \\
&= i \int \frac{d^4P}{(2\pi)^4} P^\mu [\Delta_{12}(x; P) + \Delta_{21}(x; P)],
\end{aligned} \tag{4.36}$$

where $\overleftrightarrow{\partial} \equiv \overrightarrow{\partial} - \overleftarrow{\partial}$. Straightforward manipulation using (4.35) with (4.30) yields

$$\begin{aligned}\langle j^0(x) \rangle &\simeq \int \frac{d^3p}{(2\pi)^3} [\tilde{N}_+(x; \mathbf{p}) - \tilde{N}_-(x; -\mathbf{p})] \\ \langle \mathbf{j}(x) \rangle &\simeq \int \frac{d^3p}{(2\pi)^3} \frac{\mathbf{p}}{E_p} [\tilde{N}_+(x; \mathbf{p}) + \tilde{N}_-(x; -\mathbf{p})],\end{aligned}\quad (4.37)$$

where

$$\begin{aligned}\tilde{N}_+(x; \mathbf{p}) &\equiv N_+(x; P) \Big|_{p_0 = E_p}, \\ \tilde{N}_-(x; -\mathbf{p}) &\equiv N_-(x; -P) \Big|_{p_0 = -E_p}.\end{aligned}\quad (4.38)$$

Thus $\tilde{N}_+(x; \mathbf{p})$ [$\tilde{N}_-(x; -\mathbf{p})$] is a number density of particles with momentum $\mathbf{p}$ [antiparticles with momentum $-\mathbf{p}$]. It is to be noted that (4.38) is valid in the gradient approximation, i.e., the terms with first derivative $\partial N_{\pm}(x; \pm P)/\partial x$ do not appear in (4.38). Also to be noted is that the argument x here is a macroscopic spacetime coordinates.

C. Two perturbative schemes

In this subsection, we introduce two perturbative schemes of calculating various quantities. The first scheme, which we call the “bare- f scheme,” is the naive one introduced above. In this scheme, perturbative calculation goes with (3.4) and (3.5) for vertices and (4.35) for propagators. We designate the number-density function in (4.35) in this scheme as f_B .

The second scheme, which we call the “physical- f scheme,” is defined as follows. We *redefine* the fields, $\hat{\phi}$ and $\hat{\phi}^\dagger$, by (4.15) with (4.16), where $f^{(\tau)}$ is different from $f_B^{(\tau)}$. Then, the fields in this scheme are different from the fields in the bare- f scheme. Changing the definition of fields leads to emergence of a counter term in the Lagrangian. In order to extract it, we first note that (4.33) does not hold now and (4.25) may be written as

$$\begin{aligned}\int d^3u d^3v \sum_{\tau=\pm} D_\tau(x; \mathbf{u}, \mathbf{v}) \hat{\Delta}^{(\tau)}(\mathbf{v}, \mathbf{y}; x_0, y_0) &\simeq -\hat{\tau}_3 \delta^4(x - y), \\ D_\tau(x; \mathbf{u}, \mathbf{v}) &\equiv (\partial_x^2 + m^2) \delta(\mathbf{x} - \mathbf{u}) \delta(\mathbf{u} - \mathbf{v}) - \left[2 \frac{\partial \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0)}{\partial x_0} \frac{\partial}{\partial x_0} \right. \\ &\quad \left. - \left\{ (\nabla_{\mathbf{x}}^2 - \nabla_{\mathbf{u}}^2) \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \right\} \right] \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u}, \mathbf{v}; x_0) \hat{\tau}_3.\end{aligned}\quad (4.39)$$

This means that, as a free hat-Lagrangian density, we should take

$$\hat{\mathcal{L}}_0 = - \int d^3u d^3v \sum_{\tau=\pm} \hat{\phi}^{(\tau)\dagger}(\mathbf{x}; x_0) \hat{\tau}_3 D_\tau(x; \mathbf{u}, \mathbf{v}) \hat{\phi}^{(\tau)}(\mathbf{v}; x_0), \quad (4.40)$$

where $\hat{\phi}^{(\tau)\dagger} = (\phi_1^{(\tau)\dagger}, \phi_2^{(\tau)\dagger})$ and $\hat{\phi}^{(\tau)} = {}^t(\phi_1^{(\tau)}, \phi_2^{(\tau)})$ with $\phi_i^{(+)}$ ($\phi_i^{(-)}$) a positive (negative) frequency part of ϕ_i , and $\phi_i^{(\pm)\dagger}$ a Hermitian conjugate of $\phi_i^{(\pm)}$. Instead of (4.25), starting with $(\partial_y^2 + m^2) \hat{\Delta}(x, y)$, we are led to the same $\hat{\mathcal{L}}_0$ as (4.40) above, as it should be. Equation (4.40) tells us that, for compensating the difference between (4.40) and the original free hat-Lagrangian density, the counter term should be introduced in the hat-Lagrangian density:

$$\begin{aligned}\hat{\mathcal{L}}_c &= - \int d^3u d^3v \sum_{\tau=\pm} \hat{\phi}^{(\tau)\dagger}(\mathbf{x}; x_0) \hat{\tau}_3 \left[2 \frac{\partial \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0)}{\partial x_0} \frac{\partial}{\partial x_0} \right. \\ &\quad \left. - (\nabla_{\mathbf{x}}^2 - \nabla_{\mathbf{u}}^2) \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \right] \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u}, \mathbf{v}; x_0) \hat{\tau}_3 \hat{\phi}^{(\tau)}(\mathbf{v}; x_0),\end{aligned}\quad (4.41)$$

which leads to a two-point vertex function,

$$\begin{aligned}
i\hat{\mathcal{V}}_c(x, y) &= -i\hat{\tau}_3 \int d^3u \sum_{\tau=\pm} \left[2 \frac{\partial \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0)}{\partial x_0} \frac{\partial}{\partial x_0} - (\nabla_{\mathbf{x}}^2 - \nabla_{\mathbf{u}}^2) \hat{B}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \right] \\
&\quad \times \hat{\tau}_3 \hat{B}_R^{(\tau)}(\mathbf{u}, \mathbf{y}; x_0) \hat{\tau}_3 \delta(x_0 - y_0) \\
&\simeq i\hat{A}_- \sum_{\tau=\pm} \left[2 \frac{\partial f(\mathbf{x}, \mathbf{y}, x_0)}{\partial x_0} \frac{\partial}{\partial x_0} - (\nabla_{\mathbf{x}}^2 - \nabla_{\mathbf{y}}^2) f^{(\tau)}(\mathbf{x}, \mathbf{y}, x_0) \right] \delta(x_0 - y_0) \\
&= 2 \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x-y)} \sum_{\tau=\pm} \theta(\tau p_0) \left[P \cdot \partial_X f^{(\tau)}(X; \mathbf{p}) \right] \hat{A}_-.
\end{aligned}$$

Here $X = (x + y)/2$ and $\hat{A}_-$ is defined as in (3.16). Going to the $|p_0|$ prescription, we have (cf. (4.30)),

$$i\hat{\mathcal{V}}_c(x, y) \rightarrow 2 \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x-y)} [P \cdot \partial_X f(X; P)] \hat{A}_-. \quad (4.42)$$

The form for $f(X; P)$ will be determined later so as to be the physical “number density.” If we impose (4.33), (4.42) vanishes as it should be.

The “free form” of the propagator (4.28) should be replaced by (4.20) - (4.23) with the $|p_0|$ prescription:

$$\hat{\Delta}(x, y) = \int \frac{d^4P}{(2\pi)^4} e^{-iP \cdot (x-y)} \hat{\Delta}(X; P), \quad (4.43)$$

$$\hat{\Delta}(X; P) = \hat{\Delta}^{(0)}(X; P) + \hat{\Delta}^{(1)}(X; P) \quad (4.44)$$

$$\hat{\Delta}^{(0)}(X; P) = \hat{B}_L(X; P) \hat{\Delta}_{diag}(P) \hat{B}_R(X; P) = \hat{\Delta}_{RA}(P) + \hat{\Delta}^{(0)'}(X; P), \quad (4.45)$$

$$\hat{B}_L(X; P) = \begin{pmatrix} 1 & f(X; P) \\ 1 & 1 + f(X; P) \end{pmatrix}, \quad \hat{B}_R(X; P) = \begin{pmatrix} 1 + f(X; P) & f(X; P) \\ 1 & 1 \end{pmatrix}, \quad (4.46)$$

where $\hat{\Delta}_{RA}(P)$ is as in (4.21) and (cf. (4.30))

$$\hat{\Delta}^{(0)'}(X; P) = f(X; P) (\Delta_R(P) - \Delta_A(P)) \hat{A}_+, \quad (4.47)$$

$$\hat{\Delta}^{(1)}(X; P) = \frac{i}{2} \frac{\partial f(X; P)}{\partial X^\mu} \frac{\partial}{\partial P_\mu} (\Delta_R(P) + \Delta_A(P)) \hat{A}_+ \quad (4.48)$$

$$= -iP \cdot \partial_X f(X; P) [\Delta_R^2(P) + \Delta_A^2(P)] \hat{A}_+. \quad (4.49)$$

Let us discuss two perturbative schemes in analogy with mass renormalization in vacuum theory. We take the Lagrangian density

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_0 + \mathcal{L}_{int}, \\
\mathcal{L}_0 &= \partial_\mu \phi^\dagger \partial^\mu \phi - m_B^2 \phi^\dagger \phi.
\end{aligned}$$

The “bare scheme” in vacuum theory takes $\mathcal{L}_0$ as a nonperturbative part. Then, the propagator reads,

$$\Delta(P) = \frac{1}{P^2 - m_B^2 + i\eta},$$

which corresponds to (4.28) in the bare- f scheme here. The “renormalized scheme” in vacuum theory adopts (cf. (3.2) and (3.3))

$$\mathcal{L}'_0 = \partial_\mu \phi^\dagger \partial^\mu \phi - m^2 \phi^\dagger \phi, \quad (4.50)$$

as the nonperturbative part. Here $m = Z_m^{-1/2} m_B$ is a renormalized mass. Equation (4.50) leads to

$$\Delta(P) = \frac{1}{P^2 - m^2 + i\eta},$$

for the propagator, which corresponds to (4.43) - (4.49) in the physical- f scheme here. To compensate the difference $\mathcal{L}_0 - \mathcal{L}'_0$, a counter term should be introduced into the Lagrangian density, $\mathcal{L}_c = -(Z_m Z - 1) m^2 \phi^\dagger \phi$, which corresponds to (4.41) here.

To summarize, the bare- f scheme is constructed in terms of original bare number density $f_B(X; P)$ [respect. bare mass m_B]. On the other hand, the physical- f scheme is constructed in terms of physical number density $f(X; P)$ [respect. renormalized mass m]. Both perturbative schemes are equivalent. The first scheme starts with the “bare quantity” and the “renormalization” is done at the end, while in the second scheme, the “renormalization” is done at the beginning by introducing the counter hat-Lagrangian $\hat{\mathcal{L}}_c$ [respect. $\mathcal{L}_c$]. It is worth recalling that the renormalized mass m is defined so that m is free from UV divergences. However, there is arbitrariness in the definition of the “finite part” of m . m is determined by imposing some condition. (In this paper, we are adopting the $\overline{\text{MS}}$ scheme.) Thus, m is determined order by order in perturbation series. As will be seen in the next section, this also applies to the present physical- f scheme. Namely, there is arbitrariness in defining f . In Section VC, we shall impose the condition for determining f so as to be the physical number density, under which $P \cdot \partial_X f(X; P)$ in $\hat{\Delta}^{(1)}(X; P)$, Eq. (4.49), and then also $f(X; P)$ turn out to be determined order by order in perturbation series.

V. INTERACTING FIELD

Interactions among fields give rise to reactions taking place in the system, which, in turn, causes a nontrivial change in number density of quasiparticles. In this section, we analyze full propagator in two perturbative schemes presented at the last section and demonstrate their equivalence.

A. Self-energy part

The self-energy part $\hat{\Sigma}$ takes the form

$$\begin{aligned}\hat{\Sigma}(x, y) &= \hat{\Sigma}_{int}(x, y) + \hat{\Sigma}_c(x, y), \\ \hat{\Sigma}_c(x, y) &= -\hat{\mathcal{V}}_c(x, y) \equiv -\mathcal{V}_c(x, y)\hat{A}_-, \end{aligned} \quad (5.1)$$

where $\hat{\Sigma}_{int}$ is a loop-diagram contribution, $\hat{\Sigma}_c$ comes from the counter hat-Lagrangian density $\hat{\mathcal{L}}_c$ in (4.41), and $\hat{\mathcal{V}}_c$ is as in (4.42).

It can be shown that (cf. Appendix B)

$$\sum_{i, j=1}^2 \Sigma_{ij} = \sum_{i, j=1}^2 (\Sigma_{int})_{ij} = 0, \quad (5.2)$$

with which we obtain, in the configuration space,

$$\underline{\hat{\Sigma}} \equiv \hat{B}_R \cdot \hat{\Sigma} \cdot \hat{B}_L = \begin{pmatrix} \Sigma_R & \Sigma_{off} \\ 0 & -\Sigma_A \end{pmatrix}, \quad (5.3)$$

where $\hat{B}_{R(L)}$ is as in (4.32) and

$$\begin{aligned}\Sigma_R &= \Sigma_{11} + \Sigma_{12} = (\Sigma_{int})_{11} + (\Sigma_{int})_{12}, \\ \Sigma_A &= -(\Sigma_{22} + \Sigma_{12}) = -\{(\Sigma_{int})_{22} + (\Sigma_{int})_{12}\} (= \Sigma_R^*), \\ \Sigma_{off} &= \Sigma_{12} + \Sigma_{12} \cdot f - \Sigma_{21} \cdot f + \Sigma_A \cdot f - f \cdot \Sigma_A + \mathcal{V}. \end{aligned} \quad (5.4)$$

Σ_R and Σ_A are called the retarded and advanced self-energy parts, respectively.

Going to momentum space, we obtain, after some manipulations,

$$\Sigma_{off}(X; P) \simeq -i \{f(X; P), P^2 - m^2 - Re\Sigma_R(X; P)\} + i\tilde{\Gamma}^{(p)}(X; P), \quad (5.5)$$

where

$$\{A(X; P), B(X; P)\} \equiv \frac{\partial A(X; P)}{\partial X_\mu} \frac{\partial B(X; P)}{\partial P^\mu} - \frac{\partial A(X; P)}{\partial P^\mu} \frac{\partial B(X; P)}{\partial X_\mu}, \quad (5.6)$$

$$i\tilde{\Gamma}^{(p)}(X; P) \equiv (1 + f(X; P))\Sigma_{12}(X; P) - f(X; P)\Sigma_{21}(X; P). \quad (5.7)$$

In deriving (5.5), use has been made of the relation $\Sigma_A(X; P) = [\Sigma_R(X; P)]^*$, a proof of which is given in Appendix B.

B. Self-energy-part resummed propagator

In this subsection, we compute a self-energy-part resummed full propagator through the Dyson equation,

$$\hat{G}(x, y) = \hat{\Delta}(x, y) + [\hat{\Delta} \cdot \hat{\Sigma} \cdot \hat{G}](x, y).$$

We would like to obtain $\hat{G}(X; P)$ ($X = (x + y)/2$), the Fourier transform of $\hat{G}(x, y)$ on $x - y$. Using the transformation (4.31), we obtain

$$\begin{aligned} \underline{\hat{G}}(x, y) &= [\hat{B}_L^{-1} \cdot \hat{G} \cdot \hat{B}_R^{-1}](x, y) \\ &= \hat{\Delta}_{diag}(x, y) + [\hat{\Delta}_{diag} \cdot \hat{\Sigma} \cdot \underline{\hat{G}}](x, y). \end{aligned}$$

Fourier transforming on $x - y$ using (C2) in Appendix C, we obtain

$$\begin{aligned} \underline{\hat{G}}(X; P) &\simeq \hat{\Delta}_{diag}(P) + \hat{\Delta}_{diag}(P) \hat{\Sigma}(X; P) \underline{\hat{G}}(X; P) \\ &+ \frac{i}{2} \left[\frac{\partial \hat{\Delta}_{diag}}{\partial P^\mu} \frac{\partial \hat{\Sigma}}{\partial X_\mu} \underline{\hat{G}} - \hat{\Delta}_{diag} \frac{\partial \hat{\Sigma}}{\partial X_\mu} \frac{\partial \underline{\hat{G}}}{\partial P^\mu} + \frac{\partial \hat{\Delta}_{diag}}{\partial P^\mu} \underline{\hat{\Sigma}} \frac{\partial \underline{\hat{G}}}{\partial X_\mu} \right], \end{aligned} \quad (5.8)$$

where (cf. (5.3))

$$\begin{aligned} \hat{\Sigma} &= \hat{\Sigma}_{diag} + \hat{\Sigma}_{off} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \\ \hat{\Sigma}_{diag} &= \text{diag}(\Sigma_R, -\Sigma_A). \end{aligned} \quad (5.9)$$

The terms in the square brackets in (5.8) involve X^μ -derivatives and are small when compared to the first term on the RHS.

Let us solve (5.8) iteratively. In doing so, we assume that the leading terms on the RHS of (5.8) are $\hat{\Delta}_{diag} + \hat{\Delta}_{diag} \hat{\Sigma}_{diag} \underline{\hat{G}}$ and $\hat{\Sigma}$'s in the square brackets may be approximated as $\hat{\Sigma} \simeq \hat{\Sigma}_{diag}$. This assumption will be justified a posteriori (cf. the next subsection). Straightforward manipulation yields

$$\underline{\hat{G}}(X; P) = \hat{G}_{diag}(X; P) - \Sigma_{off}(X; P) G_R(X; P) G_A(X; P) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad (5.10)$$

where

$$\begin{aligned} \hat{G}_{diag}(X; P) &= \text{diag}(G_R(X; P), -G_A(X; P)), \\ G_{R(A)}(X; P) &= \frac{1}{P^2 - m^2 - \Sigma_{R(A)}(X; P) \pm i\epsilon(p_0)\eta}. \end{aligned} \quad (5.11)$$

It is to be noted that (5.10) is exact to the derivative approximation, i.e., no term including the first derivative (with respect to X^μ) arises. Then, using again (C2) in Appendix C, we obtain

$$\hat{G}(X; P) = \hat{G}^{(0)}(X; P) + \hat{G}^{(1)}(X; P), \quad (5.12)$$

$$\begin{aligned} \hat{G}^{(0)}(X; P) &= \hat{B}_L(X; P) \hat{G}_{diag}(X; P) \hat{B}_R(X; P) \\ &= \hat{G}_{RA}(X; P) + \hat{G}^{(0)'}(X; P), \end{aligned} \quad (5.13)$$

where

$$\hat{G}_{RA}(X; P) = \begin{pmatrix} G_R(X; P) & 0 \\ G_R(X; P) - G_A(X; P) & -G_A(X; P) \end{pmatrix}, \quad (5.14)$$

$$\hat{G}^{(0)'}(X; P) = f(X; P) (G_R(X; P) - G_A(X; P)) \hat{A}_+, \quad (5.15)$$

$$\begin{aligned} \hat{G}^{(1)}(X; P) &= -\Sigma_{off}(X; P) G_R(X; P) G_A(X; P) \hat{A}_+ \\ &\quad - \frac{i}{2} [\{f(X; P), P^2 - m^2 - \Sigma_R(X; P)\} G_R^2(X; P) \end{aligned} \quad (5.16)$$

$$\begin{aligned}
& + \{f(X; P), P^2 - m^2 - \Sigma_A(X; P)\} G_A^2(X; P) \hat{A}_+ \\
& = \frac{1}{2} \Sigma_{off}(X; P) (G_R(X; P) - G_A(X; P))^2 \hat{A}_+ \\
& \quad - \frac{1}{2} \{f(X; P), Im\Sigma_R(X; P)\} (G_R^2(X; P) - G_A^2(X; P)) \hat{A}_+ \\
& \quad - \frac{1}{2} \tilde{\Gamma}^{(p)}(X; P) (G_R^2(X; P) + G_A^2(X; P)) \hat{A}_+.
\end{aligned} \tag{5.17}$$

In narrow-width approximation, $Im\Sigma_R = -Im\Sigma_A \rightarrow -\epsilon(p_0)0^+$, the first term on the RHS of (5.17) develops pinch singularity, while the rest turns out to the well-defined distributions.

For clarifying the physical meaning of (5.17), we compute the contribution to the physical number density through analyzing the contribution to the statistical average of the current density $\langle j^\mu \rangle$, Eq. (4.36). The second and third terms on the RHS of (5.17) leads to perturbative corrections to $\langle j^\mu \rangle$ in (4.37) due to quantum and medium effects, while the first term yields a large contribution,

$$\langle \delta j^\mu(x) \rangle = i \int \frac{d^4 P}{(2\pi)^4} P^\mu \Sigma_{off}(x; P) (G_R(x; P) - G_A(x; P))^2, \tag{5.18}$$

which diverges in the narrow-width approximation. (Note again that the argument x here is a macroscopic spacetime coordinates.) In the next subsection, we inspect this large contribution more closely in the bare- f and physical- f schemes.

C. Two perturbative schemes revisited

1. Bare- f scheme

Let us estimate (5.18). All the formulae displayed above may be used with $f \rightarrow f_B$ and with imposition of (4.33). Important region of integration is where $Re(G_R^{-1}(x; P)) = P^2 - m^2 - Re\Sigma_R(x; P) \simeq 0$. We define “on the mass-shell” $p_0 = \pm\omega_\pm(X; \pm\mathbf{p})$ ($\equiv \pm\omega_\pm$) through

$$[P^2 - m^2 - Re\Sigma_R(X; P)] \Big|_{p_0 = \pm\omega_\pm} = 0. \tag{5.19}$$

We make following approximations;

$$\begin{aligned}
P^\mu \Sigma_{off} & \simeq \theta(p_0) (\omega_+, \mathbf{p})^\mu \Sigma_{off}^{(bare)(+)} + \theta(-p_0) (-\omega_-, \mathbf{p})^\mu \Sigma_{off}^{(bare)(-)}, \\
\frac{-i}{2} [G_R(x; P) - G_A(x; P)] & \simeq \sum_{\tau\pm} \frac{\theta(\tau p_0) Im\Sigma_R^{(\tau)}}{[2\omega_\tau Z_\tau^{-1}(p_0 - \tau\omega_\tau)]^2 + (Im\Sigma_R^{(\tau)})^2},
\end{aligned} \tag{5.20}$$

where

$$\Sigma_{off}^{(bare)(\pm)} = \Sigma_{off}(X; p_0 = \pm\omega_\pm, \mathbf{p}), \quad \Sigma_R^{(\pm)} = \Sigma_R(X; p_0 = \pm\omega_\pm, \mathbf{p}), \tag{5.21}$$

$$Z_\pm^{-1} \equiv 1 \mp \frac{\partial Re\Sigma_R}{\partial p_0} \Big|_{p_0 = \pm\omega_\pm}. \tag{5.22}$$

It is convenient to carry out the wave-function renormalization such that $Z_\pm = 1$, which we assume to be done for a while. Using (5.20) in (5.18), we obtain

$$\langle \delta j^\mu(x) \rangle \simeq \frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \left[(1, \mathbf{p}/\omega_+)^mu \frac{-i\Sigma_{off}^{(bare)(+)}}{|Im\Sigma_R^{(+)}|} - (1, -\mathbf{p}/\omega_-)^mu \frac{-i\Sigma_{off}^{(bare)(-)}}{|Im\Sigma_R^{(-)}|} \right]. \tag{5.23}$$

In deriving this, we have assumed that $|Im\Sigma_R^{(\pm)}| \ll \omega_\pm$, which is the case in equilibrium case. As in (4.36) - (4.38), we can read off from (5.23) the contribution to the physical number density:

$$\begin{aligned}
\delta \tilde{N}_\pm(x, \pm \mathbf{p}) &= \frac{1}{2} \frac{-i \Sigma_{off}^{(bare)(\pm)}}{|Im \Sigma_R^{(\pm)}|}, \\
&= \frac{1}{2} \tau_\pm \left[-i \Gamma_\pm^{(p)} + \frac{1}{2\omega_\pm} \{f, Re \Sigma_R\} \right]_{p_0 = \pm \omega_\pm}.
\end{aligned} \tag{5.24}$$

Here $\tau_+ \equiv 2\omega_+ / |Im \Sigma_R^{(+)}|$ [$\tau_- \equiv 2\omega_- / |Im \Sigma_R^{(-)}|$] is a characteristic time during which a particle with momentum $\mathbf{p}$ [an antiparticle with momentum $-\mathbf{p}$] damps:

$$\begin{aligned}
G_R(X; x_0 - y_0; \mathbf{p}) &= \int \frac{d^3 p}{(2\pi)^3} e^{i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} G_R(x, y) \\
&\propto e^{-(x_0 - y_0)/\tau_+} \quad \text{for a particle mode,} \\
&\propto e^{-(x_0 - y_0)/\tau_-} \quad \text{for an antiparticle mode.}
\end{aligned}$$

In (5.24),

$$\Gamma_\pm^{(p)} = \frac{-i}{2\omega_\pm} [(1 + N_\pm) \Sigma_{12(21)} - N_\pm \Sigma_{21(12)}] \Big|_{p_0 = \pm \omega_\pm}, \tag{5.25}$$

which comes from $\tilde{\Gamma}^{(p)} \in -i \Sigma_{off}$ in (5.7), is a net production rate of a particle (an antiparticle) of momentum $\mathbf{p}$ ($-\mathbf{p}$). In fact, $\Gamma_\pm^{(p)}$, Eq. (5.25), is a difference between the production rate and the decay rate, so that $\Gamma_\pm^{(p)}$ is the net production rate. In the case of equilibrium system, $\Gamma_\pm^{(p)} = 0$ (detailed balance formula).

To disclose the physical meaning of (5.24), for the time being, we put aside the second term in the square brackets, whose physical interpretation will become clear later at (5.32). Then, the RHS of (5.24) is the change in the physical number density, during the time interval $\tau_\pm/2$, due to the net production rate. In Section I, we have introduced spacetime cells, whose size is L^μ ($\mu = 0, 1, 2, 3$). It is quite natural to take $\tau_\pm = L_0$, where L_0 is the size of a time direction of the spacetime cell (including the spacetime point x^μ in (5.24)). [Strictly speaking, in general, $\tau_+ \neq \tau_-$. However, in the present crude argument, we ignore this difference.] It is interesting to note that (5.24) is *half* of the net production probability, due to the reaction, during the time interval $\tau_\pm = L_0$. [In this respect, cf. [25].]

Noticing that we are concerned about the particle mode with momentum $\mathbf{p}$ [antiparticle mode with momentum $-\mathbf{p}$], we see that (5.24) leads to the relations for the physical number density,

$$\begin{aligned}
\tilde{N}_+(x_0 + L_0, \mathbf{x} + L_0 \mathbf{p} / \omega_+; \mathbf{p}) - \tilde{N}_+(x_0, \mathbf{x}; \mathbf{p}) &\simeq L_0 \frac{-i \Sigma_{off}^{(bare)}(x; \omega_+, \mathbf{p})}{2\omega_+}, \\
\tilde{N}_-(x_0 + L_0, \mathbf{x} - L_0 \mathbf{p} / \omega_-; -\mathbf{p}) - \tilde{N}_-(x_0, \mathbf{x}; -\mathbf{p}) &\simeq L_0 \frac{-i \Sigma_{off}^{(bare)}(x; -\omega_-, \mathbf{p})}{2\omega_-},
\end{aligned}$$

from which we obtain

$$\frac{\partial \tilde{N}_\pm(x; \pm \mathbf{p})}{\partial x_0} \pm \frac{\mathbf{p}}{\omega_\pm} \cdot \frac{\partial \tilde{N}_\pm}{\partial \mathbf{x}} \simeq \frac{-i \Sigma_{off}^{(bare)}(x; \omega_\pm, \mathbf{p})}{2\omega_\pm}. \tag{5.26}$$

Physical interpretation of these relations will be discussed below.

In calculating some quantity in the bare- f scheme, one carries out perturbative calculation using the propagator $\hat{\Delta}(X; P)$ given in (4.35). The calculated quantity is written in terms of the bare number density $f_B(X; P)$ or $N_{B\pm}(X; E_p, \pm \mathbf{p})$. Using the solution to (5.26), one can rewrite the quantity in terms of the “physical number density” $\tilde{N}_\pm(X; \pm \mathbf{p})$.

2. Physical- f scheme

As defined in Section IVC, we change the definition of the number-density function f . Then, in this physical- f scheme, the number density function f does not obey (4.33). It is convenient to define f , such that the number densities

$$\begin{aligned}
N_+(X; E_{\mathbf{p}}, \hat{\mathbf{p}}) &= f(X; p_0 = E_{\mathbf{p}}, \mathbf{p}), \\
N_-(X; E_{\mathbf{p}}, -\hat{\mathbf{p}}) &= -1 - f(X; p_0 = -E_{\mathbf{p}}, \mathbf{p}),
\end{aligned}$$

are as close as the physical number densities $\tilde{N}_{\pm}(X; \pm\mathbf{p})$. As seen above, the first term on the RHS of (5.17), which is proportional to $\Sigma_{off}(X; P)$, yields a large contribution to the physical number density. Thus, we demand that $f(X; P)$ changes in time X_0 so that $\Sigma_{off}(X; P)$ vanishes:

$$\begin{aligned}
i\Sigma_{off}(X; P) &= \{f(X; P), P^2 - m^2 - Re\Sigma_R(X; P)\} - \tilde{\Gamma}^{(p)}(X; P) \\
&= 0.
\end{aligned} \tag{5.27}$$

Given an initial data at the “initial” time $X_0 = T_{in}$, $f(X_0 = T_{in}, \mathbf{X}; P)$, (5.27) serves as determining $f(X_0, \mathbf{X}; P)$ at later time X_0 . Note that $Re\Sigma_R$ and $\tilde{\Gamma}^{(p)}$ are evaluated order by order in perturbation series. Then, through (5.27), $f(X; P)$ is also determined order by order. (In this relation, see the argument at Section IVC for the mass renormalization in vacuum theory.) We refer (5.27) to as the “renormalization condition” for the number density, which replaces the condition (4.33) in the bare scheme. In the present scheme, transformed self-energy part $\underline{\Sigma}$, Eq. (5.3), is diagonal:

$$\hat{B}_R \hat{\Sigma} \hat{B}_L = \begin{pmatrix} \Sigma_R & 0 \\ 0 & -\Sigma_A \end{pmatrix}. \tag{5.28}$$

It is to be noted that, with the condition (5.27), $\hat{G}(X; P)$ in (5.12) - (5.17) is free from singular contribution in the narrow-width limit (cf. above after (5.17)) and the difference between $\hat{G}$ and the bare propagator $\hat{\Delta}$ leads to a perturbative correction to the physical number density.

In the sequel of this paper, we adopt the physical- f scheme.

Let us inspect the physical implication of the “renormalization condition” (5.27):

$$2P \cdot \partial_X f(X; P) - \{f(X; P), Re\Sigma_R(X; P)\} = \tilde{\Gamma}^{(p)}(X; P). \tag{5.29}$$

Here let us recall the definition (5.19) of the “mass-shell” and find the form of (5.29) on the mass-shell. From (5.19), we have

$$\begin{aligned}
\left[\left\{ 2p_0 - \frac{\partial Re\Sigma_R}{\partial p_0} \right\} \mathbf{v}_{\pm} - 2\mathbf{p} - \frac{\partial Re\Sigma_R}{\partial \mathbf{p}} \right] \Big|_{p_0 = \pm\omega_{\pm}(X; \pm\mathbf{p})} &= 0, \\
\frac{\partial Re\Sigma_R}{\partial X} &= \pm \frac{\partial \omega_{\pm}}{\partial X} \left(2p_0 - \frac{\partial Re\Sigma_R}{\partial p_0} \right) \Big|_{p_0 = \pm\omega_{\pm}(X; \pm\mathbf{p})},
\end{aligned} \tag{5.30}$$

where $\mathbf{v}_{\pm} \equiv \pm \partial \omega_{\pm}(X; \pm\mathbf{p}) / \partial \mathbf{p}$ is the velocity of the $\pm$ mode. We also recall that $f = \theta(p_0)N_+ - \theta(-p_0)(1 + N_-)$, where $N_+ = N_+(X; p_0 = \omega_+, \hat{\mathbf{p}})$ [$N_- = N_-(X; -p_0 = \omega_-, -\hat{\mathbf{p}})$] is the number density of a (quasi)particle [an anti(quasi)particle] with momentum $\mathbf{p}$ [$-\mathbf{p}$]. It is straightforward to show that, on the mass-shell, $p_0 = \pm\omega_{\pm}(X; \pm\mathbf{p})$ ($= \pm\omega_{\pm}$), (5.29) becomes

$$\begin{aligned}
&\frac{\partial N_{\pm}}{\partial X_0} + \mathbf{v}_{\pm} \cdot \nabla_X N_{\pm} \pm \frac{\partial \omega_{\pm}}{\partial X_{\mu}} \frac{\partial N_{\pm}}{\partial P^{\mu}} \Big|_{p_0 = \pm\omega_{\pm}} \\
&= \frac{dN_{\pm}(X; \omega_{\pm}(X; \pm\mathbf{p}), \pm\hat{\mathbf{p}})}{dX_0} + \frac{d\omega_{\pm}}{d\mathbf{p}} \cdot \frac{dN_{\pm}}{d\mathbf{X}} - \frac{\partial \omega_{\pm}}{\partial \mathbf{X}} \cdot \frac{dN_{\pm}}{d\mathbf{p}} \\
&= Z_{\pm} \Gamma_{\pm}^{(p)} \Big|_{p_0 = \pm\omega_{\pm}},
\end{aligned} \tag{5.31}$$

where $Z_{\pm}$ is the wave-function renormalization factor (cf. (5.22)) and, as defined in (5.25), $\Gamma_{\pm}^{(p)}$ is the net production rate of a particle (an antiparticle) with momentum $\mathbf{p}$ [$-\mathbf{p}$]. If we employ the wave-function renormalization condition $Z_{\pm} = 1$, the RHS turns out to the net production rate $\Gamma_{\pm}^{(p)}$. The quasiparticle distribution function $N_{\pm} = N_{\pm}(X; p_0 = \omega_{\pm}, \hat{\mathbf{p}})$ here is nothing but the relativistic Wigner function and (5.31) is the generalized kinetic or Boltzmann equation (cf. [26]) for the relativistic complex-scalar-field system.

Here we make a comment on a role of the second term on the left-hand side (LHS) of (5.29). The second term represents the effect due to the change of “mass;”

$$\{f(X; P), Re\Sigma_R(X; P)\} = \frac{\partial f(X; P)}{\partial X_\mu} \frac{\partial Re\Sigma_R(X; P)}{\partial P^\mu} - \frac{\partial f(X; P)}{\partial P^\mu} \frac{\partial Re\Sigma_R(X; P)}{\partial X_\mu}. \quad (5.32)$$

The first term on the RHS together with the first term on the LHS of (5.29) yields the term being proportional to the first two terms on the LHS of (5.31), while the second term on the RHS yields the term being proportional to the third term on the LHS of (5.31).

One can show using (5.30) that (5.26) is the same in form as (5.31). [Recall that we have set $Z_\pm = 1$ in (5.26).] This observation gives a support for the qualitative argument made above in conjunction with (5.26).

D. Resummation of the lowest-order $\hat{\Sigma}$

To lowest order in λ (cf. (3.3)), a tadpole diagram contributes to $\hat{\Sigma}_{int}$:

$$\begin{aligned} \hat{\Sigma}_{int}(X) &= M^2(X) \hat{\tau}_3, \\ M^2(X) &\simeq i\lambda \int \frac{d^D P}{(2\pi)^D} \Delta_{11}^{(0)}(X; P) + (Z_m Z - 1) m^2 \\ &= i\lambda \int \frac{d^D P}{(2\pi)^D} \left[\frac{1}{P^2 - m^2 + i\eta} \right. \\ &\quad \left. - 2\pi i \{ \theta(p_0) N_+(X; p_0, \hat{\mathbf{p}}) + \theta(-p_0) N_-(X; -p_0, -\hat{\mathbf{p}}) \} \right] \delta(P^2 - m^2) \\ &\quad + (Z_m Z - 1) m^2. \end{aligned} \quad (5.33)$$

Here use has been made of the dimensional regularization: D is the spacetime dimension and $\epsilon = 4 - D$ works as a regulator for the UV divergence. For the purpose of later use, we have adopted here the $F\overline{F}$ representation (cf. (3.36)) for $\Delta_{11}^{(0)}$, (4.44) with (4.45) and (4.47). $N_\pm$ in (5.34) is as in (4.30). Note that $\Delta_{11}^{(1)}$ in (4.44) with (4.49) leads to a contribution, which is of higher order in λ . It should be kept in mind that, in dealing with $O(\lambda^2)$ contributions, (5.34) with $\Delta_{11}^{(1)}$ for $\Delta_{11}^{(0)}$, as well as the two-loop contribution, should be taken into account.

Standard manipulation yields

$$\begin{aligned} M^2(X) &= M_0^2 + M_{induced}^2(X), \\ M_0^2 &= -\frac{\lambda}{(4\pi)^2} m^2 \left[\frac{2}{\epsilon} + 1 - \ln \frac{m^2}{\mu_d^2} \right], \end{aligned} \quad (5.35)$$

$$M_{induced}^2(X) = \lambda \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_p} [N_+(X; E_p, \hat{\mathbf{p}}) + N_-(X; E_p, -\hat{\mathbf{p}})], \quad (5.36)$$

where

$$\frac{1}{\epsilon} \equiv \frac{1}{4 - D} - \frac{\gamma_E}{2} + \frac{1}{2} \ln(4\pi). \quad (5.37)$$

Here M_0^2 is the squared self mass in vacuum theory, γ_E is the Euler constant, and μ_d is an arbitrary scale parameter. As an UV renormalization scheme in vacuum theory, we are adopting the $\overline{\text{MS}}$ scheme. Thus, at the present lowest order, we have

$$Z_m = 1 + \frac{\lambda}{(4\pi)^2} \frac{2}{\epsilon}, \quad Z = 1,$$

and then

$$M_0^2 = \frac{\lambda}{(4\pi)^2} m^2 \ln \frac{m^2}{e\mu_d^2}. \quad (5.38)$$

If we choose $\mu_d = m/\sqrt{e}$, we have $M_0^2 = 0$.

In passing, for a locally thermal equilibrium system, we have

$$N_+(X; E_p, \hat{\mathbf{p}}) = N_-(X; E_p, -\hat{\mathbf{p}}) = \frac{1}{e^{E_p/T(X)} - 1},$$

where $T(X)$ is the local temperature of the system. In this case, we obtain

$$M_{induced}^2(X) = \frac{\lambda}{2\pi^2} m^2 \int_1^\infty du \frac{\sqrt{u^2 - 1}}{e^{mu/T(X)} - 1}.$$

In the high-temperature regime, $T(X) \gg m$, we have

$$M_{induced}^2(X) \simeq \frac{\lambda}{12} T^2(X) \left(1 - \frac{3}{\pi} \frac{m}{T(X)}\right) \quad (T(X) \gg m), \quad (5.39)$$

while, at low temperature, we have

$$M_{induced}^2(X) \simeq \frac{\lambda m^2}{(2\pi)^{3/2}} \left(\frac{T(X)}{m}\right)^{3/2} e^{-m/T(X)} \left(1 + \frac{3}{8} \frac{T(X)}{m}\right) \quad (T(X) \ll m). \quad (5.40)$$

Let us construct a $\hat{\Sigma}$ -resummed propagator. To the present lowest order, $\Sigma_{12} = \Sigma_{21} = 0$ and then, from (5.7), $\tilde{\Gamma}^{(p)}$ vanishes. Then, the renormalization condition (5.29), $\Sigma_{off} = 0$, reads

$$i\Sigma_{off}(X; P) = 2P \cdot \partial_X f(X; P) + \frac{\partial f(X; P)}{\partial P^\mu} \frac{\partial M^2(X)}{\partial X_\mu} = 0, \quad (5.41)$$

whose approximate solution is given by

$$f(X; P) \simeq \mathcal{F}(\underline{\mathbf{X}}; P^2 - M^2(X), \underline{\mathbf{p}}) \simeq G(\underline{\mathbf{X}}; \underline{P}). \quad (5.42)$$

Here, $\mathcal{F}$ or G is an arbitrary function and

$$\begin{aligned} \underline{p}_0 &\equiv p_0 - \frac{\partial M^2(X)}{\partial X^0} \frac{(X_0 - T_{in})}{2p_0}, \\ \underline{p}_i &\equiv p_i - \frac{\partial M^2(X)}{\partial X^i} \frac{(X_0 - T_{in})}{2p_0} \quad (\text{no summation over } i), \\ \underline{X}_i &\equiv X_i - \frac{p_i}{p_0} (X_0 - T_{in}) + \frac{p_0 \partial M^2(X) / \partial X^i - p_i \partial M^2(X) / \partial X^0}{4(p_0)^3} (X_0 - T_{in})^2. \end{aligned} \quad (5.43)$$

Given an initial data $f(X_0 = T_{in}, \underline{\mathbf{X}}; P) = G(\underline{\mathbf{X}}; P)$, (5.42) gives $f(X_0, \underline{\mathbf{X}}; P)$ at later time X_0 to $O(\lambda)$.

Since $Im\Sigma_R = \tilde{\Gamma}^{(p)} = 0$, $\hat{G}^{(1)}$ in (5.17) vanishes due to the condition (5.41). Then, the $\hat{\Sigma}$ -resummed propagator $\hat{G}(X; P)$ reads

$$\begin{aligned} \hat{G}(X; P) &\simeq \hat{G}^{(0)}(X; P) \\ &= G_{RA} + f(X; P) (G_R - G_A) \hat{A}_+, \end{aligned} \quad (5.44)$$

where

$$\begin{aligned} G_{RA} &= \begin{pmatrix} G_R & 0 \\ G_R - G_A & -G_A \end{pmatrix}, \\ G_{R(A)} &= G_{R(A)}(X; P) = \frac{1}{P^2 - m^2 - M^2(X) \pm i\epsilon(p_0)\eta}. \end{aligned} \quad (5.45)$$

E. Changing the mass parameter

The analysis in the last subsection shows that an efficient perturbative scheme is obtained as follows. Rewrite the Lagrangian density (3.3) by introducing a new “mass function” $M(x)$, which is assumed to depend weakly on x :

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_0 + \mathcal{L}_{int} + \mathcal{L}_{rc}, \\
\mathcal{L}_0 &= \partial_\mu \phi^\dagger(x) \partial^\mu \phi(x) - M^2(x) \phi^\dagger(x) \phi(x), \\
\mathcal{L}_{int} &= -\frac{\lambda}{4} (\phi^\dagger(x) \phi(x))^2, \\
\mathcal{L}_{rc} &= \chi(x) \phi^\dagger(x) \phi(x) + (Z-1) \partial_\mu \phi^\dagger(x) \partial^\mu \phi(x) - (Z_m Z - 1)(M^2(x) - \chi(x)) \phi^\dagger(x) \phi(x) \\
&\quad - \frac{\lambda}{4} (Z_\lambda Z^2 - 1) (\phi^\dagger(x) \phi(x))^2,
\end{aligned} \tag{5.46}$$

where

$$\chi(x) \equiv M^2(x) - m^2. \tag{5.47}$$

Note that M^2 here is not M^2 in (5.34). As discussed in [10], perturbation theory based on (5.46) is UV-renormalizable. We give an simple argument for this in Appendix D.

It should be emphasized [10] that the perturbation theory based on (5.46) is defined by taking $\mathcal{L}_0$ as a nonperturbative or “free” part and $\mathcal{L}_{int} + \mathcal{L}_{rc}$ as perturbative part. Then, $\chi(x)\phi^\dagger\phi$ in $\mathcal{L}_{rc}$ is a part of the perturbative terms. Thus, in traditional perturbation expansion, we assign

$$M^2(x) = O(\lambda^0), \quad \chi(x) (= M^2(x) - m^2) = O(\lambda), \tag{5.48}$$

and, in loop or δ expansion (cf. the next section), we assign

$$M^2(x) = O(\delta^0), \quad \chi(x) (= M^2(x) - m^2) = O(\delta), \tag{5.49}$$

It is obvious from the argument in the above subsection that, in the bare- f scheme in perturbation theory based on (5.46), the bare propagator is given by (5.44),

$$\hat{G}(X; P) = G_{RA} + f(X; P) (G_R - G_A) \hat{A}_+, \tag{5.50}$$

where

$$G_{R(A)} = G_{R(A)}(X; P) = \frac{1}{P^2 - M^2(X) \pm i\epsilon(p_0)\eta}. \tag{5.51}$$

The number-density function $f(X; P)$ in (5.50) obeys (5.41).

This result may also be derived through the same procedure as in Section IV. This time we start with the $|p_0|$ -prescription-formula (4.31) from the beginning,

$$\hat{G}(x, y) = \int d^4u d^4v \hat{B}_L(x, u) \hat{G}_{diag}(u, v) \hat{B}_R(v, y). \tag{5.52}$$

Here $\hat{G}_{diag} = \text{diag}(G_R, -G_A)$ with $G_{R(A)}(x, y)$ a solution to

$$(\partial_x^2 + M^2(x)) G_{R(A)}(x, y) = (\partial_y^2 + M^2(y)) G_{R(A)}(x, y) = -\delta^4(x - y), \tag{5.53}$$

with retarded (advanced) boundary condition. It can easily be shown that the solution to (5.53) is (5.51). Note that no first X^μ -derivative term appears there. In place of (4.25), we have

$$\begin{aligned}
(\partial_x^2 + M^2(x)) \hat{G}(x, y) &= -\hat{\tau}_3 \delta^4(x - y) + \int d^4u d^4v \left[\{(\partial_x^2 - \partial_u^2) \right. \\
&\quad \left. + M^2(x) - M^2(u)\} \hat{B}_L(x, u) \right] \hat{\tau}_3 \hat{B}_R(u, v) \hat{\tau}_3 \hat{G}(v, y),
\end{aligned} \tag{5.54}$$

and in place of (4.41), we have

$$\hat{\mathcal{L}}_c = - \int d^4u d^4v \hat{\phi}^\dagger(x) \hat{\tau}_3 \left[\{(\partial_x^2 - \partial_u^2) + M^2(x) - M^2(u)\} \hat{B}_L(x, u) \right] \hat{\tau}_3 \hat{B}_R(u, v) \hat{\tau}_3 \hat{\phi}(v).$$

This yields a two-point vertex function

$$\begin{aligned}
i\hat{\mathcal{V}}_c(X; P) &= 2P \cdot \partial_X f(X; P) + \frac{dM^2(X)}{dX_\mu} \frac{\partial f(X; P)}{\partial P^\mu} \hat{A}_- \\
&= \{f(X; P), P^2 - M^2(X)\} \hat{A}_-.
\end{aligned} \tag{5.55}$$

Since we are to obtain the bare propagator, $\hat{\Sigma}_{int} = \tilde{\Gamma}^{(p)} = 0$. Then, the “renormalization condition” (5.27) turns out to

$$\{f(X; P), P^2 - M^2(X)\} = 0, \tag{5.56}$$

with which we have $\hat{\mathcal{V}}_c(X; P) = 0$. Equation (5.56) is nothing but (5.41). Using this condition, we can derive (5.50) from (5.52), which is the bare propagator in the bare- f scheme. The physical- f scheme may be constructed in a similar manner as in Section IVC.

In this theory, $\hat{\Sigma}_{int}$ to $O(\lambda)$ reads

$$\hat{\Sigma}_{int}(X) = \hat{\tau}_3 \left[\frac{\lambda}{(4\pi)^2} M^2(X) \ln \frac{M^2(X)}{e\mu_d^2} + M_{induced}^2(X) - M^2(X) + m^2 \right],$$

where $M_{induced}^2(X)$ is as in (5.36) with $M^2(X)$ for m^2 and $-\hat{\tau}_3[M^2(X) - m^2]$ has come from $\chi\phi^\dagger\phi \in \mathcal{L}_{rc}$, Eq. (5.46).

Up to this point, the function $M^2(X)$ (≥ 0) is arbitrary. An efficient perturbation theory is obtained by demanding, e.g., $\hat{\Sigma}_{int}(X) = 0$;

$$M^2(X) \left[1 - \frac{\lambda}{(4\pi)^2} \ln \frac{M^2(X)}{e\mu_d^2} \right] = m^2 + M_{induced}^2(X). \tag{5.57}$$

The scheme with this $M(X)$ may be called “presummation scheme.” Equation (5.57) is the gap equation, which serves as determining $M^2(X)$ in terms of μ_d , which is still an arbitrary parameter.

It is worth mentioning that, in the case where the condition $[\rho, Q] = 0$, Eq. (3.6), is not fulfilled, we may generalize (5.46), although not necessary, by changing $\mathcal{L}_0$ to

$$\mathcal{L}_0 = \partial^\mu \phi^\dagger(x) \partial_\mu \phi(x) - M_+(x) \phi^{(+)\dagger}(x) \phi^{(+)}(x) - M_-(x) \phi^{(-)\dagger}(x) \phi^{(-)}(x),$$

and changing $\mathcal{L}_{rc}$ accordingly. Here the superscript “(+)” [“(−)”] refers to the positive [negative] frequency part, and then $M_+(x)$ [$M_-(x)$] is an effective mass of a quasiparticle [anti-quasiparticle].

VI. THE $O(N)$ LINEAR σ MODEL

A. Preliminary

In this section, we deal with a fermionless $O(N)$ linear σ model, in which N real-scalar fields constitute a vector representation $\vec{\phi} = {}^t(\phi_1, \dots, \phi_n)$ of $O(N)$. The case of $N = 4$ is of practical interest. Specifically, through computing the effective action, we analyze how the phase transition proceeds. The Lagrangian of the theory reads

$$\mathcal{L} = \frac{1}{2} \left[(\partial \vec{\phi}_B)^2 - \mu_B^2 \vec{\phi}_B^2 \right] - \frac{\lambda_B}{4!} (\vec{\phi}_B^2)^2 + H_B ({}^t\vec{e} \cdot \vec{\phi}_B), \tag{6.1}$$

where $\mu_B^2 < 0$, so that the theory describes the system whose ground state is in a broken phase in the classical limit. In (6.1), we have introduced the term $H_B ({}^t\vec{e} \cdot \vec{\phi}_B)$, with $\vec{e}$ an unit vector, which explicitly breaks $O(N)$ symmetry. Noticing the fact that an UV-renormalization scheme for the symmetric phase ($\mu_B^2 > 0$) works [27] as it is for the symmetry-broken phase ($\mu_B^2 < 0$), we introduce UV-renormalized quantities,

$$\vec{\phi}_B = \sqrt{Z} \vec{\phi}, \quad \mu_B^2 = Z_\mu \mu^2, \quad \lambda_B = Z_\lambda \lambda.$$

In terms of renormalized quantities, $\mathcal{L}$ reads

$$\begin{aligned}
\mathcal{L}(\vec{\phi}) &= \frac{1}{2} \left[(\partial \vec{\phi})^2 - \mu^2 \vec{\phi}^2 \right] - \frac{\lambda}{4!} (\vec{\phi}^2)^2 + H ({}^t\vec{e} \cdot \vec{\phi}) + \frac{1}{2} (Z - 1) \left[(\partial \vec{\phi})^2 - \mu^2 \vec{\phi}^2 \right] \\
&\quad - (Z_\lambda Z^2 - 1) \frac{\lambda}{4!} (\vec{\phi}^2)^2 + A \mu^4,
\end{aligned} \tag{6.2}$$

where $H = \sqrt{Z}H_B$. In (6.2), according to [10], we have introduced the c-number counter term $A\mu^4$, which is necessary to make the effective action finite.

For the purpose of later use, we here construct

$$\mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x)) \equiv \mathcal{L}(\vec{\phi}(x) + \vec{\varphi}(x)) - \frac{\partial \mathcal{L}(\vec{\varphi}(x))}{\partial \vec{\varphi}(x)} \cdot \vec{\phi}(x). \quad (6.3)$$

To avoid too many notations, for $\mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x))$, we have used the same letter “ $\mathcal{L}$ ” as in (6.1) and (6.2). As will be discussed below, $\vec{\varphi}(x)$ in (6.3) is the (classical) condensate or order-parameter fields and $\vec{\phi}(x)$ is the quantum fields, which describes the fluctuation around $\vec{\varphi}(x)$. Straightforward manipulation yields

$$\begin{aligned} \mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x)) &= \mathcal{L}_{cond} + \mathcal{L}'_{cond} + \mathcal{L}'_q + \mathcal{L}'_{rc}, \\ \mathcal{L}_{cond} &= \frac{1}{2} [(\partial \vec{\varphi})^2 - \mu^2 \vec{\varphi}^2] - \frac{\lambda}{4!} (\vec{\varphi}^2)^2 + H({}^t\vec{e} \cdot \vec{\varphi}), \end{aligned} \quad (6.4)$$

$$\mathcal{L}'_{cond} = \frac{1}{2} (Z - 1)(\partial \vec{\varphi})^2 - \frac{1}{2} (Z_\mu Z - 1)\mu^2 \vec{\varphi}^2 - (Z_\lambda Z^2 - 1)\frac{\lambda}{4!} (\vec{\varphi}^2)^2 + A\mu^4, \quad (6.5)$$

$$\begin{aligned} \mathcal{L}'_q &= \frac{1}{2} [(\partial \vec{\phi})^2 - \mu^2 \vec{\phi}^2] - \frac{\lambda}{12} [\vec{\varphi}^2 \vec{\phi}^2 + 2({}^t\vec{\varphi} \cdot \vec{\phi})^2] - \frac{\lambda}{3!} ({}^t\vec{\varphi} \cdot \vec{\phi}) \vec{\phi}^2 - \frac{\lambda}{4!} (\vec{\phi}^2)^2, \\ \mathcal{L}'_{rc} &= \frac{1}{2} (Z - 1)(\partial \vec{\phi})^2 - \frac{1}{2} (Z_\mu Z - 1)\mu^2 \vec{\phi}^2 - (Z_\lambda Z^2 - 1) \left[\frac{\lambda}{12} \{ \vec{\varphi}^2 \vec{\phi}^2 + 2({}^t\vec{\varphi} \cdot \vec{\phi})^2 \} \right. \\ &\quad \left. + \frac{\lambda}{3!} ({}^t\vec{\varphi} \cdot \vec{\phi}) \vec{\phi}^2 + \frac{\lambda}{4!} (\vec{\phi}^2)^2 \right]. \end{aligned} \quad (6.6)$$

It should be noted that $\mathcal{L}$ enjoys $O(N - 1)$ symmetry.

For obtaining an efficient or rather physically-sensible (perturbative) scheme, as in Section VE, it is necessary [10,28–30]) to introduce weakly x -dependent masses, $M_\pi(x)$ and $M_\sigma(x)$, for which we assume that

$$|\mu^2 - M_\pi^2(x)|, |M_\pi^2(x) - M_\sigma^2(x)| = O(\delta). \quad (6.7)$$

How to determine $M_\pi^2(x)$ and $M_\sigma^2(x)$ will be discussed in subsection D. We then rewrite $\mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x))$ in the form,

$$\mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x)) = \mathcal{L}_{cond} + \mathcal{L}'_{cond} + \mathcal{L}_0(\vec{\phi}(x); \vec{\varphi}(x)) + \mathcal{L}_{int} + \mathcal{L}_{rc}. \quad (6.8)$$

Here $\mathcal{L}_{cond}$ and $\mathcal{L}'_{cond}$ are as in (6.4) and (6.5), respectively, and

$$\mathcal{L}_0(\vec{\phi}(x); \vec{\varphi}(x)) = \frac{1}{2} {}^t\vec{\phi}(x) \mathbf{\Delta}^{-1}(x) \vec{\phi}(x), \quad (6.9)$$

$$\mathcal{L}_{int} = -\frac{\lambda}{3!} ({}^t\vec{\varphi} \cdot \vec{\phi}) \vec{\phi}^2 - \frac{\lambda}{4!} (\vec{\phi}^2)^2,$$

$$\mathcal{L}_{rc} = \mathcal{L}'_{rc} + \frac{1}{2} {}^t\vec{\phi}(x) [\chi_\pi(x) \mathbf{P}_\pi(x) + \chi_\sigma(x) \mathbf{P}_\sigma(x)] \vec{\phi}(x). \quad (6.10)$$

Here $\mathcal{L}'_{rc}$ is as in (6.6) and

$$\begin{aligned} \mathbf{\Delta}^{-1}(x) &\equiv - \left(\partial^2 + M_\pi^2(x) + \frac{\lambda}{6} \vec{\varphi}^2(x) \right) \mathbf{P}_\pi(x) - \left(\partial^2 + M_\sigma^2(x) + \frac{\lambda}{2} \vec{\varphi}^2(x) \right) \mathbf{P}_\sigma(x), \\ \chi_\xi(x) &= M_\xi^2(x) - \mu^2 \quad (\xi = \pi, \sigma), \\ \mathbf{P}_\pi(x) &= \mathbf{I} - |\varphi(x)\rangle \langle \varphi(x)|, \\ \mathbf{P}_\sigma(x) &= |\varphi(x)\rangle \langle \varphi(x)|. \end{aligned} \quad (6.11)$$

In these equations, the boldface letters denote $N \times N$ matrices in the real vector space: $\mathbf{P}_\pi$ ($\mathbf{P}_\sigma$) is the projection operator onto the π - (σ)-subspace; $\mathbf{I}$ is an unit matrix. $|\varphi(x)\rangle$ is a unit vector along $\vec{\varphi}(x)$, and $\langle \varphi(x)|$ is an adjoint of $|\varphi(x)\rangle$. As has been noted above, when compared to $M_\xi^2(x)$ ($\xi = \pi, \sigma$), $\chi_\xi(x)$ is of higher order.

As mentioned above, the quantum fields $\vec{\phi}$ describe the fluctuation around the condensate fields $\vec{\varphi}(x)$. The construction of perturbation theory based on (6.8) starts with constructing the Fock space of the quanta described by $\vec{\phi}$, which is defined “on $\vec{\varphi}(x)$.” As stated in Section I, dealing with the dynamical aspect of the system, we are assuming that microscopic or elementary reactions can be regarded as taking place in a single spacetime cell. While, in a single cell, relaxation phenomena are negligible. This means that the theory to be developed in the sequel is consistent only when $\vec{\varphi}(x)$ does not change appreciably in a single spacetime cell. We shall assume that this is the case and treat x of $\vec{\varphi}(x)$ as macroscopic spacetime coordinates.

B. Effective action

Spacetime evolution of the condensate fields $\vec{\varphi}(x)$ is governed by the Euler-Lagrange equation, which is derived from the CTP effective action $\Gamma_C[\vec{\varphi}]$. If one restricts to the case where $\vec{\varphi}(x)$ is space-time independent, $\vec{\varphi}(x) \rightarrow \vec{\varphi}$, $\Gamma_C[\vec{\varphi}]$ reduces to minus the CTP effective potential, $\Gamma_C[\vec{\varphi}] = -V_C(\vec{\varphi})$. The *physical part* of the CTP effective potential (cf. below) $V_{phys}(\vec{\varphi})$ is the minimum free energy density of the system under the constraint, $\langle \phi(x) \rangle = \varphi$, the minimum free energy density with quantum and medium effects taken into account. A computational algorithm of effective potential in vacuum theory is established in [31]. On the basis of a generalization of this algorithm to the case of equilibrium thermal field theory, a calculational scheme of the effective potential and its *physical part* have been settled [32] (see also [33]).

We generalize this algorithm to the present case of out of equilibrium systems. Following standard procedure [2] of obtaining an expression for the CTP effective action, we start with the CTP generating functional for Green functions in the interaction picture (cf. (3.7)),

$$Z_C[\vec{J}] \equiv e^{iW_C[\vec{J}]} = \langle T_C \exp i \int_C [\mathcal{L}'(\vec{\phi}(x)) + \vec{J}(x) \cdot \vec{\phi}(x)] \rangle, \quad (6.12)$$

where, as in Section II, $C (= C_1 \oplus C_2)$ is the closed time-path in the complex time-plane and $\mathcal{L}'$ stands for the perturbative part of the Lagrangian density, which is to be identified below in (6.20). In (6.12), $W_C[J]$ is the generating functional for connected Green functions. Expectation value of $\vec{\phi}(x)$, $\langle \vec{\phi}(x) \rangle_J (\equiv \vec{\varphi}(x))$, in the presence of the external source $\vec{J}(x)$ is obtained through

$$\vec{\varphi}(x) = \frac{\delta W_C[\vec{J}]}{\delta \vec{J}(x)}. \quad (6.13)$$

The CTP effective action Γ_C is defined as the functional Legendre transform,

$$\Gamma_C[\vec{\varphi}] = W_C[\vec{J}] - \int_C \vec{J}(x) \cdot \vec{\varphi}(x), \quad (6.14)$$

from which the Euler-Lagrange equation follows:

$$\frac{\delta \Gamma_C[\vec{\varphi}]}{\delta \vec{\varphi}(x)} = -\vec{J}(x). \quad (6.15)$$

The CTP effective action Γ_C is the generating functional for connected, one-particle irreducible, amputated Green functions. Substitution of (6.12) and (6.15) into (6.14) yields

$$e^{i\Gamma_C[\vec{\varphi}]} = \langle T_C \exp \left(i \int_C \left[\mathcal{L}'(\vec{\phi}(x)) - \frac{\delta \Gamma_C[\vec{\varphi}]}{\delta \vec{\varphi}(x)} \cdot \{\vec{\phi}(x) - \vec{\varphi}(x)\} \right] \right) \rangle.$$

Redefining $\vec{\phi}(x) - \vec{\varphi}(x)$ as new quantum fields $\vec{\phi}(x)$, we have

$$e^{i\Gamma_C[\vec{\varphi}]} = \langle T_C \exp \left[\frac{i}{\delta} \int_C \left\{ \mathcal{L}'(\vec{\phi}(x) + \vec{\varphi}(x)) - \frac{\delta \Gamma_C[\vec{\varphi}]}{\delta \vec{\varphi}(x)} \cdot \vec{\phi}(x) \right\} \right] \rangle. \quad (6.16)$$

Thus, the new quantum fields $\vec{\phi}(x)$ describe the fluctuation around $\vec{\varphi}(x)$. Here, for the purpose stated below, we have inserted the factor $\delta (= 1)$. Going to the two-component representation, we have

$$e^{i\hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]} = \langle T_C \exp \left[\frac{i}{\delta} \int d^4x \sum_{j=1}^2 \left\{ (-)^{j-1} \mathcal{L}'(\vec{\phi}_j(x) + \vec{\varphi}_j(x)) - \frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_j(x)} \cdot \vec{\phi}_j(x) \right\} \right] \rangle, \quad (6.17)$$

where $\vec{\phi}_j$ and $\vec{\varphi}_j$ ($j = 1, 2$) are the type- j fields and

$$\frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_j(x)} = (-)^{j-1} \frac{\delta \Gamma_C[\vec{\varphi}]}{\delta \vec{\varphi}(x)} \quad \text{with } x_0 \in C_j \quad (j = 1, 2). \quad (6.18)$$

The minus sign on the RHS of (6.18) with $j = 2$ comes from the fact that, on C_2 , the time ticks away backward from $+\infty$ to $-\infty$ (cf. (2.4)).

Equation (6.17) is of the form which allows for computing $\hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]$ perturbatively. Perturbative calculation of (6.17) goes by taking suitable

$$\hat{\mathcal{L}}_0 \equiv \sum_{j=1}^2 (-)^{j-1} \mathcal{L}_0(\vec{\phi}_j(x), \vec{\varphi}_j(x)) \quad (\in \hat{\mathcal{L}} \equiv \sum_{j=1}^2 (-)^{j-1} \mathcal{L}(\vec{\phi}_j(x) + \vec{\varphi}_j(x))), \quad (6.19)$$

being bilinear in quantum fields $\vec{\phi}_j(x)$, as the free hat-Lagrangian density, which defines 2×2 propagators of the fields $\vec{\phi}(x) = {}^t(\vec{\phi}_1(x), \vec{\phi}_2(x))$. The form of $\mathcal{L}(\vec{\phi}(x) + \vec{\varphi}(x))$ is already been obtained in the last subsection and $\mathcal{L}_0$ is defined in (6.9) with (6.11). The 2×2 matrix propagators will be constructed in the next subsection. The vertex factors can be reads off from

$$\hat{\mathcal{L}} - \sum_{j=1}^2 (-)^{j-1} \mathcal{L}(\vec{\varphi}_j(x)) - \hat{\mathcal{L}}_0 - \sum_{j=1}^2 \frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_j(x)} \cdot \vec{\phi}_j(x). \quad (6.20)$$

Note that, to zeroth order (tree level),

$$\hat{\Gamma}^{(0)}[\vec{\varphi}_1, \vec{\varphi}_2] = \sum_{j=1}^2 (-)^{j-1} \int d^4x \mathcal{L}(\vec{\varphi}_j(x)). \quad (6.21)$$

Then, the exponent in (6.17) with $\hat{\Gamma}^{(0)}$ for $\hat{\Gamma}$ on the RHS is free from the term being linear in $\vec{\phi}$, which means that the “one-point vertex” is absent. [In constructing $\mathcal{L}(\vec{\phi}(x); \vec{\varphi}(x))$ above, Eq. (6.3), this has already been taken into account.] Thus, in terms of Feynman diagrams, (6.17) consists of bubble or no-leg diagrams. The loop expansion is defined as an expansion by δ (δ -expansion), which is set to 1 at the end of calculation (see, e.g., [34]). Computation of $\hat{\Gamma}$ should be carried out successively: In computing l th loop-order contribution, the contributions to $\hat{\Gamma}$ up to $(l-1)$ th loop-order should be substituted for $\hat{\Gamma}$ on the RHS of (6.17). This part, $-i\delta^{-1} \sum_{j=1}^2 \vec{\phi}_j \cdot \delta \hat{\Gamma} / \delta \vec{\varphi}_j$, plays a roll of eliminating the contribution of one-particle reducible diagrams.

The one-loop contribution to $\hat{\Gamma}$ will be dealt with in Section VII. Computation of a contribution $\hat{\Gamma}^{(n)}$ from a multi-loop diagram, which includes n (≥ 1) vertices, goes in standard manner. Each vertex consists of two types, the type-1 and type-2. Then, $\hat{\Gamma}^{(n)}$ is obtained by summing over types of vertices, so that $\hat{\Gamma}^{(n)}$ consists of 2^n terms, which we can write, with obvious notation, as

$$\hat{\Gamma}^{(n)}[\vec{\varphi}_1, \vec{\varphi}_2] = \sum_{i_1, i_2, \dots, i_n=1}^2 \hat{\mathcal{F}}_{i_1, i_2, \dots, i_n}[\vec{\varphi}_1, \vec{\varphi}_2], \quad (6.22)$$

where i_j ($j = 1, 2, \dots, n$) denotes the type of vertex.

In the two-component representation, the Euler-Lagrange equation (6.15) reads (cf. above after (6.18)),

$$\frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_j(x)} = -(-)^{j-1} \vec{J}_j(x) \quad (j = 1, 2). \quad (6.23)$$

In the limit of absence of the external sources, $\vec{J}_1(x) = \vec{J}_2(x) = 0$, $\vec{\varphi}_1(x) = \vec{\varphi}_2(x)$ holds and (6.23) becomes

$$\left. \frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_j(x)} \right|_{\vec{\varphi}_1 = \vec{\varphi}_2 = \vec{\varphi}} = 0, \quad (j = 1, 2). \quad (6.24)$$

The equation with $j = 1$ is the equation of motion for the physical or type-1 condensate fields, while the one with $j = 2$ is for the unphysical or the type-2 condensate fields. The *physical* effective action Γ_{phys} and the *unphysical* effective action Γ_{unphys} are defined through

$$\begin{aligned} \frac{\delta \Gamma_{phys}[\vec{\varphi}]}{\delta \vec{\varphi}(x)} &= \left. \frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_1(x)} \right|_{\vec{\varphi}_1 = \vec{\varphi}_2 = \vec{\varphi}}, \\ \frac{\delta \Gamma_{unphys}[\vec{\varphi}]}{\delta \vec{\varphi}(x)} &= \left. \frac{\delta \hat{\Gamma}[\vec{\varphi}_1, \vec{\varphi}_2]}{\delta \vec{\varphi}_2(x)} \right|_{\vec{\varphi}_1 = \vec{\varphi}_2 = \vec{\varphi}}. \end{aligned} \quad (6.25)$$

With suitable (functional) integration constant, (6.25) determines Γ_{phys} and Γ_{unphys} . In the case of thermally equilibrium system, the physical effective potential (Γ_{phys} with constant $\vec{\varphi}$) has been computed in [32] up to two-loop order.

At the previous subsection, in defining $\mathcal{L}_0$, we have introduced (arbitrary) “mass functions” $M_\xi^2(x)$ ($\xi = \pi, \sigma$). The free hat-Lagrangian density in the present CTP formalism is given by (6.19). Then, as in the fields, there are two types of $M_\xi^2(x)$, the type-1 and type-2, which we write as $M_{\xi j}^2(x)$ ($j = 1, 2$). [$M_{\xi j}^2(x) = M_\xi^2(x)$ with $x_0 \in C_j$ ($j = 1, 2$).] As will be discussed in subsection D, various methods of determining $M_{\xi j}^2(x)$ are available. All those methods are based on the notion of “optimization” of some physical quantity. The determined $M_{\xi j}^2(x)$ generally depends on $\vec{\varphi}_1(x)$ and $\vec{\varphi}_2(x)$. Let $\hat{\Gamma}[\vec{\varphi}_1; \vec{\varphi}_2; M_{\pi 1}^2, M_{\pi 2}^2, M_{\sigma 1}^2, M_{\sigma 2}^2]$ be the effective action computed to some order of loop expansion. Here a question arises: In computing the equation of motion (6.24), does the functional derivative $\delta/\delta\vec{\varphi}_j(x)$ ($j = 1, 2$) act on $M_{\xi 1}^2(x)$ and $M_{\xi 2}^2(x)$? The answer is as follows: If we intend to “optimize” the effective action, from which the equation of motion is derived, $\delta/\delta\vec{\varphi}_j(x)$ should act on $M_{\xi 1}^2(x)$ and $M_{\xi 2}^2(x)$. On the contrary, if we intend to “optimize” the equation of motion, $\delta/\delta\vec{\varphi}_j(x)$ should not act on $M_{\xi 1}^2(x)$ and $M_{\xi 2}^2(x)$. We take the stand that the effective action is an intermediate device to derive an equation of motion for $\vec{\varphi}_j(x)$ ($j = 1, 2$), etc., and then the equation of motion is of physical importance over the effective action. On the basis of this observation, following [10], we take the latter throughout in the sequel, i.e., $\delta/\delta\vec{\varphi}_j(x)$ does not act on $M_{\xi 1}^2(x)$ and $M_{\xi 2}^2(x)$.

C. Propagator

1. Retarded/advanced propagator

Here, we obtain a retarded/advanced propagator, $\hat{\Delta}_{diag}$, which is an inverse of

$$\hat{\Delta}^{-1}(x) = \Delta^{-1}(x) \hat{\tau}_3, \quad (6.26)$$

with $\Delta^{-1}(x)$ as in (6.11), under the retarded/advanced boundary condition (cf. (4.2) and (4.4)).

We start with finding an inverse of $\Delta^{-1}(x)$ under the retarded boundary condition:

$$\Delta^{-1}(x) \Delta_R(x, y) = \mathbf{I} \delta^4(x - y). \quad (6.27)$$

To this end, we introduce

$$\begin{aligned} \Delta_R^{(0)}(x, y) &= \Delta_R^{(\pi)}(x, y) + \Delta_R^{(\sigma)}(x, y), \\ \Delta_R^{(\xi)}(x, y) &= \mathbf{P}_\xi(x) \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \frac{1}{P^2 - \mathcal{M}_\xi^2(X)} \quad (\xi = \pi, \sigma), \end{aligned} \quad (6.28)$$

where $X = (x + y)/2$ and

$$\mathcal{M}_\pi^2(X) = M_\pi^2(X) + \frac{\lambda}{6} \vec{\varphi}^2(X), \quad \mathcal{M}_\sigma^2(X) = M_\sigma^2(X) + \frac{\lambda}{2} \vec{\varphi}^2(X). \quad (6.29)$$

$\mathcal{M}_\xi^2(X)$ in (6.28) should be understood to be $\mathcal{M}_\xi^2(X) - i\epsilon(p_0)\eta$.

Straightforward manipulation using (C2) in Appendix C yields

$$\begin{aligned} \Delta^{-1}(x) \Delta_R^{(0)}(x, y) &= \mathbf{I} \delta^4(x - y) - 2i \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} P^\mu \frac{\partial |\varphi(X)\rangle \langle \varphi(X)|}{\partial X^\mu} \\ &\quad \times \left[\frac{1}{P^2 - \mathcal{M}_\pi^2(X)} - \frac{1}{P^2 - \mathcal{M}_\sigma^2(X)} \right]. \end{aligned} \quad (6.30)$$

With (6.30) in hand, we are now ready to obtain $\Delta_R(x, y)$ in (6.27) to the gradient approximation:

$$\begin{aligned} \Delta_R(x, y) &\simeq \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(x) \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \frac{1}{P^2 - \mathcal{M}_\xi^2(X)} \\ &\quad + 2i \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \frac{P \cdot \partial_X \mathbf{P}_\sigma(X)}{P^2 - \mathcal{M}_\xi^2(X)} \\ &\quad \times \left[\frac{1}{P^2 - \mathcal{M}_\pi^2(X)} - \frac{1}{P^2 - \mathcal{M}_\sigma^2(X)} \right]. \end{aligned}$$

Using (C2) in Appendix C and Fourier transforming the resultant expression, we obtain

$$\begin{aligned}\Delta_R(X; P) &\simeq \sum_{\xi=\pi, \sigma} \frac{\mathbf{P}_\xi(X)}{P^2 - \mathcal{M}_\xi^2(X)} \\ &\quad - iP_\mu \left(|\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)| \right) \left[\frac{1}{P^2 - \mathcal{M}_\pi^2(X)} - \frac{1}{P^2 - \mathcal{M}_\sigma^2(X)} \right]^2.\end{aligned}\quad (6.31)$$

It can be shown that $\Delta_R(x, y)$, inverse Fourier transform of $\Delta_R(X; P)$, satisfies

$$\Delta_R(x, y) \overleftarrow{\Delta}^{-1}(y) = \mathbf{I} \delta^4(x - y).$$

Form for $\Delta_A(x, y)$ may be obtained in a similar manner. Thus we obtain for $\hat{\Delta}_{diag}(X; P)$, the counterpart to the Fourier transform of $\hat{\Delta}_{diag}(u - v)$ in (4.31),

$$\hat{\Delta}_{diag}(X; P) \simeq \hat{\Delta}_{diag}^{(0)}(X; P) + \hat{\Delta}_{diag}^{(1)}(X; P) \quad (6.32)$$

$$\begin{aligned}\hat{\Delta}_{diag}^{(0)}(X; P) &= \text{diag} \left(\Delta_R^{(0)}(X; P), -\Delta_A^{(0)}(X; P) \right), \\ &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \text{diag} \left(\Delta_R^{(\xi)}(X; P), -\Delta_A^{(\xi)}(X; P) \right) \\ &\equiv \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \hat{\Delta}_{diag}^{(\xi)}(X; P),\end{aligned}\quad (6.33)$$

$$\hat{\Delta}_{diag}^{(1)}(X; P) = \text{diag} \left(\Delta_R^{(1)}(X; P), -\Delta_A^{(1)}(X; P) \right), \quad (6.34)$$

where

$$\begin{aligned}\Delta_{R(A)}^{(\xi)}(X; P) &\equiv \frac{1}{P^2 - \mathcal{M}_\xi^2(X) \pm i\epsilon(p_0)\eta}, \\ \Delta_{R(A)}^{(1)}(X; P) &= -iP_\mu \left(|\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)| \right) \\ &\quad \times \left[\frac{1}{P^2 - \mathcal{M}_\pi^2(X) \pm i\epsilon(p_0)\eta} - \frac{1}{P^2 - \mathcal{M}_\sigma^2(X) \pm i\epsilon(p_0)\eta} \right]^2.\end{aligned}\quad (6.35)$$

$\hat{\Delta}_{diag}(X; P)$ corresponds to $\Delta_{R(A)}(P)$, Eq. (4.6), in the complex-scalar field theory.

The following form for $\hat{\Delta}_{diag}(x, y)$ is also useful for later analysis:

$$\begin{aligned}\hat{\Delta}_{diag}(x, y) &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(x) \hat{\Delta}_{diag}^{(\xi)}(x, y) \mathbf{P}_\xi(y) + \hat{\Delta}_{diag}^{(1)'}(x, y), \\ \hat{\Delta}_{diag}^{(\xi)}(x, y) &= \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} \hat{\Delta}_{diag}^{(\xi)}(X; P), \\ \hat{\Delta}_{diag}^{(1)'}(x, y) &= 2i \int \frac{d^4 P}{(2\pi)^4} e^{-iP \cdot (x-y)} P_\mu \left(|\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)| \right) \\ &\quad \times \text{diag} \left(\Delta_R^{(\pi)}(X; P) \Delta_R^{(\sigma)}(X; P), -\Delta_A^{(\pi)}(X; P) \Delta_A^{(\sigma)}(X; P) \right).\end{aligned}\quad (6.36)$$

Let us find the property of the operators $|\varphi(X)\rangle \partial_{X_\mu} \langle \varphi(X)|$ and $|\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)|$. As defined above, $|\varphi(X)\rangle$ is the unit vector along $\vec{\varphi}(X)$ in the $O(N)$ vector space, so that $\langle \varphi(X) | \varphi(X) \rangle = 1$. Then $\langle \varphi(X) | \partial_{X_\mu} \varphi(X) \rangle = \langle \varphi(X) | \overleftrightarrow{\partial}_{X_\mu} \varphi(X) \rangle = 0$. From these relations, we obtain

$$\begin{aligned}\mathbf{P}_\xi(X) \left(|\varphi(X)\rangle \partial_{X_\mu} \langle \varphi(X)| \right) \mathbf{P}_\zeta(X) &= \begin{cases} |\varphi(X)\rangle \partial_{X_\mu} \langle \varphi(X)| & \text{if } \xi = \sigma \text{ and } \zeta = \pi, \\ 0 & \text{otherwise,} \end{cases} \\ \mathbf{P}_\xi(X) \left(|\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)| \right) \mathbf{P}_\zeta(X) &= \begin{cases} |\varphi(X)\rangle \overleftrightarrow{\partial}_{X_\mu} \langle \varphi(X)| & \text{if } \xi = \pi \text{ and } \zeta = \sigma, \\ 0 & \text{otherwise.} \end{cases}\end{aligned}\quad (6.37)$$

Thus, $\hat{\Delta}_{diag}^{(1)}$ in (6.32) and $\hat{\Delta}_{diag}^{(1)'}$ in (6.36) represent mixing between the π sector and the σ sector.

2. 2×2 matrix propagator

Let us introduce ‘‘Bogoliubov’’ transformation (cf. (4.15)),

$$\begin{aligned}\hat{\vec{\phi}}(x) &= \int d^3u \sum_{\tau=\pm} \hat{\mathbf{B}}_L^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) \hat{\vec{\phi}}_{RA}^{(\tau)}(\mathbf{u}, x_0), \\ \hat{\vec{\phi}}(y) &= \int d^3v \sum_{\tau=\pm} \hat{\vec{\phi}}_{RA}^{(\tau)}(\mathbf{v}, y_0) \hat{\mathbf{B}}_R^{(\tau)}(\mathbf{v}, \mathbf{y}; y_0).\end{aligned}$$

As seen above after (6.6), $\mathcal{L}_q$ enjoys $O(N-1)$ symmetry. We assume that the density matrix enjoys $O(N-1)$ symmetry in the sense that the $N \times N$ matrices $\hat{\mathbf{B}}_{L(R)}^{(\tau)}$ take the form,

$$\hat{\mathbf{B}}_{L(R)}^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) = \sum_{\xi, \zeta=\pi, \sigma} \mathbf{P}_\xi(\mathbf{x}, x_0) \hat{B}_{L(R)}^{(\tau)(\xi, \zeta)}(\mathbf{x}, \mathbf{u}; x_0) \mathbf{P}_\zeta(\mathbf{u}, x_0).$$

The condition that the canonical commutation relations are preserved (cf. (4.13)) yields a solution,

$$\hat{\mathbf{B}}_{L(R)}^{(\tau)}(\mathbf{x}, \mathbf{u}; x_0) = \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(\mathbf{x}, x_0) \hat{B}_{L(R)}^{(\tau)(\xi)}(\mathbf{x}, \mathbf{u}; x_0) \mathbf{P}_\xi(\mathbf{u}, x_0),$$

where $\hat{B}_{L(R)}^{(\tau)(\xi)}$ is as in (4.16) with $f^{(\tau)} \rightarrow f_\xi^{(\tau)}$. Translated into the $|p_0|$ prescription (cf. (4.31)), we obtain for the propagator,

$$\begin{aligned}\hat{\Delta}(x, y) &= \int d^4u d^4v \hat{\mathbf{B}}_L(x, u) \hat{\Delta}_{diag}(u, v) \hat{\mathbf{B}}_R(v, y) \\ &= \int d^4u d^4v \sum_{\xi, \zeta=\pi, \sigma} \mathbf{P}_\xi(x) \hat{B}_L^{(\xi)}(x, u) \mathbf{P}_\xi(u) \hat{\Delta}_{diag}(u, v) \mathbf{P}_\zeta(v) \hat{B}_R^{(\zeta)}(v, y) \mathbf{P}_\zeta(y), \\ \hat{B}_L^{(\xi)}(x, u) &= \begin{pmatrix} \delta^4(x-u) & f_\xi(x, u) \\ \delta^4(x-u) & \delta^4(x-u) + f_\xi(x, u) \end{pmatrix}, \\ \hat{B}_R^{(\xi)}(v, y) &= \begin{pmatrix} \delta^4(v-y) + f_\xi(v, y) & f_\xi(v, y) \\ \delta^4(v-y) & \delta^4(v-y) \end{pmatrix}.\end{aligned}\tag{6.38}$$

Substituting (6.36) into (6.38), we obtain

$$\begin{aligned}\hat{\Delta}(x, y) &= \int d^4u d^4v \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(x) \hat{B}_L^{(\xi)}(x, u) \mathbf{P}_\xi(u) \hat{\Delta}_{diag}^{(\xi)}(u, v) \mathbf{P}_\xi(v) \hat{B}_R^{(\xi)}(v, y) \mathbf{P}_\xi(y) \\ &\quad + \int d^4u d^4v \sum_{\xi, \zeta=\pi, \sigma} \mathbf{P}_\xi(x) \hat{B}_L^{(\xi)}(x, u) \mathbf{P}_\xi(u) \hat{\Delta}_{diag}^{(1)'}(u, v) \mathbf{P}_\zeta(v) \hat{B}_R^{(\zeta)}(v, y) \mathbf{P}_\zeta(y).\end{aligned}\tag{6.39}$$

From the first term on the RHS, we pick out

$$\mathbf{P}_\xi(x) \mathbf{P}_\xi(u) \mathbf{P}_\xi(v) \mathbf{P}_\xi(y).\tag{6.40}$$

Using $\mathbf{P}_\xi(u) \simeq \mathbf{P}_\xi(x) + (u-x)_\mu \partial_{X_\mu} \mathbf{P}_\xi(x)$, etc., we obtain

$$\begin{aligned}\text{Eq. (6.40)} &= \mathbf{P}_\xi(x) \mathbf{P}_\xi(y) + (u-x)_\mu \mathbf{P}_\xi(x) (\partial_{x_\mu} \mathbf{P}_\xi(x)) \mathbf{P}_\xi(y) \\ &\quad + (v-y)_\mu \mathbf{P}_\xi(y) (\partial_{y_\mu} \mathbf{P}_\xi(y)) \mathbf{P}_\xi(x).\end{aligned}$$

In the second (third) term on the RHS, we have set $\mathbf{P}_\xi(v) \simeq \mathbf{P}_\xi(y) \simeq \mathbf{P}_\xi(x)$ ($\mathbf{P}_\xi(x) \simeq \mathbf{P}_\xi(u) \simeq \mathbf{P}_\xi(y)$), which is allowed since the term already contains the X_μ -derivative. The relations (6.37) tell us that the second and third terms on the RHS vanish. Thus, we obtain, for the first term on the RHS of (6.39),

$$\sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(x) \left[\int d^4u d^4v \hat{B}_L^{(\xi)}(x, u) \hat{\Delta}_{diag}^{(\xi)}(u, v) \hat{B}_R^{(\xi)}(v, y) \right] \mathbf{P}_\xi(y).\tag{6.41}$$

The quantity in the square brackets is the same as in complex-scalar theory dealt with in Section IV. Fourier transformation of (6.41) may be carried out using (6.38), (6.36) and (C2) - (C4). The last term in (6.39) may also be computed by using (6.36) and (6.37). Fourier transforming the resultant expression, we have

$$\begin{aligned} & -2iP_\mu \frac{\partial |\varphi(X)\rangle}{\partial X_\mu} \langle \varphi(X) | \left[\begin{pmatrix} \Delta_R^\pi \Delta_R^\sigma & 0 \\ \Delta_R^\pi \Delta_R^\sigma - \Delta_A^\pi \Delta_A^\sigma & -\Delta_A^\pi \Delta_A^\sigma \end{pmatrix} + (f_\sigma \Delta_R^\pi \Delta_R^\sigma - f_\pi \Delta_A^\pi \Delta_A^\sigma) \hat{A}_+ \right] \\ & + 2iP_\mu |\varphi(X)\rangle \frac{\partial \langle \varphi(X) |}{\partial X_\mu} [\pi \longleftrightarrow \sigma], \end{aligned}$$

where $\Delta_R^\pi = \Delta_R^\pi(X; P)$, $f_\sigma = f_\sigma(X)$, etc.

After all this, we finally obtain for the Fourier transform of $\hat{\Delta}(x, y)$ on $x - y$:

$$\hat{\Delta}(X; P) = \hat{\Delta}^{(0)}(X; P) + \hat{\Delta}^{(1)}(X; P), \quad (6.42)$$

$$\begin{aligned} \hat{\Delta}^{(0)}(X; P) &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \hat{B}_L^{(\xi)}(X; P) \hat{\Delta}_{diag}^{(\xi)}(X; P) \hat{B}_R^{(\xi)}(X; P) \\ &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \hat{\Delta}^{(\xi)}(X; P), \end{aligned} \quad (6.43)$$

$$\begin{aligned} \hat{B}_L^{(\xi)}(X; P) &= \begin{pmatrix} 1 & f_\xi(X; P) \\ 1 & 1 + f_\xi(X; P) \end{pmatrix}, \quad \hat{B}_R^{(\xi)}(X; P) = \begin{pmatrix} 1 + f_\xi(X; P) & f_\xi(X; P) \\ 1 & 1 \end{pmatrix}, \\ \hat{\Delta}^{(\xi)}(X; P) &= \hat{\Delta}_{RA}^{(\xi)}(X; P) + \hat{\Delta}^{(\xi)'}(X; P), \end{aligned} \quad (6.44)$$

where

$$\hat{\Delta}_{RA}^{(\xi)}(X; P) = \begin{pmatrix} \Delta_R^{(\xi)} & 0 \\ \Delta_R^{(\xi)} - \Delta_A^{(\xi)} & -\Delta_A^{(\xi)} \end{pmatrix}, \quad (6.45)$$

$$\Delta_{R(A)}^{(\xi)}(X; P) = \frac{1}{P^2 - \mathcal{M}_\xi^2(X) \pm i\epsilon(p_0)\eta}, \quad (6.46)$$

$$\hat{\Delta}^{(\xi)'}(X; P) = f_\xi(X; P) [\Delta_R^{(\xi)}(X; P) - \Delta_A^{(\xi)}(X; P)] \hat{A}_+, \quad (6.47)$$

$$\begin{aligned} \hat{\Delta}^{(1)}(X; P) &= -\frac{i}{2} \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \{f_\xi(X; P), P^2 - \mathcal{M}_\xi^2(X)\} [(\Delta_R^{(\xi)})^2 + (\Delta_A^{(\xi)})^2] \hat{A}_+ \\ &\quad + iP^\mu \frac{\partial |\varphi(X)\rangle}{\partial X^\mu} \langle \varphi(X) | \hat{\Omega} - iP^\mu |\varphi(X)\rangle \frac{\partial \langle \varphi(X) |}{\partial X^\mu} \hat{\Omega} \Big|_{\pi \leftrightarrow \sigma} \\ &\quad + \frac{i}{2} \left[|\varphi(X)\rangle \frac{\leftrightarrow}{\partial X_\mu} \langle \varphi(X) | \right] \sum_{\xi=\pi, \sigma} \frac{\partial f_\xi(X; P)}{\partial P_\mu} (\Delta_R^{(\xi)} - \Delta_A^{(\xi)}) \hat{A}_+. \end{aligned} \quad (6.48)$$

Here

$$\hat{\Omega} = \hat{\Omega}_0 + \left(f_\sigma [(\Delta_R^{(\sigma)})^2 - 2\Delta_R^{(\pi)} \Delta_R^{(\sigma)} - (\Delta_A^{(\sigma)})^2] - f_\pi [(\Delta_A^{(\pi)})^2 - 2\Delta_A^{(\pi)} \Delta_A^{(\sigma)} - (\Delta_R^{(\pi)})^2] \right) \hat{A}_+, \quad (6.49)$$

$$\hat{\Omega}_0 = \begin{pmatrix} (\Delta_R^{(\pi)} - \Delta_R^{(\sigma)})^2 & 0 \\ (\Delta_R^{(\pi)} - \Delta_R^{(\sigma)})^2 - (\Delta_A^{(\pi)} - \Delta_A^{(\sigma)})^2 & -(\Delta_A^{(\pi)} - \Delta_A^{(\sigma)})^2 \end{pmatrix}. \quad (6.50)$$

Note that f_ξ is related to the “genuine” number density $N^{(\xi)}$ through (cf. (4.30))

$$f_\xi(X; P) = \theta(p_0) N^{(\xi)}(X; E_p^{(\xi)}; \hat{\mathbf{p}}) - (1 + \theta(-p_0)) N^{(\xi)}(X; E_p^{(\xi)}; -\hat{\mathbf{p}}).$$

3. Counter Lagrangian and renormalized scheme

Following the procedure in Section VE (cf. (5.54) to (5.55)), we can extract the counter hat-Lagrangian density $\hat{\mathcal{L}}_c$ through analyzing $\hat{\Delta}^{-1}(x) \hat{\Delta}(x, y)$ and $\hat{\Delta}(x, y) \hat{\Delta}^{-1}(y)$. $\hat{\mathcal{L}}_c$ thus found gives a two-point vertex function $\hat{\mathbf{V}}_c(X; P)$. Straightforward manipulation yields

$$i\hat{\mathbf{V}}_c(X;P) \simeq \sum_{\xi=\pi,\sigma} \{ \mathbf{P}_\xi(X) f_\xi(X;P), P^2 - \mathcal{M}_\xi^2(X) \} \hat{A}_- \quad (6.51)$$

$$= \sum_{\xi=\pi,\sigma} \mathbf{P}_\xi(X) \{ f_\xi(X;P), P^2 - \mathcal{M}_\xi^2(X) \} \hat{A}_- \\ - 2P_\mu [f_\pi(X;P) - f_\sigma(X;P)] \frac{\partial |\varphi(X)\rangle \langle \varphi(X)|}{\partial X_\mu} \hat{A}_-. \quad (6.52)$$

Let us adopt the physical- f scheme. In the case of complex-scalar field, we have adopted the “renormalization condition” $\hat{\mathbf{V}}_c(X;P) = 0$ (cf. (5.55) and (5.56)). In the present case, we cannot set $\hat{\mathbf{V}}_c(X;P) = 0$. Then, we adopt the condition,

$$\{ f_\xi(X;P), P^2 - \mathcal{M}_\xi^2(X) \} = 0, \quad (6.53)$$

which is the same as in the complex-scalar-field case. Then, in $\hat{\mathbf{V}}_c$ in (6.52), the second term on the RHS survives. Let us compute the $\hat{\mathbf{V}}_c$ -resummed propagator $\hat{\Delta}_{c-res}$. Since $\hat{\mathbf{V}}_c$ contains the X_μ -derivative, one can directly write down the Fourier-transformed form:

$$\hat{\Delta}(X;P) \left[1 + \sum_{n=1} \left\{ -\hat{\mathbf{V}}_c(X;P) \hat{\Delta}(X;P) \right\}^n \right]. \quad (6.54)$$

In 2×2 matrix space, $\hat{\mathbf{V}}_c$ is proportional to $\hat{A}_-$, Eq. (3.16), which shows that, in (6.54), the contributions with $n \geq 2$ vanish. Thus, after some manipulations, we obtain,

$$\hat{\Delta}_{c-resum}(X;P) \simeq -2i(f_\pi - f_\sigma)P_\mu \\ \times \left[|\varphi(X)\rangle \overleftarrow{\partial}_{X_\mu} \langle \varphi(X)| \Delta_R^\pi \Delta_A^\sigma + |\varphi(X)\rangle \partial_{X_\mu} \langle \varphi(X)| \Delta_R^\sigma \Delta_A^\pi \right] \hat{A}_+.$$

Summing this to $\hat{\Delta}(X;P)$ in (6.42), we obtain (6.42) - (6.48) with (6.53), provided that $\hat{\Omega}$ in (6.49) is changed to

$$\hat{\Omega} \rightarrow \hat{\Omega}_{renorm} = \hat{\Omega}_0 + \left(f_\sigma [\Delta_R^{(\sigma)} - \Delta_A^{(\sigma)}] [\Delta_R^{(\sigma)} + \Delta_A^{(\sigma)} - 2\Delta_R^{(\pi)}] \right. \\ \left. - f_\pi [\Delta_A^{(\pi)} - \Delta_R^{(\pi)}] [\Delta_A^{(\pi)} + \Delta_R^{(\pi)} - 2\Delta_A^{(\sigma)}] \right) \hat{A}_+. \quad (6.55)$$

This completes the deduction of bare propagators in the bare- f scheme. The physical- f scheme may be constructed in a similar manner as in Section IVC. In place of (5.27) or (5.29), we have, with obvious notation,

$$\Sigma_{off}^{(\xi)}(X;P) = 0 \quad (\xi = \pi, \sigma), \quad (6.56)$$

which lead to the generalized Boltzmann equations.

D. Determining $M_\pi^2(X)$ and $M_\sigma^2(X)$

In this subsection, we present a scheme for determining so far arbitrary $M_\xi^2(X)$ ($\xi = \pi, \sigma$), introduced in (6.7) - (6.11). Various methods of fixing $M_\xi^2(X)$ are available. Among those, we mention the following three methods.

1) *Resummation method.* Argument in Section V shows that the quantities to be resummed are the real part of the retarded and advanced self-energy parts,

$$\Sigma_R(X;R) = \Sigma_{11}(X;R) + \Sigma_{12}(X;R) = (\Sigma_A(X;R))^*.$$

Determine $M_\xi^2(X)$ ($\xi = \pi, \sigma$) such that the real part of $\Sigma_R(X;R)$ computed up to required L th loop order, $Re \hat{\Sigma}_R^{(L)}(X;R)$, vanishes at $R^\mu = R_0^\mu$ with R_0^μ a suitable four momentum:

$$Re \hat{\Sigma}_R^{(L)}(X;R = R_0) = 0. \quad (6.57)$$

2) *The principles of minimal sensitivity* [35]. Total Lagrangian density $\mathcal{L}$ in (6.8) is independent of arbitrary functions $M_\pi^2(X)$ and $M_\sigma^2(X)$. Let Ξ be a quantity, in which we are interested. Then, if we could evaluate Ξ to

all orders of perturbation series or loop expansion, it is independent of $M_\xi^2(X)$ ($\xi = \pi, \sigma$). On the other hand, Ξ computed on the basis of $\mathcal{L}$ in (6.8) up to the L th order, $\Xi^{(L)}$, depends on $M_\xi^2(X)$. This observation leads one to *impose* the condition

$$\frac{\delta \Xi^{(L)}}{\delta M_\xi^2(X)} = 0 \quad (\xi = \pi, \sigma),$$

which serves as a determining equation for $M_\xi^2(X)$. Note that, for $\Xi = \Gamma_{phys}$ or $\Xi = \delta \Gamma_{phys} / \delta \vec{\varphi}(X)$, this method works for $L \geq 2$.

3) *The criterion of the fastest apparent convergence* [36]. Compute Ξ up to L th order, $\Xi^{(L)}$ ($L \geq 2$). Then, determine all the parameters of the theory, such as the coupling constant, mass functions $M_\xi^2(X)$ ($\xi = \pi, \sigma$), etc., so that all the terms but the first term of the series vanish.

A number of works, which employ the methods 2) and 3) above for analyzing the equilibrium effective potential, is cited in [10].

In this paper, we follow the method 1). The requirement in 1) gives the gap equation.

E. Summary of this section

In this section, we have found that the spacetime evolution of the condensate fields $\vec{\varphi}(X)$ is governed by the following system of coupled equations:

E1) Generalized relativistic kinetic or Boltzmann equation (6.56).

E2) Gap equation (6.57).

E3) Equation of motion for $\vec{\varphi}(X)$, (6.24) with $j = 1$.

As seen in Section VC, (6.53) in E1) is a determining equation for the number-density function $f_\xi(X; P)$ ($\xi = \pi, \sigma$) under a given initial data $f_\xi(X_0 = T_{in}, \mathbf{X}; P)$. This equation involves yet unknown functions $M_\xi^2(X)$ ($\xi = \pi, \sigma$). Equation (6.57) determines $M_\xi^2(X)$ in terms of $f_\xi(X; P)$. Thus, (6.53) and (6.57) interplay to determine $M_\xi^2(X)$ and $f_\xi(X; P)$. Through computing the effective action $\hat{\Gamma}$ using $M_\xi^2(X)$ and $f_\xi(X; P)$ thus determined, the equation of motion (6.25) for $\vec{\varphi}(X)$ is obtained. Solving it under a given initial data determines the spacetime evolution of $\vec{\varphi}(X)$, which describes the progress of phase transition.

VII. LOWEST-ORDER EFFECTIVE ACTION IN THE σ MODEL

In this section, we compute the CTP effective action to the lowest nontrivial order, which comes³ from an one-loop diagram with $\hat{\mathcal{L}}_0$ as in (6.9). The contribution is

$$\hat{\Gamma} = \frac{i}{2} \text{Tr} \ln \left(\frac{1}{2} \hat{\tau}_3 \left\{ \mathbf{I} + \sum_{\xi=\pi, \sigma} \hat{\mathbf{P}}_\xi \hat{\mathcal{M}}_\xi^2 \partial^{-2} \right\} \right), \quad (7.1)$$

where ‘Tr’ acts on matrices in spacetime, 2×2 matrices in the “complex-time plane,” and $N \times N$ matrices in the $O(N)$ vector space. In (7.1),

$$(\hat{\tau}_3 \partial^{-2})(x, y) = -\hat{\Delta}(x, y) \Big|_{m=0},$$

with $\hat{\Delta}(x, y)$ as in (4.31) and

³The difference between $\hat{\Gamma}$ in (7.1) and $\hat{\Gamma}$ computed using the physical- f scheme, whose form has not been displayed explicitly, is of higher order.

$$\begin{aligned}
\hat{\mathcal{M}}_\xi^2(x) &= \text{diag} [\mathcal{M}_{\xi 1}^2(x), -\mathcal{M}_{\xi 2}^2(x)] \quad (\xi = \pi, \sigma) \\
\hat{\vec{\varphi}}^2 &= \text{diag} (\vec{\varphi}_1^2(x), -\vec{\varphi}_2^2(x)), \\
\hat{\mathbf{P}}_\xi(x) &= \text{diag} (\mathbf{P}_{\xi 1}(x), -\mathbf{P}_{\xi 2}(x)).
\end{aligned}$$

Here $\vec{\varphi}_i$ ($i = 1, 2$) are the type- i condensate fields, $\mathcal{M}_{\xi i}^2$ and $\mathbf{P}_{\xi i}$ ($i = 1, 2$) are as in (6.29) and (6.11), respectively, with $\vec{\varphi}_i$ in place of $\vec{\varphi}$.

For extracting the *physical* effective action from (7.1), we use a method [32] of computing *physical* effective potential, in which the tadpole method is employed. The method starts with taking the derivatives,

$$\frac{\delta \hat{\Gamma}}{\delta \mathcal{M}_{\xi 1}^2(x)} = \frac{i}{2} \text{tr} \mathbf{P}_{\xi 1}(x) \left[\mathbf{I} \partial^{-2}(x, x) - \left\{ \partial^{-2} \left(\hat{\mathbf{P}}_\pi \hat{\mathcal{M}}_\pi^2 + \hat{\mathbf{P}}_\sigma \hat{\mathcal{M}}_\sigma^2 \right) \partial^{-2} \right\} (x, x) + \dots \right]_{11}$$

($\xi = \pi, \sigma$), (7.2)

where “tr” acts on 2×2 matrices and $N \times N$ matrices. Then, we set $\vec{\varphi}_1^2(x) = \vec{\varphi}_2^2(x) \equiv \vec{\varphi}^2(x)$. Equation (7.2) now becomes (cf. (6.25))

$$\begin{aligned}
\left. \frac{\delta \hat{\Gamma}}{\delta \mathcal{M}_{\xi 1}^2(x)} \right|_{\vec{\varphi}_1 = \vec{\varphi}_2 = \vec{\varphi}} &= -\frac{i}{2} \text{tr} \mathbf{P}_\xi(x) \mathbf{G}_{11}(x, x) \\
&= \frac{\delta \Gamma_{phys}}{\delta \mathcal{M}_\xi^2(x)},
\end{aligned}$$

(7.3)

where

$$\hat{\mathbf{G}} = -\hat{\tau}_3 \left[\mathbf{I} \partial^2 + \hat{\mathbf{P}}_\pi \hat{\mathcal{M}}_\pi^2 + \hat{\mathbf{P}}_\sigma \hat{\mathcal{M}}_\sigma^2 \right]^{-1} \Big|_{\vec{\varphi}_1 = \vec{\varphi}_2 = \vec{\varphi}}.$$

(7.4)

The one-loop contribution to the *physical* effective action Γ_{phys} is obtained [32] by integrating (7.3) over $\mathcal{M}_\xi^2(x)$.

Incidentally, the “unphysical part” of Γ , which comes from $\delta \hat{\Gamma} / \delta \mathcal{M}_{\xi 2}^2(X)$, becomes

$$\Gamma_{unphys} = -[\Gamma_{phys}]^*.$$

(7.5)

So far, we have made no assumption on the x -dependence of $\mathcal{M}_\xi^2(x)$ and $\vec{\varphi}(x)$. Here we recall that $\mathcal{M}_\xi^2(x)$ and $\vec{\varphi}(x)$ weakly depend on x . We first note that

$$\text{Sp}[\partial_{X_\mu} \mathbf{P}_\xi(X)] = \text{Sp}[\mathbf{P}_\xi \partial_{X_\mu} \mathbf{P}_\zeta(X)] = 0 \quad (\xi, \zeta = \pi, \sigma),$$

where ‘Sp’ stands for the trace operation that acts on an $N \times N$ matrix. Then, the term with $\partial_{X_\mu} \mathbf{P}_\xi(X)$ does not appear. This means that, within the gradient approximation, we can assign common spacetime coordinates for all $\hat{\mathbf{P}}_\xi$ in (7.4). Thus, including the zeroth-order contribution (cf. (6.21)), we obtain

$$\Gamma_{phys} = \int d^4 X \gamma_{phys}(X),$$

(7.6)

$$\begin{aligned}
\gamma_{phys}(X) &= \frac{1}{2} [(\partial \vec{\varphi}(X))^2 - \mu^2 \vec{\varphi}^2(X)] - \frac{\lambda}{4!} (\vec{\varphi}^2(X))^2 + H({}^t \vec{e} \cdot \vec{\varphi}(X)) \\
&\quad + \frac{1}{2} [(N-1)\gamma_\pi(X) + \gamma_\sigma(X)],
\end{aligned}$$

(7.7)

$$\gamma_\xi(X) = - \int d\mathcal{M}_\xi^2(X) \mathcal{I}_\xi(X) \quad (\xi = \pi, \sigma).$$

(7.8)

Here

$$\mathcal{I}_\xi(X) = i \int \frac{d^4 P}{(2\pi)^4} \Delta_{11}^{(\xi)}(X; P) = \mathcal{I}_\xi^{(0)}(X) + \mathcal{I}_\xi^{(\beta)}(X),$$

(7.9)

$$\mathcal{I}_\xi^{(0)}(X) \equiv \int \frac{d^4 P}{(2\pi)^4} \frac{i}{P^2 - \mathcal{M}_\xi^2 + i\eta},$$

(7.10)

$$\mathcal{I}_\xi^{(\beta)}(X) \equiv \int \frac{d^3 p}{(2\pi)^3} \frac{N^{(\xi)}(X; \mathcal{E}_p^{(\xi)}(X), \hat{\mathbf{p}})}{\mathcal{E}_p^{(\xi)}(X)},$$

(7.11)

where $\mathcal{E}_p^{(\xi)}(X) = \sqrt{p^2 + \mathcal{M}_\xi^2(X)}$. Thus the physical effective action Γ_{phys} is given by an integral of the “local effective action” $\gamma_{phys}(X)$ over the macroscopic spacetime coordinates X^μ . Computation of $\gamma_{phys}(X)$ goes essentially in the same manner as that of the physical effective potential.

In the following two subsections, we separately analyze the positive-mass-square region and negative-mass-square region. As mentioned in Section I, our scheme does not apply to the region near the transition point between the two regions.

A. The region $\mathcal{M}_\pi^2(X), \mathcal{M}_\sigma^2(X) > 0$.

The lowest-order contribution to $\mathcal{I}_\xi(X)$, Eq. (7.9), has already been computed in Section VD (see (5.34) - (5.37)):

$$\begin{aligned}\mathcal{I}_\xi^{(0)}(X) &= -\frac{\mathcal{M}_\xi^2(X)}{(4\pi)^2} \left[\frac{2}{\bar{\epsilon}} - \ln \frac{\mathcal{M}_\xi^2(X)}{e\mu_d^2} \right] \\ \mathcal{I}_\xi^{(\beta)}(X) &= \frac{1}{2\pi^2} \int_{\mathcal{M}_\xi(X)}^\infty d\mathcal{E} \sqrt{\mathcal{E}^2 - \mathcal{M}_\xi^2(X)} \langle N^{(\xi)}(X; \mathcal{E}) \rangle,\end{aligned}\tag{7.12}$$

where

$$\langle N^{(\xi)}(X; \mathcal{E}) \rangle \equiv \int \frac{d\Omega_{\mathbf{p}}}{4\pi} N^{(\xi)}(X; \mathcal{E}, \mathbf{p})\tag{7.13}$$

with $d\Omega_{\mathbf{p}}$ a solid-angle element in a $\mathbf{p}$ -space. Substituting these into (7.8), we obtain

$$\gamma_\xi(X) = \gamma_\xi^{(0)}(X) + \gamma_\xi^{(\beta)}(X)\tag{7.14}$$

with

$$\gamma_\xi^{(0)}(X) = i \int \frac{d^4 P}{(2\pi)^4} \ln(P^2 - \mathcal{M}_\xi^2(X) + i\eta)\tag{7.15}$$

$$= \frac{1}{32\pi^2} \mathcal{M}_\xi^4(X) \left[\frac{2}{\bar{\epsilon}} + \frac{3}{2} - \ln \frac{\mathcal{M}_\xi^2(X)}{\mu_d^2} \right],\tag{7.16}$$

$$\gamma_\xi^{(\beta)}(X) = \frac{1}{3\pi^2} \int_{\mathcal{M}_\xi(X)}^\infty d\mathcal{E} (\mathcal{E}^2 - \mathcal{M}_\xi^2(X))^{3/2} \langle N^{(\xi)}(X; \mathcal{E}) \rangle.\tag{7.17}$$

$2/\bar{\epsilon}$ in (7.16) is canceled by the counter Lagrangian $\mathcal{L}'_{cond}$, Eq. (6.5). We note that, to the present one-loop order, we can replace $-(Z_\mu Z - 1)\mu^2 \vec{\varphi}^2/2$ in (6.5) as

$$-(Z_\mu Z - 1)\mu^2 \vec{\varphi}^2/2 \rightarrow -\frac{1}{2(N+2)}(Z_\mu Z - 1) [(N-1)M_\pi^2 + 3M_\sigma^2] \vec{\varphi}^2.\tag{7.18}$$

The difference between the LHS and the RHS, which is proportional to the UV-diverging factor $1/\bar{\epsilon}$, is of higher order (cf. above after (6.11)). This difference is canceled by the higher-order one-loop diagrams, one of which includes the vertex coming from ${}^t\vec{\phi}\chi_\pi \mathbf{P}_\pi \vec{\phi}/2$ ($\in \mathcal{L}_{rc}$) and the other one includes the vertex coming from ${}^t\vec{\phi}\chi_\sigma \mathbf{P}_\sigma \vec{\phi}/2$ ($\in \mathcal{L}_{rc}$). Since we are choosing the $\overline{\text{MS}}$ scheme, we have

$$Z = 1, \quad Z_\mu = 1 + \lambda \frac{N+2}{48\pi^2} \frac{1}{\bar{\epsilon}}, \quad Z_\lambda = 1 + \lambda \frac{N+8}{48\pi^2} \frac{1}{\bar{\epsilon}}.\tag{7.19}$$

In a similar manner, we obtain for A in (6.5),

$$A = -\frac{N}{32\pi^2} \frac{1}{\bar{\epsilon}}.$$

After all this, we obtain for the local effective action, to one-loop order,

$$\gamma_{phys}(X) = \gamma_{phys}^{(0)}(X) + \gamma_{phys}^{(\beta)}(X),\tag{7.20}$$

where

$$\begin{aligned} \gamma_{phys}^{(0)}(X) &= \frac{1}{2} [(\partial\vec{\varphi}(X))^2 - \mu^2\vec{\varphi}^2(X)] - \frac{\lambda}{4!}(\vec{\varphi}^2(X))^2 + H({}^t\vec{e} \cdot \vec{\varphi}(X)) \\ &\quad - \frac{1}{(8\pi)^2} \left[(N-1)\mathcal{M}_\pi^4(X) \ln \frac{\mathcal{M}_\pi^2(X)}{e^{3/2}\mu_d^2} + \mathcal{M}_\sigma^4(X) \ln \frac{\mathcal{M}_\sigma^2(X)}{e^{3/2}\mu_d^2} \right], \end{aligned} \quad (7.21)$$

$$\gamma_{phys}^{(\beta)}(X) = \frac{1}{2} \left[(N-1)\gamma_\pi^{(\beta)}(X) + \gamma_\sigma^{(\beta)}(X) \right]. \quad (7.22)$$

It should be emphasized that the factor of the “ $\vec{\varphi}^2$ term” in (7.21) is neither $\mathcal{M}_\pi^2(X)$ nor $\mathcal{M}_\sigma^2(X)$ but μ^2 .

In the case of thermal equilibrium system with constant $\vec{\varphi}$, $-\gamma_{phys}$ reduces to the effective potential (density) of the system, $V(\vec{\varphi}) = -\gamma_{phys}(\vec{\varphi})$. Besides the terms due to the different UV-renormalization scheme, $V(\vec{\varphi})$ obtained above agrees with the result of [10], where $\mathcal{M}_\pi^2(X) = \mathcal{M}_\sigma^2(X)$, being constant, is assumed.

Now we are ready to write down the equation of motion for the condensate fields (cf. Section VIB),

$$\frac{\delta\Gamma_{phys}}{\delta\vec{\varphi}(X)} = -[\partial^2 + \Omega(\vec{\varphi}^2(X))] \vec{\varphi}(X) + H\vec{e} = 0, \quad (7.23)$$

$$\Omega(\vec{\varphi}^2(X)) = -2\frac{\partial\gamma_{phys}(X)}{\partial\vec{\varphi}^2(X)} \quad (7.24)$$

$$= \mu^2 + \frac{\lambda}{6}\vec{\varphi}^2(X) + \frac{\lambda}{6}[(N-1)\mathcal{I}_\pi(X) + 3\mathcal{I}_\sigma(X)] \quad (7.25)$$

$$= \Omega_0(\vec{\varphi}^2(X)) + \Omega_\beta(\vec{\varphi}^2(X)), \quad (7.26)$$

where

$$\begin{aligned} \Omega_0(\vec{\varphi}^2(X)) &= \mu^2 + \frac{\lambda}{6}\vec{\varphi}^2(X) \\ &\quad + \frac{\lambda}{96\pi^2} \left[(N-1)\mathcal{M}_\pi^2(X) \ln \frac{\mathcal{M}_\pi^2(X)}{e\mu_d^2} + 3\mathcal{M}_\sigma^2(X) \ln \frac{\mathcal{M}_\sigma^2(X)}{e\mu_d^2} \right], \end{aligned} \quad (7.27)$$

$$\Omega_\beta(\vec{\varphi}^2(X)) = \frac{\lambda}{6} \left[(N-1)\mathcal{I}_\pi^{(\beta)}(X) + 3\mathcal{I}_\sigma^{(\beta)}(X) \right]. \quad (7.28)$$

Here $\mathcal{I}_\xi^{(\beta)}(X)$ ($\xi = \pi, \sigma$) is as in (7.12).

It is worth mentioning in passing that the equation of motion for unphysical or type-2 condensate fields are obtained from (7.5), which is of the same form as (7.23). Since the time argument of the type-2 field is on C_2 , the type-2 field “lives” in the “time-reversed world.” Then, one can say that the equation of motion is time-reversal invariant.

Since we are assuming that $\vec{\varphi}(X)$ does not appreciably change in a single spacetime cell (cf. Section I), (7.23) applies only in the region of $\vec{\varphi}(X)$ where $|\Omega(\vec{\varphi}^2(X))| \lesssim 1/(L^\mu)^2$ ($\mu = 0, 1, 2, 3$) with L^μ the size of the spacetime cell.

Let us study the small oscillation of $\vec{\varphi}(X)$ at the vicinity of the quasiminimum point $\vec{\varphi}(X) = \varphi_0(X)\vec{e}$:

$$\Omega(\varphi_0^2(X))\varphi_0(X) = H. \quad (7.29)$$

Setting $\vec{\varphi}(X) = \varphi_0(X)\vec{e} + \vec{\epsilon}(X)$, we obtain,

$$\left[\partial^2 + \Omega^{(\sigma)}(\vec{\varphi}_0^2(X)) + \frac{H}{\varphi_0(X)} \right] ({}^t\vec{e} \cdot \vec{\epsilon}(X)) + \partial^2\varphi_0(X) \simeq 0, \quad (7.30)$$

$$\left[\partial^2 + \frac{h}{\varphi_0(X)} \right] \vec{\epsilon}_\perp(X) \simeq 0, \quad (7.31)$$

where $\vec{\epsilon}_\perp(X) \equiv \vec{\epsilon}(X) - ({}^t\vec{e} \cdot \vec{\epsilon}(X))\vec{e}$ and

$$\Omega^{(\sigma)}(\vec{\varphi}_0^2(X)) = \Omega_0^{(\sigma)}(\vec{\varphi}_0^2(X)) + \Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X))$$

with

$$\Omega_0^{(\sigma)}(\vec{\varphi}_0^2(X)) = \frac{\lambda\varphi_0^2(X)}{3} \left[1 + \frac{\lambda}{96\pi^2} \left\{ (N-1) \ln \frac{\mathcal{M}_\pi^2(X)}{\mu_d^2} + 9 \ln \frac{\mathcal{M}_\sigma^2(X)}{\mu_d^2} \right\} \right]_{\vec{\varphi}(X)=\varphi_0(X)\vec{e}}, \quad (7.32)$$

$$\Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X)) = \frac{\lambda^2\varphi_0^2(X)}{18} \left[(N-1) \frac{\partial\mathcal{I}_\pi(X)}{\partial\mathcal{M}_\pi^2(X)} + 9 \frac{\partial\mathcal{I}_\sigma(X)}{\partial\mathcal{M}_\sigma^2(X)} \right]_{\vec{\varphi}(X)=\varphi_0(X)\vec{e}}. \quad (7.33)$$

$\vec{\epsilon}_\perp(X)$ describes quasi Goldstone mode. Equation (7.30) may be used only in the region of $\varphi_0(X)$ where $|\Omega^{(\sigma)}(\vec{\varphi}_0^2(X))| \lesssim 1/(L^\mu)^2$.

1. “High-temperature” expansion

The system is characterized by a scale parameter or several scale parameters of mass dimension. Let us assume that, for small $\mathcal{E}$, $\langle N^{(\xi)}(X; \mathcal{E}) \rangle$ in (7.13) behaves as

$$\langle N^{(\xi)}(X; \mathcal{E}) \rangle \simeq \frac{\mathcal{T}_\xi(X)}{\mathcal{E}} + \mathcal{C}_\xi(X) \quad \text{for } \mathcal{T}_\xi(X) \gg \mathcal{E}, \quad (7.34)$$

where $\mathcal{T}_\xi(X)$, which we call the temperature, is proportional to the scale parameter or to some combination of the scale parameters. In the case of locally thermal equilibrium system, the averaged number-density function reads

$$\langle N^{(\xi)}(X; \mathcal{E}) \rangle = \frac{1}{e^{\mathcal{E}/T(X)} - 1}, \quad (7.35)$$

where $T(X)$ is the local temperature of the system. Then, we have

$$\mathcal{T}_\xi(X) = T(X), \quad \mathcal{C}_\xi(X) = -\frac{1}{2}.$$

Let us analyze $\gamma_{phys}(X)$ in the high-temperature region. We start with rewriting (7.17) as

$$\begin{aligned} \gamma_\xi^{(\beta)}(X) &= \gamma_{\xi 1}^{(\beta)}(X) + \gamma_{\xi 2}^{(\beta)}(X) \\ \gamma_{\xi 1}^{(\beta)}(X) &= \frac{1}{3\pi^2} \int_0^\infty d\mathcal{E} \left[\mathcal{E}^3 - \frac{3}{2} \mathcal{E} \mathcal{M}_\xi^2(X) \right] \langle N^{(\xi)}(X; \mathcal{E}) \rangle \\ &\equiv \frac{\pi^2}{45} b_\xi(X) \mathcal{T}_\xi^4(X) - \frac{1}{12} c_\xi(X) \mathcal{M}_\xi^2(X) \mathcal{T}_\xi^2(X), \end{aligned} \quad (7.36)$$

$$\begin{aligned} \gamma_{\xi 2}^{(\beta)}(X) &= \frac{1}{3\pi^2} \int_0^\infty d\mathcal{E} \left[\theta(\mathcal{E} - \mathcal{M}_\xi(X)) (\mathcal{E}^2 - \mathcal{M}_\xi^2(X))^{3/2} - \mathcal{E}^3 + \frac{3}{2} \mathcal{E} \mathcal{M}_\xi^2(X) \right] \\ &\quad \times \langle N^{(\xi)}(X; \mathcal{E}) \rangle. \end{aligned} \quad (7.37)$$

In the case of locally equilibrium system, $b_\xi(X) = c_\xi(X) = 1$. At “high temperature,” $\gamma_{\xi 1}^{(\beta)}$ dominates, $|\gamma_{\xi 2}^{(\beta)}| \ll \gamma_{\xi 1}^{(\beta)}$. $\gamma_{\xi 2}^{(\beta)}$ may be estimated by substituting (7.34) into (7.37):

$$\gamma_{\xi 2}^{(\beta)}(X) \simeq \frac{1}{6\pi} \mathcal{M}_\xi^3(X) \mathcal{T}_\xi(X).$$

Putting all this together, we obtain

$$\gamma_\xi^{(\beta)}(X) \simeq \frac{\pi^2}{45} b_\xi(X) \mathcal{T}_\xi^4(X) - \frac{1}{12} c_\xi(X) \mathcal{M}_\xi^2(X) \mathcal{T}_\xi^2(X) + \frac{1}{6\pi} \mathcal{M}_\xi^3(X) \mathcal{T}_\xi(X). \quad (7.38)$$

The local effective action $\gamma_{phys}(X)$, Eq. (7.20), now becomes

$$\gamma_{phys}(X) \simeq \gamma_{phys}^{(0)}(X) + \gamma_{phys}^{(\beta)}(X),$$

where $\gamma_{phys}^{(0)}(X)$ is as in (7.21), and

$$\begin{aligned} \gamma_{phys}^{(\beta)}(X) &\simeq \frac{1}{90} \left[(N-1) \left\{ \pi^2 b_\pi(X) \mathcal{T}_\pi^4(X) - \frac{15}{4} c_\pi(X) \mathcal{M}_\pi^2(X) \mathcal{T}_\pi^2(X) \right\} \right. \\ &\quad \left. + \left\{ \pi^2 b_\sigma(X) \mathcal{T}_\sigma^4(X) - \frac{15}{4} c_\sigma(X) \mathcal{M}_\sigma^2(X) \mathcal{T}_\sigma^2(X) \right\} \right] \\ &\quad + \frac{1}{12\pi} \left[(N-1) \mathcal{M}_\pi^3(X) \mathcal{T}^{(\pi)}(X) + \mathcal{M}_\sigma^3(X) \mathcal{T}^{(\sigma)}(X) \right]. \end{aligned}$$

In the case where $\mathcal{M}_\pi^2(X) \ll \mathcal{T}_\pi(X)$ and $\mathcal{M}_\sigma^2(X) \gg \mathcal{T}_\sigma(X)$, the “ σ -terms” here and below are dropped.

High-temperature expansion of $\Omega_\beta(\vec{\varphi}^2(X))$, Eq. (7.28), and $\Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X))$, Eq. (7.33), reads, in respective order,

$$\begin{aligned} \Omega_\beta(\vec{\varphi}^2(X)) &\simeq \frac{\lambda}{72} \left[(N-1) \left\{ c_\pi \mathcal{T}_\pi^2(X) - \frac{3}{\pi} \mathcal{M}_\pi(X) \mathcal{T}_\pi(X) \right\} \right. \\ &\quad \left. + 3 \left\{ c_\sigma \mathcal{T}_\sigma^2(X) - \frac{3}{\pi} \mathcal{M}_\sigma(X) \mathcal{T}_\sigma(X) \right\} \right] \end{aligned}$$

and

$$\Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X)) \simeq -\frac{\lambda^2 \varphi_0^2(X)}{144\pi} \left[(N-1) \frac{\mathcal{T}_\pi(X)}{\mathcal{M}_\pi(X)} + 9 \frac{\mathcal{T}_\sigma(X)}{\mathcal{M}_\sigma(X)} \right].$$

2. “Low-temperature” expansion

Here we consider locally thermal equilibrium system, which is characterized by (7.35). In the low-temperature limit, $T(X) \ll \mathcal{M}_\xi^2(X)$, $\gamma_\xi^{(\beta)}(X)$ in (7.17) may be computed as

$$\begin{aligned} \gamma_\xi^{(\beta)}(X) &\simeq \frac{1}{3\pi^2} \int_{\mathcal{M}_\xi(X)}^\infty dE (E^2 - \mathcal{M}_\xi^2(X))^{3/2} \frac{1}{e^{E/T(X)} - 1} \\ &\simeq \frac{1}{3\pi^2} (\mathcal{M}_\xi^2(X))^{3/4} e^{-\mathcal{M}_\xi^2(X)/T(X)} \int_0^\infty dx x^{3/2} e^{-x/T(X)} \\ &= \frac{1}{\sqrt{2}\pi} (\mathcal{M}_\xi^2(X))^{3/4} (T(X))^{5/2} e^{-\mathcal{M}_\xi(X)/T(X)}. \end{aligned} \quad (7.39)$$

Then, $\gamma_{phys}^{(\beta)}(X)$ in (7.22) is approximated as

$$\begin{aligned} \gamma_{phys}^{(\beta)}(X) &\simeq \frac{1}{2^{3/2}\pi} (T(X))^{5/2} \left[(N-1) (\mathcal{M}_\pi^2(X))^{3/4} e^{-\mathcal{M}_\pi(X)/T(X)} \right. \\ &\quad \left. + (\mathcal{M}_\sigma^2(X))^{3/4} e^{-\mathcal{M}_\sigma(X)/T(X)} \right]. \end{aligned}$$

$\Omega_\beta(\vec{\varphi}_0^2(X))$ in (7.28) becomes

$$\begin{aligned} \Omega_\beta(\vec{\varphi}_0^2(X)) &\simeq \frac{\lambda}{12\sqrt{2}\pi} (T(X))^{3/2} \left[(N-1) (\mathcal{M}_\pi^2(X))^{1/4} e^{-\mathcal{M}_\pi(X)/T(X)} \right. \\ &\quad \left. + 3 (\mathcal{M}_\sigma^2(X))^{1/4} e^{-\mathcal{M}_\sigma(X)/T(X)} \right] \end{aligned}$$

and $\Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X))$ in (7.33) becomes

$$\begin{aligned} \Omega_\beta^{(\sigma)}(\vec{\varphi}_0^2(X)) &\simeq -\frac{\lambda^2 \varphi_0^2(X)}{72\sqrt{2}\pi} (T(X))^{1/2} \left[(N-1) (\mathcal{M}_\pi^2(X))^{-1/4} e^{-\mathcal{M}_\pi(X)/T(X)} \right. \\ &\quad \left. + 9 (\mathcal{M}_\sigma^2(X))^{-1/4} e^{-\mathcal{M}_\sigma(X)/T(X)} \right]. \end{aligned}$$

B. The region $\mathcal{M}_\pi^2(X)$ and/or $\mathcal{M}_\sigma^2(X) < 0$

It can be seen that, through analytically continuing the one-loop effective action obtained above to the region where $\mathcal{M}_\pi^2(X)$ and/or $\mathcal{M}_\sigma^2(X) < 0$, an imaginary part emerges and then we face the issue of interpreting it. In the case of effective potential in vacuum theory, this issue has been studied, e.g., in [37]. Let us summarize the observation in [37]. The quantity analyzed there is essentially (7.15). In (7.15), the quantum modes with $p^2 \geq |\mathcal{M}_\xi^2(X)|$ remain harmonic oscillator as in the case of $\mathcal{M}_\xi^2(X) \geq 0$. However, the long-wavelength modes, $p^2 < |\mathcal{M}_\xi^2(X)|$, have imaginary

frequencies and thus unstable. These unstable modes turn out to develop imaginary part of $\gamma_\xi^{(0)}(X)$, $Im \gamma_\xi^{(0)}(X)$, the emergence of which is a signal of appearance of perturbative instability. The $Im \gamma_\xi^{(0)}(X)$ may be interpreted as half the decay rate per unit volume of a particular quantum state. More precisely, $Im \gamma_\xi^{(0)}(X)$ is the sum of half the decay rate per unit volume over all unstable modes, $\phi_{long}(x)$, which are localized at $\langle \phi_{long}(x_0 = (x_0)_{in}, \mathbf{x}) \rangle \simeq 0$ at the initial time $x_0 = (x_0)_{in}$. Here $\phi_{long}(x)$ stands for the long-wavelength components of the quantum field $\phi(x)$. The perturbative instability arises due to the growth of $\langle \phi_{long}(x) \rangle$. Owing to this instability, the original quasiuniform configuration beaks up into many uncorrelated domains. Within a domain the quantum field is correlated and $\langle \phi_{long}(x) \rangle$ is increasing in time. From one domain to the next, $\langle \phi_{long}(x) \rangle$'s vary randomly from positive to negative. These domains is expected to have size at least as great as $|\mathcal{M}_\xi^2(X)|^{-1/2}$, the minimum wavelength for an unstable mode.

As seen above, leading one-loop contribution to the effective action is given by an integral of the “local effective action” $\gamma_{phys}(X)$ over the whole macroscopic spacetime region. We recall that $\gamma_{phys}(X)$ is a function of X that labels the spacetime cells. Then, in a single cell, X of $\gamma_{phys}(X)$ is approximately constant. Moreover $\gamma_{phys}(X)$ is the same in form as the effective potential (density). From this observation, together with the conclusion [37] for the zero-temperature effective potential, we assume that the imaginary part of the effective action in the present case also reflects the proper physical situation.

For the purpose of performing analytic continuation, we return back to (7.14) - (7.17) with $\mathcal{M}_\xi^2(X) \geq 0$. Equations (5.34) and (7.15) tell us that $\mathcal{M}_\xi^2(X)$ in (7.17) should be understood to be $\mathcal{M}_\xi^2(X) - i\eta$ with η a positive infinitesimal constant. Now let us analytically continue the function $\gamma_\xi = \gamma_\xi^{(0)} + \gamma_\xi^{(\beta)}$ from the region $\mathcal{M}_\xi^2(X) \geq 0$ to the region $\mathcal{M}_\xi^2(X) < 0$. The UV-renormalized vacuum part $\gamma_\xi^{(0)}$, Eq. (7.16), goes to

$$\gamma_\xi^{(0)} = -\frac{1}{32\pi^2} \mathcal{M}_\xi^4(X) \ln \frac{-\mathcal{M}_\xi^2(X)}{e^{3/2} \mu_d^2} + \frac{i}{32\pi} \mathcal{M}_\xi^4(X) \quad (\mathcal{M}_\xi^2(X) < 0). \quad (7.40)$$

It is worth mentioning that a “direct calculation” of (7.10) yields the correct result for $Im \gamma_\xi^{(0)}$:

$$\begin{aligned} i Im \mathcal{I}_\xi^{(0)}(X) &= i Im \int \frac{d^4 P}{(2\pi)^4} \theta(P^2 - \mathcal{M}_\xi^2(X)) \frac{i}{P^2 - \mathcal{M}_\xi^2(X) + i\eta} \\ &= i \int \frac{dp}{4\pi^2} \frac{p^2}{\sqrt{-p^2 - \mathcal{M}_\xi^2(X)}} \\ &= -\frac{i}{16\pi} \mathcal{M}_\xi^2(X), \end{aligned}$$

from which we obtain

$$\begin{aligned} i Im \gamma_\xi^{(0)} &\equiv - \int d\mathcal{M}_\xi^2(X) i Im \mathcal{I}_\xi^{(0)}(X) \\ &= \frac{i}{32\pi} \mathcal{M}_\xi^4(X), \end{aligned} \quad (7.41)$$

in accord with (7.40).

As to $\gamma_\xi^{(\beta)}$, we rewrite (7.17) as

$$\gamma_\xi^{(\beta)} = \frac{i}{6\pi^2} \int_{C_\mathcal{E}} d\mathcal{E} (\mathcal{E}^2 - \mathcal{M}_\xi^2(X) + i\eta) (\mathcal{M}_\xi^2(X) - \mathcal{E}^2 - i\eta)^{1/2} \langle N^{(\xi)}(X; \mathcal{E}) \rangle \quad (\mathcal{M}_\xi^2(X) > 0), \quad (7.42)$$

where $\langle N^{(\xi)}(X; \mathcal{E}) \rangle$ is as in (7.13). In a complex- $\mathcal{E}$ plane, the function $(\mathcal{M}_\xi^2(X) - \mathcal{E}^2 - i\eta)^{1/2}$ has branch points at $\mathcal{E} = \pm \mathcal{E}_{br}$ with $\mathcal{E}_{br} = (\mathcal{M}_\xi^2(X) - i\eta)^{1/2}$. We take a branch cut associated with $+\mathcal{E}_{br}$ to be a straight line $[\mathcal{E}_{br}, +\infty - i\alpha)$, where $\alpha = -Im(\mathcal{M}_\xi^2(X) - i\eta)^{1/2}$. The integration contour $C_\mathcal{E}$ in (7.42) is defined so as to enclose the branch cut $[\mathcal{E}_{br}, +\infty - i\alpha)$ in a clock-wise direction,

$$\begin{aligned} C_\mathcal{E} : +\infty - i(\alpha + \epsilon) &\rightarrow \mathcal{M}_\xi(X) - \epsilon - i(\alpha + \epsilon) \\ &\rightarrow \mathcal{M}_\xi(X) - \epsilon + i(\epsilon - \alpha) \rightarrow +\infty + i(\epsilon - \alpha), \end{aligned}$$

where $0 < \epsilon < \alpha$.

Continuing to the negative $\mathcal{M}_\xi^2(X)$ region, the branch points $\pm E_{br}$ go to

$$\pm E_{br} \rightarrow \pm \left(-i(-\mathcal{M}_\xi^2(X))^{1/2} + \alpha \right) \quad (\alpha = \text{Im}(-\mathcal{M}_\xi^2(X) + i\eta)^{1/2}) \quad (7.43)$$

and the branch cut associated with $\mathcal{E}_{br}$ goes to $[\mathcal{E}_{br}, (1-i)\alpha] \oplus [(1-i)\alpha, +\infty - i\alpha]$. The integration contour $C_\mathcal{E}$ is deformed accordingly. We assume that $\langle N^{(\xi)}(X; \mathcal{E}) \rangle$ is analytic in the region

$$\begin{aligned} \alpha - \epsilon < \text{Re}\mathcal{E} < (-\mathcal{M}_\xi^2(X))^{1/2} + \alpha + \epsilon, \\ -(-\mathcal{M}_\xi^2(X))^{1/2} - \epsilon < \text{Im}\mathcal{E} < \epsilon - \alpha. \end{aligned}$$

Then, we can deform the branch cut associated with $\mathcal{E}_{br}$ to

$$\begin{aligned} & [\mathcal{E}_{br}, \mathcal{E}_{br} + (-\mathcal{M}_\xi^2(X))^{1/2}] \oplus [\mathcal{E}_{br} + (-\mathcal{M}_\xi^2(X))^{1/2}, (-\mathcal{M}_\xi^2(X))^{1/2} - i\alpha] \\ & \oplus [(-\mathcal{M}_\xi^2(X))^{1/2} - i\alpha, +\infty - i\alpha]. \end{aligned}$$

The integration contour $C_\mathcal{E}$ is deformed accordingly.

Then, we obtain, after some manipulations,

$$\gamma_\xi^{(\beta)} = \frac{1}{3\pi^2} \int_{C'} dz (z^2 - \mathcal{M}_\xi^2(X))^{3/2} \langle N^{(\xi)}(X; z) \rangle, \quad (7.44)$$

where $C' = C'_1 \oplus C'_2 \oplus C'_3$ with $C'_1 = -i(-\mathcal{M}_\xi^2(X))^{1/2} \rightarrow (1-i)(-\mathcal{M}_\xi^2(X))^{1/2}$, $C'_2 = (1-i)(-\mathcal{M}_\xi^2(X))^{1/2} \rightarrow (-\mathcal{M}_\xi^2(X))^{1/2}$, and $C'_3 = (-\mathcal{M}_\xi^2(X))^{1/2} \rightarrow +\infty$.

After all this, we obtain, for the local effective action,

$$\gamma_{phys}(X) = \gamma_{phys}^{(0)}(X) + \gamma_{phys}^{(\beta)}(X), \quad (7.45)$$

where

$$\begin{aligned} \text{Re } \gamma_{phys}^{(0)}(X) &= \frac{1}{2} [(\partial\vec{\varphi}(X))^2 - \mu^2\vec{\varphi}^2(X)] - \frac{\lambda}{4!} (\vec{\varphi}^2(X))^2 + h(t\vec{e} \cdot \vec{\varphi}(X)) \\ &\quad - \frac{1}{(8\pi)^2} \left[(N-1)\mathcal{M}_\pi^4(X) \ln \frac{|\mathcal{M}_\pi^2(X)|}{e^{2/3}\mu_d^2} + \mathcal{M}_\sigma^4(X) \ln \frac{|\mathcal{M}_\sigma^2(X)|}{e^{2/3}\mu_d^2} \right] \end{aligned} \quad (7.46)$$

$$\text{Im } \gamma_{phys}^{(0)}(X) = \frac{1}{64\pi} [(N-1)\theta(-\mathcal{M}_\pi^2(X))\mathcal{M}_\pi^4(X) + \theta(-\mathcal{M}_\sigma^2(X))\mathcal{M}_\sigma^4(X)], \quad (7.47)$$

$$\gamma_{phys}^{(\beta)}(X) = \frac{1}{2} [(N-1)\gamma_\pi^{(\beta)}(X) + \gamma_\sigma^{(\beta)}(X)]. \quad (7.48)$$

As mentioned at the beginning of this subsection, $\text{Im } \Gamma_{phys}(X)$ causes the perturbative instability, which in turn leads to formation of a domain structure. In most cases analyzed so far in the literature, imaginary part appears in the CTP effective action Γ in the form,

$$\text{Im}\Gamma = i \sum_{i,j=1}^N \int d^4x d^4y (\varphi_1(x) - \varphi_2(x))_i \gamma_{ij}(x, y) (\varphi_1(y) - \varphi_2(y))_j, \quad (7.49)$$

where γ_{ij} are real functions. Recalling that the limit $\vec{\varphi}_2(x) \rightarrow \vec{\varphi}_1(x)$ is taken at the final stage, we see that (7.49) describes small fluctuations. In the present case, (7.5) shows that $\text{Im}\Gamma$ under consideration is not of the form (7.49) and describes “large” fluctuations.⁴ Here, for dealing with $\text{Im}\Gamma_{phys}$, we follow a standard procedure [38], which starts with writing $\exp i\Gamma_{phys}$ in the form,

⁴Even in the present case, the small fluctuations of the type (7.49) also appears, which we do not consider in this paper.

$$\begin{aligned}
e^{i\Gamma_{phys}} &= \exp i \int d^4 X \gamma_{phys}(X) \\
&= \exp \left[i \int d^4 X \operatorname{Re} \gamma_{phys}(X) - \int d^4 X \operatorname{Im} \gamma_{phys}(X) \right].
\end{aligned} \tag{7.50}$$

Provided that

$$\int [d\vec{\varphi}] \exp \left[- \int d^4 X \operatorname{Im} \gamma_{phys}(X) \right] < \infty, \tag{7.51}$$

we can make a functional Fourier transformation of $\exp[-\int d^4 X \operatorname{Im} \gamma_{phys}(X)]$:

$$\begin{aligned}
\exp \left[- \int d^4 X \operatorname{Im} \gamma_{phys}(X) \right] &= \exp \left[- \int d^4 X \operatorname{Im} \gamma_{phys}(X) \right]_{\vec{\varphi}=0} \\
&\times \int [d\vec{\zeta}] \exp \left[- \int d^4 X G(\vec{\zeta}^2(X)) + i \int d^4 X {}^t\vec{\zeta}(X) \cdot \vec{\varphi}(X) \right].
\end{aligned} \tag{7.52}$$

Substituting (7.52) into (7.50), we obtain

$$\begin{aligned}
e^{i\Gamma_{phys}} &= \exp \left[- \int d^4 X \operatorname{Im} \gamma_{phys}(X) \right]_{\vec{\varphi}=0} \\
&\times \langle \exp \left[i \int d^4 X \left\{ \operatorname{Re} \gamma_{phys}(X) + {}^t\vec{\zeta}(X) \cdot \vec{\varphi}(X) \right\} \right] \rangle_{\zeta},
\end{aligned} \tag{7.53}$$

where the $\vec{\zeta}$ -average $\langle \dots \rangle_{\zeta}$ is defined as

$$\langle \dots \rangle_{\zeta} \equiv \int [d\vec{\zeta}] e^{-\int d^4 X G(\vec{\zeta}^2(X))} \dots$$

Equation (7.53) tells us that $\operatorname{Re} \gamma_{phys}(X) + {}^t\vec{\zeta}(X) \cdot \vec{\varphi}(X)$ may be interpreted as the local effective action, from which follows the equation of motion for $\vec{\varphi}(X)$,

$$\frac{\partial \operatorname{Re} \gamma_{phys}(X)}{\partial \vec{\varphi}(X)} + \vec{\zeta}(X) = 0. \tag{7.54}$$

Thus, the auxiliary fields $\vec{\zeta}(X)$ represent random forces, whose character is determined by $G(\vec{\zeta}^2(X))$.

Taking the functional derivatives of (7.52) with respect to $\vec{\varphi}$ and then setting $\vec{\varphi} = 0$, we obtain the relations:

$$\begin{aligned}
\langle 1 \rangle_{\zeta} &= 1, \\
\langle \vec{\zeta}(X) \rangle_{\zeta} &= 0, \\
\langle \zeta_i(X) \zeta_j(Y) \rangle_{\zeta} &= 2\delta_{ij} \delta^4(X - Y) \frac{\partial \operatorname{Im} \gamma_{phys}(X)}{\partial \vec{\varphi}^2(X)} \Big|_{\vec{\varphi}=0} \\
&= \delta_{ij} \delta^4(X - Y) \left[\frac{\lambda}{96\pi} \{ (N-1)\theta(-M_{\pi}^2(X)) M_{\pi}^2(X) \right. \\
&\quad \left. + 3\theta(-M_{\sigma}^2(X)) M_{\sigma}^2(X) \} + (N-1)\theta(-M_{\pi}^2(X)) \frac{\partial \operatorname{Im} \gamma_{\pi}^{(\beta)}(X)}{\partial \vec{\varphi}^2(X)} \Big|_{\vec{\varphi}=0} \right. \\
&\quad \left. + \theta(-M_{\sigma}^2(X)) \frac{\partial \operatorname{Im} \gamma_{\sigma}^{(\beta)}(X)}{\partial \vec{\varphi}^2(X)} \Big|_{\vec{\varphi}=0} \right],
\end{aligned} \tag{7.55}$$

$$\begin{aligned}
&\langle \zeta_i(X) \zeta_j(Y) \zeta_k(U) \zeta_l(V) \rangle_{\zeta c} \\
&\equiv \langle \zeta_i(X) \zeta_j(Y) \zeta_k(U) \zeta_l(V) \rangle_{\zeta} - [\langle \zeta_i(X) \zeta_j(Y) \rangle_{\zeta} \langle \zeta_k(U) \zeta_l(V) \rangle_{\zeta}
\end{aligned}$$

$$\begin{aligned}
& + \langle \zeta_i(X) \zeta_k(U) \rangle_\zeta \langle \zeta_j(Y) \zeta_l(V) \rangle_\zeta + \langle \zeta_i(X) \zeta_l(V) \rangle_\zeta \langle \zeta_j(Y) \zeta_k(U) \rangle_\zeta] \\
& = -2 \delta^4(X-U) \delta^4(V-Y) \delta^4(X-Y) [\delta_{ik} \delta_{lj} + \delta_{kj} \delta_{li} + \delta_{ij} \delta_{lk}] \\
& \times \left\{ \frac{\lambda^2}{576\pi} [(N-1) \theta(-M_\pi^2(X)) + 9 \theta(-M_\sigma^2(X))] \right. \\
& + (N-1) \theta(-M_\pi^2(X)) \left. \frac{\partial^2 \text{Im} \gamma_\pi^{(\beta)}(X)}{\partial(\varphi^2(X))^2} \right|_{\vec{\varphi}=0} \\
& + \theta(-M_\sigma^2(X)) \left. \frac{\partial^2 \text{Im} \gamma_\sigma^{(\beta)}(X)}{\partial(\varphi^2(X))^2} \right|_{\vec{\varphi}=0} \Big\}, \tag{7.56}
\end{aligned}$$

etc. As mentioned in Section I, the macroscopic spacetime coordinates are the indices that labels the spacetime cells, whose size is L^μ ($\mu = 0, 1, 2, 3$). Then, $\delta(X^\mu - Y^\mu)$ in (7.55) and (7.56) should be understood to be

$$\delta(X^\mu - Y^\mu) \simeq \frac{\delta_{i^\mu j^\mu}}{L^\mu} \quad (\text{no summation over } \mu),$$

where i^μ , etc., are the μ -component of the label for the spacetime cells.

Let $\vec{\varphi}(X) = \vec{\varphi}(X; [\vec{\zeta}])$ be a solution to (7.54) under a given initial data (at the initial time $X_0 = T_{in}$), $\vec{\varphi}(X_0 = T_{in}, \mathbf{X})$ and $\partial \vec{\varphi}(X_0, \mathbf{X}) / \partial X_0 \Big|_{X_0 = T_{in}}$. Then using (7.55), (7.56), etc., one can compute the ζ -averaged correlation functions

$$\begin{aligned}
& \langle \varphi_i(X) \rangle_\zeta, \quad \langle \varphi_i(X) \varphi_j(Y) \rangle_\zeta, \\
& \langle \varphi_i(X) \varphi_j(Y) \varphi_k(Z) \rangle_\zeta, \quad \langle \varphi_i(X) \varphi_j(Y) \varphi_k(U) \varphi_l(V) \rangle_\zeta,
\end{aligned}$$

etc.

A multi-point correlation function consists of “connected” and “disconnected” parts. For example, a two-point correlation function $\langle \varphi_i(X) \varphi_j(Y) \rangle_\zeta$ takes the form

$$\langle \varphi_i(X) \varphi_j(Y) \rangle_\zeta = \langle \varphi_i(X) \varphi_j(Y) \rangle_{\zeta c} + \langle \varphi_i(X) \rangle_\zeta \langle \varphi_j(Y) \rangle_\zeta,$$

where the first term on the RHS stands for “connected” part and the second term stands for “disconnected” part. It is convenient to introduce a generating functional

$$\mathcal{G}[\vec{y}, \vec{\varphi}] \equiv \langle e^{i \int d^4 Z \vec{y}(Z) \cdot \vec{\varphi}(Z)} \rangle_\zeta, \tag{7.57}$$

which generates $\langle \varphi_{i_1}(X_1) \dots \varphi_{i_n}(X_n) \rangle_\zeta$ through

$$\langle \varphi_{i_1}(X_1) \dots \varphi_{i_n}(X_n) \rangle_\zeta = (-i)^n \prod_{j=1}^n \frac{\delta}{\delta y_{i_j}(X_j)} \mathcal{G}[\vec{y}, \vec{\varphi}] \Big|_{\vec{y}=0}.$$

$\vec{\zeta}$ -averaged “connected” correlation functions are obtained through

$$\langle \varphi_{i_1}(X_1) \dots \varphi_{i_n}(X_n) \rangle_{\zeta c} = (-i)^n \prod_{j=1}^n \frac{\delta}{\delta y_{i_j}(X_j)} \ln \mathcal{G}[\vec{y}, \vec{\varphi}] \Big|_{\vec{y}=0}. \tag{7.58}$$

It is worth mentioning here the equation of motion for type-2 or unphysical condensate fields. From (7.5), it is obvious that, for the unphysical condensate fields, all the above formulae apply with the substitutions,

$$\text{Re} \gamma_{phys} \rightarrow \text{Re} \gamma_{phys}, \quad \text{Im} \gamma_{phys} \rightarrow -\text{Im} \gamma_{phys}.$$

This means that (7.55) and (7.56), etc., change sign. Recalling again that the type-2 field “lives” in the “time-reversed world,” we see that the random forces that statistically upsetting the system act in completely opposite way in the time-reversed world.

1. “High-temperature” expansion

In a similar manner as in the case of $\mathcal{M}_\pi^2(X)$, $\mathcal{M}_\sigma^2(X) \geq 0$, one can obtain a “high-temperature” expansion of $\gamma_{phys}(X)$. Using (7.34) in (7.44), we obtain (cf. (7.36) - (7.38))

$$\begin{aligned} \gamma_\xi^{(\beta)}(X) &\simeq \frac{1}{3\pi^2} \int_0^\infty d\mathcal{E} \left[\mathcal{E}^3 - \frac{3}{2} \mathcal{E} \mathcal{M}_\xi^2(X) \right] \langle N^{(\xi)}(X; \mathcal{E}) \rangle \\ &\quad + \frac{1}{3\pi^2} \left[\int_{C'} dz (z^2 - \mathcal{M}_\xi^2(X))^{3/2} - \int_0^\infty dz \left(z^3 - \frac{3}{2} z \mathcal{M}_\xi^2(X) \right) \right] \frac{\mathcal{T}_\xi(X)}{z} \\ &\simeq \frac{\pi^2}{45} b_\xi(X) \mathcal{T}_\xi^4(X) - \frac{1}{12} c_\xi(X) \mathcal{M}_\xi^2(X) \mathcal{T}_\xi^2(X) + \frac{i}{6\pi} (-\mathcal{M}_\xi^2(X))^{3/2} \mathcal{T}_\xi(X). \end{aligned} \quad (7.59)$$

It is to be noted that (7.59) is directly obtained by analytically continuing the asymptotic form (7.38) from the region $\mathcal{M}_\xi^2(X) (> 0)$ to the region $\mathcal{M}_\xi^2(X) - i\eta$ ($\mathcal{M}_\xi^2(X) < 0$). It is also worth mentioning that the following “direct calculation” reproduces (7.59). We start from (7.37),

$$\begin{aligned} \gamma_{\xi 2}^{(\beta)}(X) &= \frac{1}{3\pi^2} \left[\int_0^\infty dp \frac{p^4}{\sqrt{p^2 + \mathcal{M}_\xi^2(X)}} \langle N^{(\xi)}(X; \sqrt{p^2 + \mathcal{M}_\xi^2(X)}) \rangle \right. \\ &\quad \left. - \int_0^\infty d\mathcal{E} \left(\mathcal{E}^3 - \frac{3}{2} \mathcal{E} \mathcal{M}_\xi^2(X) \right) \langle N^{(\xi)}(X; \mathcal{E}) \rangle \right], \end{aligned}$$

which is valid for $\mathcal{M}_\xi^2 \geq 0$. At this stage, let us naively change $\mathcal{M}_\xi^2(X)$ from positive to negative. Then we have

$$\begin{aligned} \gamma_{\xi 2}^{(\beta)}(X) &= \frac{1}{3\pi^2} \left[\int_0^\infty dp \frac{p^4}{\sqrt{p^2 - |\mathcal{M}_\xi^2(X)|}} \langle N^{(\xi)}(X; \sqrt{p^2 - |\mathcal{M}_\xi^2(X)|}) \rangle \right. \\ &\quad \left. - \int_0^\infty d\mathcal{E} \left(\mathcal{E}^3 - \frac{3}{2} \mathcal{E} \mathcal{M}_\xi^2(X) \right) \langle N^{(\xi)}(X; \mathcal{E}) \rangle \right]. \end{aligned}$$

Substituting (7.34) and using (7.36), we obtain

$$\begin{aligned} \gamma_{\xi 2}^{(\beta)}(X) &\simeq \frac{\mathcal{T}_\xi(X)}{3\pi^2} \left[\int_0^\infty dp \frac{p^4}{p^2 - |\mathcal{M}_\xi^2(X)| - i\eta} \right. \\ &\quad \left. - \int_{\sqrt{|\mathcal{M}_\xi^2(X)|}}^\infty dp p \left(\sqrt{p^2 - |\mathcal{M}_\xi^2(X)|} + \frac{3}{2} \frac{|\mathcal{M}_\xi^2(X)|}{\sqrt{p^2 - |\mathcal{M}_\xi^2(X)|}} \right) \right] \\ &= \frac{i\mathcal{T}_\xi(X)}{3\pi} \int_0^\infty dp p^4 \delta(p^2 - |\mathcal{M}_\xi^2(X)|) \\ &= \frac{i}{6\pi} |\mathcal{M}_\xi^2(X)|^{3/2} \mathcal{T}_\xi(X). \end{aligned}$$

This together with (7.36) reproduces (7.59).

Thus, $\gamma_{phys}^{(\beta)}(X)$ in (7.48) is approximated as

$$\begin{aligned} \gamma_{phys}^{(\beta)}(X) &\simeq \frac{1}{90} \left[(N-1) \left(\pi^2 b_\pi(X) \mathcal{T}_\pi^4(X) - \frac{15}{4} c_\pi(X) \mathcal{M}_\pi^2(X) \mathcal{T}_\pi^2(X) \right) \right. \\ &\quad \left. + \left(\pi^2 b_\sigma(X) \mathcal{T}_\sigma^4(X) - \frac{15}{4} c_\sigma(X) \mathcal{M}_\sigma^2(X) \mathcal{T}_\sigma^2(X) \right) \right] \\ &\quad + \frac{1}{12\pi} \left[(N-1) \theta(\mathcal{M}_\pi^2(X)) \mathcal{M}_\pi^3(X) \mathcal{T}^{(\pi)}(X) + \theta(\mathcal{M}_\sigma^2(X)) \mathcal{M}_\sigma^3(X) \mathcal{T}^{(\sigma)}(X) \right] \\ &\quad + \frac{i}{12\pi} \left[(N-1) \theta(-\mathcal{M}_\pi^2(X)) (-\mathcal{M}_\pi^2(X))^{3/2} \mathcal{T}^{(\pi)}(X) \right. \\ &\quad \left. + \theta(-\mathcal{M}_\sigma^2(X)) (-\mathcal{M}_\sigma^2(X))^{3/2} \mathcal{T}^{(\sigma)}(X) \right]. \end{aligned} \quad (7.60)$$

The moment (7.55) and (7.56) are approximated, in respective order, as

$$\begin{aligned}
\langle \zeta_i(X) \zeta_j(Y) \rangle_\zeta &= 2\delta_{ij} \delta(X-Y) \left. \frac{\partial \text{Im} \gamma_{phys}(X)}{\partial \vec{\varphi}^2(X)} \right|_{\vec{\varphi}=0} \\
&\simeq -\frac{\lambda}{24\pi} \delta_{ij} \delta(X-Y) \left[(N-1)\theta(-M_\pi^2(X)) (-M_\pi^2(X))^{1/2} \mathcal{T}_\pi(X) \right. \\
&\quad \left. + 3\theta(-M_\sigma^2(X)) (-M_\sigma^2(X))^{1/2} \mathcal{T}_\sigma(X) \right. \\
&\quad \left. - \frac{1}{4}(N-1)\theta(-M_\pi^2(X)) M_\pi^2(X) + \frac{3}{4}\theta(-M_\sigma^2(X)) M_\sigma^2(X) \right]
\end{aligned}$$

and

$$\begin{aligned}
&\langle \zeta_i(X) \zeta_j(Y) \zeta_k(U) \zeta_l(V) \rangle_{\zeta_c} \\
&\simeq -\frac{\lambda^2}{288\pi} \delta^4(X-U) \delta^4(V-Y) \delta^4(X-Y) [\delta_{ik} \delta_{lj} + \delta_{kj} \delta_{li} + \delta_{ij} \delta_{lk}] \\
&\quad \times \left\{ (N-1)\theta(-M_\pi^2(X)) + 9\theta(-M_\sigma^2(X)) \right. \\
&\quad \left. + 2 \left[(N-1)\theta(-M_\pi^2(X)) (-M_\pi^2(X))^{-1/2} \mathcal{T}_\pi(X) \right. \right. \\
&\quad \left. \left. + 9\theta(-M_\sigma^2(X)) (-M_\sigma^2(X))^{-1/2} \mathcal{T}_\sigma(X) \right] \right\}.
\end{aligned}$$

$\text{Im} \gamma_{phys}(X) = \text{Im} \gamma_{phys}^{(0)}(X) + \text{Im} \gamma_{phys}^{(\beta)}(X)$ vanishes for $\mathcal{M}_\pi^2(X) > 0$ and $\mathcal{M}_\sigma^2(X) > 0$, i.e., for

$$\vec{\varphi}^2(X) > \text{Max} \left(-\frac{6}{\lambda} M_\pi^2(X), -\frac{2}{\lambda} M_\sigma^2(X) \right).$$

Then, the condition (7.51) is met.

2. Low-temperature limit

We consider locally thermal equilibrium system, which is characterized by (7.35), and obtain $\gamma_{phys}^{(\beta)}(X)$ in the low-temperature limit, $T(X) \ll -\mathcal{M}_\xi^2(X)$. The function (7.35) satisfy the assumption made above after (7.43), we can use the representation (7.44). From (7.44), one can easily see that the contribution from the segment C'_1 dominates over others. Changing the integration variable from z to ρ through $z = (-\mathcal{M}_\xi^2(X))[\rho - i]$, we have

$$\begin{aligned}
\gamma_\xi^{(\beta)} &= \frac{1}{3\pi^2} (-\mathcal{M}_\xi^2(X))^2 \int_0^\infty d\rho \frac{\rho^{3/2}(\rho - 2i)^{3/2}}{\exp\left(\frac{(-\mathcal{M}_\xi^2(X))^{1/2}}{T(X)}(\rho - i)\right) - 1} \\
&\simeq \frac{2\sqrt{2}}{3\pi^2} e^{-3\pi i/4} e^{i(-\mathcal{M}_\xi^2(X))^{1/2}/T(X)} (-\mathcal{M}_\xi^2(X))^2 \int_0^\infty d\rho \rho^{3/2} \exp\left(-\frac{(-\mathcal{M}_\xi^2(X))^{1/2}}{T(X)} \rho\right) \\
&\simeq \frac{1}{\sqrt{2}\pi} e^{-3\pi i/4} e^{i(-\mathcal{M}_\xi^2(X))^{1/2}/T(X)} (-\mathcal{M}_\xi^2(X))^{3/4} (T(X))^{5/2},
\end{aligned}$$

from which, we obtain

$$\begin{aligned}
\text{Re} \gamma_\xi^{(\beta)} &= \frac{1}{\sqrt{2}\pi} (-\mathcal{M}_\xi^2(X))^{3/4} (T(X))^{5/2} \cos\left(\frac{(-\mathcal{M}_\xi^2(X))^{1/2}}{T(X)} - \frac{3\pi}{4}\right), \\
\text{Im} \gamma_\xi^{(\beta)} &= \frac{1}{\sqrt{2}\pi} (-\mathcal{M}_\xi^2(X))^{3/4} (T(X))^{5/2} \sin\left(\frac{(-\mathcal{M}_\xi^2(X))^{1/2}}{T(X)} - \frac{3\pi}{4}\right).
\end{aligned} \tag{7.61}$$

Incidentally, (7.61) is directly obtained from (7.39) through an analytic continuation, $\mathcal{M}_\xi^2(X) (\geq 0) \rightarrow \mathcal{M}_\xi^2(X) - i\eta$ ($\mathcal{M}_\xi^2 < 0$).

The form of $\gamma_{phys}^{(\beta)}(X)$ at low temperature is obtained by substituting (7.61) into (7.48). It can be shown that the condition (7.51) holds.

C. “Determining $M_\pi^2(X)$ and $M_\sigma^2(X)$ ”

In this subsection, following the method 1) in Section VID, we derive the gap equation (the determining equation for $M_\xi^2(X)$ ($\xi = \pi, \sigma$)) to the leading one-loop order. Main task is to compute $\Sigma_R(X; R)$. Let $\mathcal{D}[\mathcal{L}_1, \mathcal{L}_2, \dots]$ ($\mathcal{L}_j \in \mathcal{L}_{int} + \mathcal{L}_{rc}$) be an one-loop diagram contributing to $\Sigma_R(X; R)$, which includes the vertices coming from $\mathcal{L}_1, \mathcal{L}_2, \dots$. The relevant ones are

I. $\mathcal{D}[-\lambda(\vec{\phi}^2)^2/4!]$ (tadpole diagram).

II. $\mathcal{D}[-\lambda(t\vec{\varphi} \cdot \vec{\phi})\vec{\phi}^2/3!, -\lambda(t\vec{\varphi} \cdot \vec{\phi})\vec{\phi}^2/3!]$.

Including the contributions coming from $\mathcal{L}_{rc}$ in (6.10), we write $\Sigma_R(X; R)$, with obvious notation, in the form

$$\begin{aligned} \Sigma_R(X; R) &= (\Sigma_R(X))_I + (\Sigma_R(X; R))_{II} + (Z_\mu Z - 1)\mu^2 \\ &\quad + \frac{\lambda}{6}(Z_\lambda Z^2 - 1)\varphi^2(X) (\mathbf{P}_\pi(X) + 3\mathbf{P}_\sigma(X)) - \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X)\chi_\xi(X). \end{aligned} \quad (7.62)$$

The leading contribution to (7.62) takes the form

$$\begin{aligned} \Sigma_R(X; R) &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X)\Sigma_R^{(\xi)}(X; R) \\ &= \sum_{\xi=\pi, \sigma} \mathbf{P}_\xi(X) \left[\left(\Sigma_R^{(\xi)}(X) \right)_I + \left(\Sigma_R^{(\xi)}(X; R) \right)_{II} + (Z_\mu Z - 1)\mu^2 - \chi_\xi(X) \right. \\ &\quad \left. + \frac{\lambda}{6} d_\xi (Z_\lambda Z^2 - 1)\varphi^2(X) \right], \end{aligned} \quad (7.63)$$

with $d_\pi = 1$ and $d_\sigma = 3$. The resummation program completes by imposing the conditions

$$Re \Sigma_R^{(\xi)}(X; R^2 = \mathcal{M}_\xi^2(X)) = 0 \quad (\xi = \pi, \sigma). \quad (7.64)$$

This is the gap equation, which serves as determining $M_\xi^2(X)$ ($\xi = \pi, \sigma$).

1. Computation of $(\Sigma_R(X))_I$

We start with computing the leading $O(\lambda)$ contribution from the tadpole diagram, $(\Sigma_R(X))_I$:

$$\begin{aligned} (\Sigma_R(X))_I &= \frac{i\lambda}{6} \int \frac{d^4 P}{(2\pi)^4} \left[2 \left\{ \mathbf{P}_\pi(X)\Delta_{11}^{(\pi)}(X; P) + \mathbf{P}_\sigma(X)\Delta_{11}^{(\sigma)}(X; P) \right\} \right. \\ &\quad \left. + \mathbf{I} \left\{ (N-1)\Delta_{11}^{(\pi)}(X; P) + \Delta_{11}^{(\sigma)}(X; P) \right\} \right], \end{aligned} \quad (7.65)$$

where $\Delta_{11}^{(\pi)}$ is the (1,1) component of $\hat{\Delta}^{(\pi)}$ in (6.44), etc. Note that $\hat{\Delta}^{(1)}$ in (6.42) yields higher-order contribution to $\Sigma_R(X)$, which is pure imaginary and consists of terms that are proportional to $(\partial|\varphi(X)\rangle/\partial X^\mu)\langle\varphi(X)|$ and $|\varphi(X)\rangle(\partial\langle\varphi(X)|/\partial X^\mu)$. When one performs two-loop calculation, this contribution should be taken into account. We rewrite (7.65) as

$$\begin{aligned} (\Sigma_{11}(X))_I &= \mathbf{P}_\pi(X) \left(\Sigma_R^{(\pi)}(X) \right)_I + \mathbf{P}_\sigma(X) \left(\Sigma_R^{(\sigma)}(X) \right)_I, \\ \left(\Sigma_R^{(\xi)}(X) \right)_I &= \frac{\lambda}{6} [\alpha_\xi \mathcal{I}_\pi(X) + \beta_\xi \mathcal{I}_\sigma(X)] \quad (\xi = \pi, \sigma) \\ &\quad ((\alpha_\pi, \beta_\pi) = (N+1, 1), (\alpha_\sigma, \beta_\sigma) = (N-1, 3)), \end{aligned} \quad (7.66)$$

where

$$\mathcal{I}_\xi(X) \equiv i \int \frac{d^4 P}{(2\pi)^4} \Delta_{11}^{(\xi)}(X; P). \quad (7.67)$$

The $O(\lambda)$ contribution to $\mathcal{I}_\xi(X)$ has already been computed in the above subsections.

2. Computation of $Re(\Sigma_R(X; R))_{\text{II}}$

Here we compute the leading $O(\lambda^2)$ contribution to $Re(\Sigma_R(X; R))_{\text{II}}$. Straightforward manipulation yields

$$\begin{aligned} (\Sigma_R(X; R))_{\text{II}} &= \mathbf{P}_\pi(X) \left(\Sigma_R^{(\pi)}(X; R) \right)_{\text{II}} + \mathbf{P}_\sigma(X) \left(\Sigma_R^{(\sigma)}(X; R) \right)_{\text{II}}, \\ \left(\Sigma_R^{(\pi)}(X; R) \right)_{\text{II}} &= \frac{\lambda^2}{9} \varphi^2(X) \mathcal{J}_{\pi\sigma}(X; R), \\ \left(\Sigma_R^{(\sigma)}(X; R) \right)_{\text{II}} &= \frac{\lambda^2}{18} \varphi^2(X) [(N-1) \mathcal{J}_{\pi\pi}(X; R) + 9 \mathcal{J}_{\sigma\sigma}(X; R)]. \end{aligned}$$

Here

$$\begin{aligned} \mathcal{J}_{\xi\zeta}(X; R) &= i \int \frac{d^4 P}{(2\pi)^4} \left[\Delta_{11}^{(\xi)}(X; P) \Delta_{11}^{(\zeta)}(X; R-P) - \Delta_{12}^{(\xi)}(X; P) \Delta_{12}^{(\zeta)}(X; R-P) \right] \\ &\equiv \mathcal{J}_{\xi\zeta}^{(0)}(X; R) + \mathcal{J}_{\xi\zeta}^{(\beta)}(X; R) \quad (\xi, \zeta = \pi, \sigma), \end{aligned}$$

with

$$\begin{aligned} \mathcal{J}_{\xi\zeta}^{(0)}(X; R) &= i \int \frac{d^D P}{(2\pi)^D} \left[\frac{1}{P^2 - \mathcal{M}_\xi^2(X) + i\eta} \frac{1}{(R-P)^2 - \mathcal{M}_\zeta^2(X) + i\eta} \right. \\ &\quad \left. + 4\pi^2 \theta(-p_0) \theta(p_0 - r_0) \delta(P^2 - \mathcal{M}_\xi^2(X)) \delta((R-P)^2 - \mathcal{M}_\zeta^2(X)) \right], \\ \mathcal{J}_{\xi\zeta}^{(\beta)}(X; R) &= \int \frac{d^4 P}{(2\pi)^3} \left[\frac{1}{P^2 - \mathcal{M}_\xi^2(X) + i\epsilon(p_0)\eta} N^{(\zeta)}(X; |r_0 - p_0|; \hat{\mathbf{p}}) \delta((R-P)^2 - \mathcal{M}_\zeta^2(X)) \right. \\ &\quad \left. + \frac{1}{(R-P)^2 - \mathcal{M}_\zeta^2(X) - i\epsilon(p_0 - r_0)\eta} N^{(\xi)}(X; |p_0|; \hat{\mathbf{p}}) \delta(P^2 - \mathcal{M}_\xi^2(X)) \right]. \end{aligned}$$

Here we have dropped the contributions that have to be dealt with in two-loop order (cf. above after (7.65)).

Calculation of $\mathcal{J}_{\xi\zeta}^{(0)}$ with $\mathcal{M}_\xi^2(X), \mathcal{M}_\zeta^2(X) \geq 0$ is straightforward. The resultant form for $\mathcal{J}_{\xi\zeta}^{(0)}$ may be analytically continued to other region than $\mathcal{M}_\xi^2(X), \mathcal{M}_\zeta^2(X) \geq 0$:

$$\begin{aligned} -(4\pi)^2 \mathcal{J}_{\pi\sigma}^{(0)}(X; R) &= \frac{2}{\epsilon} - \ln \frac{\mathcal{M}_\sigma^2(X)}{e\mu_d^2} + \frac{R^2 - \mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X)}{2R^2} \ln \frac{\mathcal{M}_\sigma^2(X)}{\mathcal{M}_\pi^2(X)} \\ &\quad + \frac{\mathcal{S}_1}{2R^2} \left[\ln \frac{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) + R^2 - \mathcal{S}_1}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) + R^2 + \mathcal{S}_1} \right. \\ &\quad \left. + \ln \frac{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - R^2 + \mathcal{S}_1}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - R^2 - \mathcal{S}_1} \right], \end{aligned} \tag{7.68}$$

where $\mathcal{M}_\xi^2(X)$ is understood to be $\mathcal{M}_\sigma^2(X) - i\epsilon(r_0)\eta$ and

$$\mathcal{S}_1 \equiv \sqrt{(R^2 - \mathcal{M}_\pi^2(X))^2 - 2\mathcal{M}_\sigma^2(X)(R^2 + \mathcal{M}_\pi^2(X)) + \mathcal{M}_\sigma^4(X)}.$$

Equation (7.68) is valid in the regions,

R1) $\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X) \geq 0$ and

$$\begin{aligned} R^2 &\leq \mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X) - 2\sqrt{\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X)} \\ \text{or} \\ \mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X) + 2\sqrt{\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X)} &\leq R^2. \end{aligned}$$

R2) $\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X) < 0$.

In the region,

$$\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - |R^2| \geq 0, \quad \mathcal{M}_\sigma^2(X), \mathcal{M}_\pi^2(X) \geq 0,$$

Equation (7.68) is as it is. The expression in the regions $\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - |R^2| < 0$ and/or $\mathcal{M}_\sigma^2(X), \mathcal{M}_\pi^2(X) < 0$ is obtained from (7.68) through simple analytic continuation.

Continuing (7.68) to the region

$$\mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X) - 2\sqrt{\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X)} < R^2 < \mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X) + 2\sqrt{\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X)} \\ (\mathcal{M}_\sigma^2(X)\mathcal{M}_\pi^2(X) > 0),$$

we obtain

$$\begin{aligned} -(4\pi)^2 \mathcal{J}_{\pi\sigma}^{(0)}(X; R) = & \frac{2}{\bar{\epsilon}} - \ln \frac{\mathcal{M}_\sigma^2(X)}{e\mu_d^2} + \frac{R^2 - \mathcal{M}_\sigma^2(X) + \mathcal{M}_\pi^2(X)}{2R^2} \ln \frac{\mathcal{M}_\sigma^2(X)}{\mathcal{M}_\pi^2(X)} \\ & + \frac{\mathcal{S}_2}{R^2} \left[\arctan \frac{\mathcal{S}_2}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) + R^2} \right. \\ & \left. - \arctan \frac{\mathcal{S}_2}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - R^2} \right], \end{aligned} \quad (7.69)$$

where

$$\mathcal{S}_2 \equiv \sqrt{2\mathcal{M}_\sigma^2(X)(R^2 + \mathcal{M}_\pi^2(X)) - (R^2 - \mathcal{M}_\pi^2(X))^2 - \mathcal{M}_\sigma^4(X)}.$$

Equation (7.69) is valid in the region

$$\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) - |R^2| \geq 0.$$

If $\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) + \tau R^2$ ($\tau = \pm$) in (7.69) is negative, the replacement

$$\arctan \frac{\mathcal{S}_2}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X) + \tau R^2} \longrightarrow \pi - \arctan \frac{\mathcal{S}_2}{\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X) - \tau R^2}$$

should be made.

$\mathcal{J}_{\xi\xi}^{(0)}(R)$ ($\xi = \pi, \sigma$) is obtained from (7.68) and (7.69) by setting $\mathcal{M}_\pi^2(X) = \mathcal{M}_\sigma^2(X) = \mathcal{M}_\xi^2(X)$.

As seen in (7.64), we need $Re \Sigma_R^{(\xi)}(X; R)$ at $R^2 = \mathcal{M}_\xi^2(X)$. As to $\mathcal{J}_{\xi\xi}^{(\beta)}(X; R)$, following [10], we compute it at $R = 0$ for simplicity. The difference between $\mathcal{J}_{\xi\xi}^{(\beta)}(X; R = 0)$ and $\mathcal{J}_{\xi\xi}^{(\beta)}(X; R)$ at $R^2 = \mathcal{M}_\xi^2(X)$ is of higher order. Then, we have

$$\begin{aligned} Re \left(\Sigma_R^{(\pi)} \right)_\Pi & \simeq \frac{\lambda^2}{9} \varphi^2(X) \left[\mathcal{J}_{\pi\sigma}^{(0)}(X; R^2 = \mathcal{M}_\pi^2(X)) + \mathcal{J}_{\pi\sigma}^{(\beta)}(X; R = 0) \right], \\ Re \left(\Sigma_R^{(\sigma)} \right)_\Pi & \simeq \frac{\lambda^2}{18} \varphi^2(X) \left[(N-1) \left\{ \mathcal{J}_{\pi\pi}^{(0)}(X; R^2 = \mathcal{M}_\sigma^2(X)) + \mathcal{J}_{\pi\pi}^{(\beta)}(X; R = 0) \right\} \right. \\ & \left. + 9 \left\{ \mathcal{J}_{\sigma\sigma}^{(0)}(X; R^2 = \mathcal{M}_\sigma^2(X)) + \mathcal{J}_{\sigma\sigma}^{(\beta)}(X; R = 0) \right\} \right], \end{aligned}$$

where

$$\begin{aligned} \mathcal{J}_{\pi\sigma}^{(0)}(X; R^2 = \mathcal{M}_\pi^2(X)) & = -\frac{1}{(4\pi)^2} \left[\frac{2}{\bar{\epsilon}} - \frac{\mathcal{M}_\sigma^2(X)}{2\mathcal{M}_\pi^2(X)} \ln \frac{|\mathcal{M}_\sigma^2(X)|}{|\mathcal{M}_\pi^2(X)|} - \ln \frac{|\mathcal{M}_\pi^2(X)|}{e^2\mu_d^2} + \mathcal{K}_{\pi\sigma}^{(\pi)} \right], \\ \mathcal{J}_{\pi\pi}^{(0)}(X; R^2 = \mathcal{M}_\sigma^2(X)) & = -\frac{1}{(4\pi)^2} \left[\frac{2}{\bar{\epsilon}} - \ln \frac{|\mathcal{M}_\pi^2(X)|}{e^2\mu_d^2} + \mathcal{K}_{\pi\pi}^{(\sigma)} \right], \\ \mathcal{J}_{\sigma\sigma}^{(0)}(X; R^2 = \mathcal{M}_\sigma^2(X)) & = -\frac{1}{(4\pi)^2} \left[\frac{2}{\bar{\epsilon}} - \frac{\pi}{\sqrt{3}} - \ln \frac{|\mathcal{M}_\sigma^2(X)|}{e^2\mu_d^2} \right] \end{aligned} \quad (7.70)$$

with

$$\begin{aligned}
\mathcal{K}_{\pi\sigma}^{(\pi)} &= \theta(\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))) \frac{\sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}}{2\mathcal{M}_\pi^2(X)} \\
&\times \left[\ln \frac{|\mathcal{M}_\sigma^2(X) - \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|}{|\mathcal{M}_\sigma^2(X) + \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|} \right. \\
&- \ln \frac{|\mathcal{M}_\sigma^2(X) - 2\mathcal{M}_\pi^2(X) - \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|}{|\mathcal{M}_\sigma^2(X) - 2\mathcal{M}_\pi^2(X) + \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|} \Big], \\
&+ \theta(\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))) \frac{\sqrt{\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))}}{\mathcal{M}_\pi^2(X)} \\
&\times \left[\epsilon(\mathcal{M}_\sigma^2(X)) \arctan \frac{\sqrt{\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))}}{|\mathcal{M}_\sigma^2(X)|} \right. \\
&- \epsilon(\mathcal{M}_\sigma^2(X) - 2\mathcal{M}_\pi^2(X)) \arctan \frac{\sqrt{\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))}}{|\mathcal{M}_\sigma^2(X) - 2\mathcal{M}_\pi^2(X)|} \\
&+ \pi \{ \theta(-\mathcal{M}_\sigma^2(X)) - \theta(2\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X)) \} \Big], \tag{7.71}
\end{aligned}$$

$$\begin{aligned}
\mathcal{K}_{\pi\pi}^{(\sigma)} &= \theta(\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))) \\
&\times \frac{\sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}}{\mathcal{M}_\sigma^2(X)} \ln \frac{|\mathcal{M}_\sigma^2(X) - \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|}{|\mathcal{M}_\sigma^2(X) + \sqrt{\mathcal{M}_\sigma^2(X)(\mathcal{M}_\sigma^2(X) - 4\mathcal{M}_\pi^2(X))}|} \\
&+ \theta(\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))) \frac{\sqrt{\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))}}{|\mathcal{M}_\sigma^2(X)|} \\
&\times \left[2 \arctan \frac{\sqrt{\mathcal{M}_\sigma^2(X)(4\mathcal{M}_\pi^2(X) - \mathcal{M}_\sigma^2(X))}}{|\mathcal{M}_\sigma^2(X)|} - \pi \right], \tag{7.72}
\end{aligned}$$

and

$$\mathcal{J}_{\pi\sigma}^{(\beta)}(X; R=0) = \frac{\mathcal{I}_\sigma^{(\beta)}(X) - \mathcal{I}_\pi^{(\beta)}(X)}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X)}, \tag{7.73}$$

$$\mathcal{J}_{\xi\xi}^{(\beta)}(X; R=0) = \frac{\partial \mathcal{I}_\xi^{(\beta)}(X)}{\partial \mathcal{M}_\xi^2(X)}. \tag{7.74}$$

Here $\mathcal{I}_\xi^{(\beta)}(X)$ is as in (7.11) or (7.12). Expressions (7.70), (7.71), and (7.72) do not diverge at $\mathcal{M}_\pi^2(X) = 0$ and at $\mathcal{M}_\sigma^2(X) = 0$. Equations (7.73) and (7.74) are valid for $\mathcal{M}_\xi^2(X) \geq 0$ ($\xi = \pi, \sigma$). Analytic continuation of them to the region $\mathcal{M}_\xi^2(X) < 0$ may be performed as in subsection B:

$$\mathcal{I}_\xi^{(\beta)}(X) \rightarrow \frac{1}{2\pi^2} \text{Re} \int_{C'} dz \sqrt{z^2 - \mathcal{M}_\xi^2(X)} \langle N^{(\xi)}(X; z) \rangle, \tag{7.75}$$

where the integration contour C' is as in (7.44).

3. Gap equation

Let us turn back to the condition (7.64), which may be written as

$$\begin{aligned}
\chi_\xi &= \mathcal{M}_\xi^2(X) - \mu^2 \\
&= \left(\Sigma_R^{(\xi)}(X) \right)_I + \left(\Sigma_R^{(\xi)}(X) \right)_{II} + (Z_\mu Z - 1) \mu^2 + \frac{\lambda}{6} d_\xi (Z_\lambda Z^2 - 1) \varphi^2(X) \quad (\xi = \pi, \sigma). \tag{7.76}
\end{aligned}$$

The UV-diverging contributions in (7.76) cancels. In fact, substituting (7.19) into (7.76), we obtain the UV-divergence free equation,

$$\left(\tilde{M}_\xi^2(X)\right)_I + \left(\tilde{M}_\xi^2(X)\right)_{II} = M_\xi^2(X) - \mu^2 \quad (\xi = \pi, \sigma). \quad (7.77)$$

Here (cf. (7.66))

$$\left(\tilde{M}_\xi^2(X)\right)_I = \frac{\lambda}{6} \left[\alpha_\xi \tilde{\mathcal{I}}_\pi + \beta_\xi \tilde{\mathcal{I}}_\sigma \right] \quad (7.78)$$

with

$$\tilde{\mathcal{I}}_\xi = \frac{1}{(4\pi)^2} \mathcal{M}_\xi^2(X) \ln \frac{|\mathcal{M}_\xi^2(X)|}{e\mu_d^2} + \text{Re} \mathcal{I}_\xi^{(\beta)}(X) \quad (7.79)$$

and

$$\begin{aligned} \left(\tilde{M}_\pi^2(X)\right)_{II} = & \frac{\lambda^2}{9} \varphi^2(X) \left[\frac{1}{(4\pi)^2} \left\{ \frac{\mathcal{M}_\sigma^2(X)}{2\mathcal{M}_\pi^2(X)} \ln \frac{|\mathcal{M}_\sigma^2(X)|}{|\mathcal{M}_\pi^2(X)|} + \ln \frac{|\mathcal{M}_\pi^2(X)|}{e^2\mu_d^2} - \mathcal{K}_{\pi\sigma}^{(\pi)} \right\} \right. \\ & \left. + \text{Re} \frac{\mathcal{I}_\sigma^{(\beta)}(X) - \mathcal{I}_\pi^{(\beta)}(X)}{\mathcal{M}_\sigma^2(X) - \mathcal{M}_\pi^2(X)} \right], \end{aligned} \quad (7.80)$$

$$\begin{aligned} \left(\tilde{M}_\sigma^2(X)\right)_{II} = & \frac{\lambda^2}{18} \varphi^2(X) \left[\frac{1}{(4\pi)^2} \left\{ (N-1) \left(\ln \frac{|\mathcal{M}_\pi^2(X)|}{e^2\mu_d^2} - \mathcal{K}_{\pi\pi}^{(\sigma)} \right) \right. \right. \\ & \left. \left. + 3\sqrt{3}\pi + 9 \ln \frac{|\mathcal{M}_\sigma^2(X)|}{e^2\mu_d^2} \right\} + \text{Re} \left\{ (N-1) \frac{\partial \mathcal{I}_\pi^{(\beta)}(X)}{\partial \mathcal{M}_\pi^2(X)} + 9 \frac{\partial \mathcal{I}_\sigma^{(\beta)}(X)}{\partial \mathcal{M}_\sigma^2(X)} \right\} \right]. \end{aligned} \quad (7.81)$$

In the above equations, $\mathcal{I}_\xi^{(\beta)}(X)$ is as in (7.12) for $\mathcal{M}_\xi^2(X) \geq 0$ and as in (7.75) for $\mathcal{M}_\xi^2(X) < 0$. $\mathcal{K}_{\pi\sigma}^{(\pi)}$ and $\mathcal{K}_{\pi\pi}^{(\sigma)}$ are as in (7.71) and (7.72), respectively. For the region,

$$\mathcal{M}_\sigma^2(X) \geq 0, \quad \mathcal{M}_\sigma^2(X) \geq 4\mathcal{M}_\pi^2(X)$$

or

$$\mathcal{M}_\sigma^2(X) < 0, \quad \mathcal{M}_\sigma^2(X) < 4\mathcal{M}_\pi^2(X),$$

the first terms on the RHS of (7.71) and of (7.72) should be used and the second terms should be used for the region,

$$\mathcal{M}_\sigma^2(X) \geq 0, \quad \mathcal{M}_\sigma^2(X) < 4\mathcal{M}_\pi^2(X)$$

or

$$\mathcal{M}_\sigma^2(X) < 0, \quad \mathcal{M}_\sigma^2(X) > 4\mathcal{M}_\pi^2(X).$$

In the LHS of (7.77), we have dropped the higher-order term,

$$- \frac{\lambda}{48\pi^2} \frac{1}{\epsilon} [c_\xi \chi_\pi + d_\xi \chi_\sigma], \quad (7.82)$$

which should be dealt with when one computes next to the leading order. The one-loop diagram $\mathcal{D} \left[{}^t\vec{\phi}\chi_\xi \mathbf{P}_\xi \vec{\phi}/2, -\lambda(\vec{\phi}^2)^2/4! \right]$, which contributes at next to the leading order, cancels the diverging contribution (7.82).

Equation (7.77) is the gap equation, by which $M_\xi^2(X)$ ($\xi = \pi, \sigma$) are determined self consistently. $M_\xi^2(X)$ thus determined depend on $\vec{\varphi}^2(X)$. With $M_\xi^2(X)$ in hand, one can judge if $\mathcal{M}_\xi^2(X)$ in (6.29) is positive or negative.

4. “High-temperature” expansion

“High-temperature” expansion of $\mathcal{I}_\xi^{(\beta)}(X)$, Eq. (7.75), is obtained as in subsections A and B,

$$\text{Re} \mathcal{I}_\xi^{(\beta)}(X) \simeq \frac{1}{12} c_\xi(X) \mathcal{T}_\xi^2(X) - \frac{1}{4\pi} \theta(\mathcal{M}_\xi^2(X)) (\mathcal{M}_\xi^2(X))^{1/2} \mathcal{T}_\xi(X). \quad (7.83)$$

From this the “high-temperature” expansion of $\partial \mathcal{I}_\xi^{(\beta)}(X)/\partial \mathcal{M}_\xi^2(X)$ ($\xi = \pi, \sigma$) in (7.81) is obtained:

$$\frac{\partial \mathcal{I}_\xi^{(\beta)}(X)}{\partial \mathcal{M}_\xi^2(X)} \simeq -\frac{1}{8\pi} \theta(\mathcal{M}_\xi^2(X)) (\mathcal{M}_\xi^2(X))^{-1/2} \mathcal{T}_\xi(X).$$

When $\mathcal{M}_\xi^2(X) < 0$, $\partial \mathcal{I}_\xi^{(\beta)}(X)/\partial \mathcal{M}_\xi^2(X)$ is negligibly small.

5. “Low-temperature” expansion

In the case of locally thermal equilibrium system at low temperature, we have, for $\mathcal{M}_\xi^2(X) \geq 0$,

$$\mathcal{I}_\xi^{(\beta)}(X) \simeq \frac{1}{2^{3/2}\pi} (\mathcal{M}_\xi^2(X))^{1/4} (T(X))^{3/2} e^{-\mathcal{M}_\xi^2(X)/T(X)} \quad (\mathcal{M}_\xi^2(X) \geq 0). \quad (7.84)$$

For $\mathcal{M}_\xi^2(X) < 0$, computation may be carried out as in subsection B. The result agrees with that obtained from (7.84) by an analytic continuation;

$$\text{Re } \mathcal{I}_\xi^{(\beta)}(X) \simeq \frac{1}{2^{3/2}\pi} (-\mathcal{M}_\xi^2(X))^{1/4} (T(X))^{3/2} \cos\left(\frac{(-\mathcal{M}_\xi(X))^{1/2}}{T(X)} - \frac{\pi}{4}\right) \quad (\mathcal{M}_\xi^2(X) < 0). \quad (7.85)$$

From (7.84) and (7.85), the low-temperature expansion of $\partial \mathcal{I}_\xi^{(\beta)}(X)/\partial \mathcal{M}_\xi^2(X)$ ($\xi = \pi, \sigma$) in (7.81) is obtained:

$$\begin{aligned} \text{Re } \frac{\partial \mathcal{I}_\xi^{(\beta)}(X)}{\partial \mathcal{M}_\xi^2(X)} \simeq & -\frac{1}{4\sqrt{2}\pi} \left[\theta(\mathcal{M}_\xi^2(X)) (\mathcal{M}_\xi^2(X))^{-1/4} (T(X))^{1/2} e^{-\mathcal{M}_\xi^2(X)/T(X)} \right. \\ & \left. - \theta(-\mathcal{M}_\xi^2(X)) \sin\left(\frac{(-\mathcal{M}_\xi(X))^{1/2}}{T(X)} - \frac{\pi}{4}\right) (-\mathcal{M}_\xi^2(X))^{-1/4} (T(X))^{1/2} \right]. \end{aligned}$$

VIII. APPLICATION — EARLY STAGE OF PHASE TRANSITION

There are few cases, for which approximate analytic calculations yield reliable results. In this section, for the purpose of illustrating how the formalism constructed above works, we apply it to very simple cases, in which the system of coupled equations obtained in the previous section can approximately be solved analytically.

A. Spatially uniform system

In this subsection, we apply the formalism to a spatially uniform system. We suppose the following situation. At the initial time $X_0 = 0$, i) the system is in a “highly super-cooled state” with $\vec{\varphi}(X_0 = 0) \simeq 0$ and ii) the temperature T is still high and the system is in quasi thermal equilibrium,

$$N^{(\xi)}(X_0 = 0; |p_0|) = \frac{1}{e^{|p_0|/T} - 1}. \quad (8.1)$$

We are only interested in the small time interval, during which iii) the high-temperature expansion is applicable, so that keeping terms that “explicitly” depend on T is a good approximation (cf. below) and iv) the condensate fields $\vec{\varphi}(X_0)$ remains small in the sense stated below. As mentioned above, our formalism applies to the system away from the critical region, which is the reason to include i) into the list.

We start with the determining equation for $f^{(\xi)}(X_0; p_0) = \theta(p_0)N^{(\xi)}(X_0; p_0) - \theta(-p_0)(1 + N^{(\xi)}(X_0; -p_0))$ ($\xi = \pi, \sigma$), Eq. (6.53),

$$2p_0 \frac{\partial N^{(\xi)}(X_0; p_0)}{\partial X_0} + \frac{\partial N^{(\xi)}(X_0; p_0)}{\partial p_0} \frac{d\mathcal{M}_\xi(X_0)}{dX_0} = 0. \quad (8.2)$$

Since $N^{(\xi)}(X_0; p_0)$ in the propagator accompanies $\delta(P^2 - \mathcal{M}_\xi^2(X_0))$, p_0 in (8.2) is $p_0 = \sqrt{p^2 + \mathcal{M}_\xi^2(X_0)}$. Equation (8.2) may be solved as

$$N^{(\xi)}(X_0; p_0) = \mathcal{F}(P^2 - \mathcal{M}_\xi^2(X_0)), \quad (8.3)$$

where $\mathcal{F}$ is an arbitrary function. For simplicity of presentation, we take $\vec{\varphi}(X_0 = 0) = d\vec{\varphi}(X_0)/dX_0 \Big|_{X_0=0} = 0$ in the

sequel. In the case of $\vec{\varphi}(X_0 = 0) \neq 0$ and/or $d\vec{\varphi}(X_0)/dX_0 \Big|_{X_0=0} \neq 0$, no new feature arises, as long as they are small.

Using (8.1) in (8.3), we obtain,

$$\begin{aligned}
N^{(\xi)}(X_0; p_0) &= \frac{1}{e^{\sqrt{p_0^2 - \mathcal{M}_\xi^2(X_0) + M_\xi^2(X_0=0)/T}} - 1} \\
&= \frac{1}{e^{\sqrt{p^2 + M_\xi^2(0)/T}} - 1},
\end{aligned} \tag{8.4}$$

where use has been made of the fact that $N^{(\xi)}(X_0; p_0)$ accompanies $\delta(P^2 - \mathcal{M}_\xi^2(X_0))$ in the propagator.

With (8.4), let us compute the fundamental quantity $\mathcal{I}_\xi(X_0)$, Eq. (7.9). Computation goes in a similar manner as the “direct calculation” in Section VIII B:

$$\begin{aligned}
\mathcal{I}_\xi^{(\beta)}(X_0) &= \frac{T^2}{12} + \frac{1}{2\pi^2} \left[\int_0^\infty dp \frac{p^2}{\sqrt{p^2 + \mathcal{M}_\xi^2(X_0)}} \frac{1}{e^{\sqrt{p^2 + M_\xi^2(0)/T}} - 1} \right. \\
&\quad \left. - \int_0^\infty d\mathcal{E} \frac{\mathcal{E}}{e^{\mathcal{E}/T} - 1} \right] \\
&\simeq \frac{T^2}{12} + \frac{T}{2\pi^2} \left[\int_0^\infty dp \frac{p^2}{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)|} \sqrt{p^2 - |M_\xi^2(0)|}} \right. \\
&\quad \left. - \int_{\sqrt{|M_\xi^2(0)|}}^\infty dp \frac{p}{\sqrt{p^2 - |M_\xi^2(0)|}} \right],
\end{aligned} \tag{8.5}$$

where high- T approximation has been used. Noticing that,

$$\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)|} = -i\sqrt{|\mathcal{M}_\xi^2(X_0)| - p^2} \quad \text{for } p^2 < |\mathcal{M}_\xi^2(X_0)|,$$

etc., we obtain

$$\begin{aligned}
\mathcal{I}_\xi^{(\beta)}(X_0) &\simeq \frac{T^2}{12} - \frac{\sqrt{|M_\xi^2(0)|} T}{2\pi^2} H\left(\frac{|\mathcal{M}_\xi^2(X_0)|}{|M_\xi^2(0)|}\right) \\
&\quad + \frac{iT}{2\pi^2} \sqrt{|M_\xi^2(0)|} E\left(\frac{\pi}{2}, \sqrt{\frac{|\mathcal{M}_\xi^2(X_0)| - |M_\xi^2(0)|}{|M_\xi^2(0)|}}\right),
\end{aligned} \tag{8.6}$$

where

$$\begin{aligned}
H(\rho^2) &\equiv \int_0^\rho dx \frac{x^2}{\sqrt{1-x^2}\sqrt{\rho^2-x^2}} - \int_1^\infty dx \frac{x}{\sqrt{x^2-1}} \left[\frac{x}{\sqrt{x^2-\rho^2}} - 1 \right], \\
E(\pi/2, k) &\equiv \int_0^{\pi/2} d\theta \sqrt{1 - k^2 \sin^2 \theta}.
\end{aligned}$$

Here E is an elliptic function of second kind. Note that $|M_\xi^2(0)| > |\mathcal{M}_\xi^2(X_0)|$, so that $\rho < 1$. From the setup iv) above,

$$|M_\xi^2(0)| - |\mathcal{M}_\xi^2(X_0)| = \lambda e_\xi \vec{\varphi}^2(X_0) \ll |M_\xi^2(0)| \quad (e_\pi = 1/6, e_\sigma = 1/2),$$

and then (8.6) may be approximated as

$$\begin{aligned}
\mathcal{I}_\xi^{(\beta)}(X_0) &\simeq \frac{T^2}{12} - \frac{1}{4\pi^2} \frac{T}{\sqrt{|M_\xi^2(0)|}} [|M_\xi^2(0)| - |\mathcal{M}_\xi^2(X_0)|] \\
&\quad + \frac{iT}{4\pi} \left[\sqrt{|\mathcal{M}_\xi^2(X_0)|} + \frac{1}{4} \frac{|M_\xi^2(0)| - |\mathcal{M}_\xi^2(X_0)|}{\sqrt{|M_\xi^2(0)|}} \right] \\
&= \frac{T^2}{12} - e_\xi \frac{\lambda}{4\pi^2} \frac{T}{\sqrt{|M_\xi^2(0)|}} \vec{\varphi}^2(X_0) + \frac{iT}{4\pi} \left[\sqrt{|\mathcal{M}_\xi^2(X_0)|} + \frac{\lambda e_\xi}{4} \frac{\vec{\varphi}^2(X_0)}{\sqrt{|M_\xi^2(0)|}} \right].
\end{aligned} \tag{8.7}$$

Let us turn to the determining equation for $M_\xi(X_0)$ ($\xi = \pi, \sigma$), Eq. (7.77). As mentioned above, we ignore the temperature-independent terms. Using (8.7) in (7.79) - (7.81), we see that (7.77) yields

$$M_\pi^2(X_0) \simeq -\omega^2 - \left(\frac{\lambda}{12\pi}\right)^2 (N+4) \frac{T}{\sqrt{|M_\pi^2(0)|}} \vec{\varphi}^2(X_0) - \frac{\lambda^3}{108\pi^2} \frac{T}{\sqrt{|M_\pi^2(0)|}} \frac{(\vec{\varphi}^2(X_0))^2}{M_\sigma^2(X_0) - M_\pi^2(X_0) + \lambda \vec{\varphi}^2(X_0)/3}, \quad (8.8)$$

$$M_\sigma^2(X_0) \simeq -\omega^2 - 3 \left(\frac{\lambda}{12\pi}\right)^2 (N+8) \frac{T}{\sqrt{|M_\pi^2(0)|}} \vec{\varphi}^2(X_0), \quad (8.9)$$

where

$$\omega^2 \equiv -\mu^2 - \frac{\lambda}{72} (N+2) T^2 (> 0).$$

In obtaining (8.8) and (8.9), use has been made of the approximation $(|M_\sigma^2(0)| - |M_\pi^2(0)|) \ll |M_\pi^2(0)|$, which will be justified a posteriori.

Approximate solution to (8.8) may be written in the form

$$M_\pi^2(X_0) \simeq -\omega^2 - \left(\frac{\lambda}{12\pi}\right)^2 (N+g) \frac{T}{\sqrt{|M_\pi^2(0)|}} \vec{\varphi}^2(X_0), \quad (8.10)$$

where

$$g = 4 \left[1 + \left\{ 1 - 3 \left(\frac{1}{12\pi}\right)^2 (2N+24-g) \frac{\lambda T}{\sqrt{|M_\pi^2(0)|}} \right\}^{-1} \right].$$

For $\lambda T < O(\sqrt{|M_\pi^2(0)|})$, $g = 8$ and, for $\lambda T > O(\sqrt{|M_\pi^2(0)|})$, $g = 4$.

Let us turn to deriving the equation of motion for $\vec{\varphi}(X)$. $\Omega(\vec{\varphi}^2(X_0))$ in (7.24) reads

$$\Omega(\vec{\varphi}^2(X_0)) \simeq -\omega^2 + \frac{\lambda}{6} \vec{\varphi}^2(X_0) - \frac{\lambda T}{12\pi^2} \left[(N-1) \sqrt{|M_\pi^2(0)|} H\left(\frac{|M_\pi^2(X_0)|}{|M_\pi^2(0)|}\right) + 3 \sqrt{|M_\sigma^2(0)|} H\left(\frac{|M_\sigma^2(X_0)|}{|M_\sigma^2(0)|}\right) \right].$$

Up to $O(\vec{\varphi}^2)$ term, this becomes

$$\Omega(\vec{\varphi}^2(X_0)) \simeq -\omega^2 + \frac{\lambda'}{6} \vec{\varphi}^2(X_0),$$

where

$$\lambda' \equiv \lambda \left[1 - \frac{\lambda}{24\pi^2} (N+8) \frac{T}{\omega} \right]. \quad (8.11)$$

It should be emphasized that $\lambda' < \lambda$. Above equations are valid for $\lambda T < O(\omega)$ (cf. the setup i) at the beginning of this subsection). The equation of motion for $\vec{\varphi}(X_0)$, Eq. (7.54), reads

$$\ddot{\vec{\varphi}}(X_0) = \left[\omega^2 - \frac{\lambda'}{6} \vec{\varphi}^2(X_0) \right] \vec{\varphi}(X_0) + \vec{\zeta}(X_0) + H \vec{e} \quad (8.12)$$

$$= -\frac{\partial \mathcal{V}}{\partial \vec{\varphi}(X_0)} + \vec{\zeta}(X_0), \quad (8.13)$$

where $\ddot{\vec{\varphi}}(X_0) \equiv d^2 \vec{\varphi}(X_0)/dX_0^2$ and

$$\mathcal{V} \simeq \frac{\pi^2}{90} N T^4 - \frac{\omega^2}{2} \vec{\varphi}^2(X_0) + \frac{\lambda'}{4!} (\vec{\varphi}^2(X_0))^2 - H(\vec{e} \cdot \vec{\varphi}(X_0)). \quad (8.14)$$

is related to γ_{phys} through

$$\mathcal{V} = -Re \gamma_{phys} + \frac{1}{2} \dot{\vec{\varphi}}^2(X_0)$$

with $\dot{\vec{\varphi}}(X_0) \equiv d\vec{\varphi}(X_0)/dX_0$. In the following we simply call $\mathcal{V}$ the potential. From the setup i) at the beginning of this section, $\omega^2 \simeq -M_\pi^2(0) \simeq -M_\sigma^2(0)$ (> 0) is quite large. The setup iv) means that we are interested in the time interval during which $\omega^2 \gg \lambda' \vec{\varphi}^2(X_0)$. [Recalling that X_0 of $\vec{\varphi}(X_0)$ is macroscopic time coordinate, ω should be $\omega \lesssim 1/L_0$, where L_0 is the size of the time-direction of the spacetime cell (cf. Section I).] Above observation allows us to solve (8.12) iteratively. Ignoring the term $\lambda' \vec{\varphi}^2(X_0) \vec{\varphi}(X_0)/6$, we obtain the zeroth-order solution, $\vec{\varphi}^{(0)}(X_0)$, of (8.12),

$$\vec{\varphi}^{(0)}(X_0) = \frac{H}{\omega^2} [\cosh(\omega X_0) - 1] \vec{e} + \frac{1}{\omega} \int_0^{X_0} dX'_0 \vec{\zeta}(X'_0) \sinh(\omega(X_0 - X'_0)). \quad (8.15)$$

Substituting (8.15) back into the term $\lambda' \vec{\varphi}^2(X_0) \vec{\varphi}(X_0)/6$ in (8.12), we obtain the first-order solution,

$$\begin{aligned} \vec{\varphi}(X_0) &= \frac{H}{\omega^2} [\cosh(\omega X_0) - 1] \vec{e} \\ &+ \frac{1}{\omega} \int_0^{X_0} dX'_0 \left[\vec{\zeta}(X'_0) - \frac{\lambda'}{6} \left(\vec{\varphi}^{(0)}(X'_0) \right)^2 \vec{\varphi}^{(0)}(X'_0) \right] \sinh(\omega(X_0 - X'_0)), \end{aligned} \quad (8.16)$$

where $\vec{\varphi}^{(0)}(X'_0)$ is as in (8.15). More accurate solution may be obtained by further iterations.

For the purpose of computing the ζ -averaged correlation functions, we compute the partial derivatives of $Im \gamma_{phys}(X_0)$ with respect to $\vec{\varphi}^2(X_0)$ using (7.24) and (7.25) with (8.7):

$$\left. \frac{\partial Im \gamma_{phys}(X_0)}{\partial \vec{\varphi}^2(X_0)} \right|_{\vec{\varphi}=0} \simeq -\frac{\lambda}{48\pi} (N+2) \omega T, \quad (8.17)$$

$$\left. \frac{\partial^2 Im \gamma_{phys}(X_0)}{\partial (\vec{\varphi}^2(X_0))^2} \right|_{\vec{\varphi}=0} \simeq \frac{\lambda^2}{1152\pi} (N+8) \frac{T}{\omega}. \quad (8.18)$$

Noticing that we are concerned about the spatially uniform system, (7.55) and (7.56) become, in respective order,

$$\langle \zeta_i(X_0) \zeta_j(Y_0) \rangle_\zeta \simeq -\frac{\lambda}{24\pi} (N+2) \frac{\omega T}{V} \delta_{ij} \delta(X_0 - Y_0) \quad (8.19)$$

and

$$\begin{aligned} \langle \zeta_i(X_0) \zeta_j(Y_0) \zeta_k(U_0) \zeta_l(V_0) \rangle_{\zeta c} &\simeq -\frac{\lambda^2}{288\pi} \frac{N+8}{V^3} \frac{T}{\omega} [\delta_{ik} \delta_{lj} + \delta_{kj} \delta_{li} + \delta_{ij} \delta_{lk}] \\ &\times \delta(X_0 - Y_0) \delta(X_0 - U_0) \delta(V_0 - Y_0), \end{aligned} \quad (8.20)$$

where V is the (large) volume of the system.

We are now in a position to compute the ζ -averaged “connected” correlation functions (7.58). From the zeroth-order solution (8.15), we obtain using (8.19) and (8.20),

$$\begin{aligned} \langle \vec{\varphi}^{(0)}(X_0) \rangle_\zeta &= \frac{h}{\omega^2} \vec{e} [\cosh(\omega X_0) - 1], \quad (8.21) \\ \langle \varphi_i^{(0)}(X_0) \varphi_j^{(0)}(Y_0) \rangle_{\zeta c} &= \frac{1}{4\omega^2} \sum_{\rho, \sigma=\pm} \rho \sigma e^{\omega(\rho X_0 - \sigma Y_0)} \\ &\times \int_0^{X_0} dX'_0 \int_0^{Y_0} dY'_0 e^{-\omega(\rho X'_0 - \sigma Y'_0)} \langle \zeta_i(X'_0) \zeta_j(Y'_0) \rangle_\zeta \\ &= -\frac{\lambda}{48\pi} \frac{T}{\omega^2} \frac{\delta_{ij}}{V} (N+2) [\cosh(\omega X_0) \sinh(\omega Y_0) - \omega_0 Y_0 \cosh(\omega(X_0 - Y_0))], \end{aligned} \quad (8.22)$$

and

$$\begin{aligned}
& \langle \varphi_i^{(0)}(X_0) \varphi_j^{(0)}(Y_0) \varphi_k^{(0)}(U_0) \varphi_l^{(0)}(V_0) \rangle_{\zeta c} \\
& \simeq -\frac{N+8}{2\pi} \left(\frac{\lambda}{48} \right)^2 \frac{T}{\omega^6 V^3} [\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}] \\
& \quad \times [\cosh(\omega(X_0 + Y_0 + U_0 - V_0)) \sinh(2\omega V_0) \\
& \quad - 4 \sinh(\omega V_0) \{ \cosh(\omega(X_0 + Y_0 - V_0)) \cosh(\omega(U_0 - V_0)) \\
& \quad + \cosh(\omega(X_0 - Y_0)) \cosh(\omega U_0) \} \\
& \quad + 2\omega V_0 \{ 2 \cosh(\omega(X_0 - V_0)) \cosh(\omega(Y_0 - U_0)) + \cosh(\omega(X_0 - Y_0 - U_0 + V_0)) \}].
\end{aligned} \tag{8.23}$$

In (8.22), $X_0 \geq Y_0$ has been assumed and in (8.23), $X_0, Y_0, U_0 \geq V_0$ has been assumed. Let us study (8.21) - (8.23) in turn.

$$1. \langle \vec{\varphi}^{(0)}(X_0) \rangle_{\zeta}$$

Equation (8.21) shows that, as X_0 increases, the ζ -averaged $\vec{\varphi}^{(0)}(X_0)$ blows up exponentially,

$$\langle \vec{\varphi}^{(0)}(X_0) \rangle_{\zeta} \sim \vec{\Phi} e^{\omega X_0} \quad (\omega X_0 \gg 1), \tag{8.24}$$

where $\vec{\Phi} \equiv H\vec{e}/(2\omega^2)$. We should recall here that, for too large X_0 , $\vec{\varphi}(X_0)$ is large and our approximation does not apply. For obtaining an improved expression for $\langle \vec{\varphi}(X_0) \rangle_{\zeta}$, we should use the improved solution (8.16), which yields, for large $e^{\omega X_0} \gg 1$,

$$\langle \vec{\varphi}(X_0) \rangle_{\zeta} \simeq \vec{\Phi} e^{\omega X_0} - \frac{\lambda'}{48\omega^2} \vec{\Phi} e^{3\omega X_0} \left[\vec{\Phi}^2 - \frac{\lambda}{192\pi} (N+2)^2 \frac{1}{\omega^2} \frac{T}{V} \right]. \tag{8.25}$$

Comparing (8.25) with (8.24), we see that (8.25) is reliable in the region

$$e^{2\omega X_0} < O \left(\frac{\omega^2}{\lambda'} \left| \vec{\Phi}^2 - \frac{\lambda}{192\pi} (N+2)^2 \frac{T}{\omega^2 V} \right|^{-1} \right),$$

which generally forces λ to small values. When compared to the first term on the RHS of (8.25), the term with $\vec{\Phi}^2$ in the second term prevents $|\langle \vec{\varphi}(X_0) \rangle|$ to become large, while the other term expedites it. The former can easily be understood from (8.12), since the term $\lambda' \vec{\varphi}^{(0)2}(X_0) \vec{\varphi}^{(0)}(X_0)/6$ in (8.12) acts as decreasing ω . The latter comes from

$$\frac{\lambda'}{6} \left(\vec{\varphi}^{(0)}(X_0) \right)^2 \varphi_i^{(0)}(X_0) \rightarrow \frac{\lambda'}{6} \langle \left(\vec{\varphi}^{(0)}(X_0) \right)^2 \rangle_{\zeta} \varphi_i^{(0)}(X_0) + \frac{\lambda'}{3} \langle \varphi_j^{(0)}(X_0) \varphi_i^{(0)}(X_0) \rangle_{\zeta} \varphi_j^{(0)}(X_0) \tag{8.26}$$

in (8.12). As seen from (8.22), the ζ -averaged quantities in (8.26) are negative so that, in (8.12), (8.26) acts as increasing $|\vec{\varphi}(X_0)|$ as X_0 increases.

2. Two-point ζ -correlation function

We now turn to (8.22). One can easily see that (8.22) is negative definite. Thus, $Im \gamma_{phys}(X_0)$ causes a negative correlation between the same component of $\vec{\varphi}$, $\varphi_i(X_0)$ and $\varphi_{j=i}(Y_0)$. If the term $\vec{\zeta}(X_0)$ in (8.12), which comes from $Im \gamma_{phys}(X_0)$, is absent, (8.15) or (8.16) with $\vec{\zeta} = 0$ uniquely determines $\vec{\varphi}(X_0)$. $Im \gamma_{phys}$ works as stochastically upsetting $\vec{\varphi}(X_0)$ through the random forces $\vec{\zeta}$. Since $\langle \varphi_i^{(0)}(X_0) \varphi_i^{(0)}(Y_0) \rangle_{\zeta}$ (with no summation over i) in (8.19) is negative, $\langle \varphi_i(X_0) \varphi_{j=i}(Y_0) \rangle_{\zeta}$ tends to decrease when compared to the case where no random force acts, $\vec{\zeta}(X_0) = 0$.

As X_0 increases, the ζ -averaged two-point correlation function (8.22) blows up exponentially. For $X_0 = Y_0$ and $e^{\omega X_0} \gg 1$, we have

$$\langle \varphi_i^{(0)}(X_0) \varphi_j^{(0)}(X_0) \rangle_{\zeta c} \simeq -\frac{\lambda}{192\pi} (N+2) \frac{T}{\omega^2} \frac{\delta_{ij}}{V} e^{2\omega X_0}. \tag{8.27}$$

Using the improved solution (8.16), we may compute the improved ζ -averaged two-point correlation function. Up to the first order with respect to the difference between (8.16) and (8.15), we obtain

$$\langle \varphi_i(X_0) \varphi_j(X_0) \rangle_{\zeta c} \simeq \text{Eq. (8.27)} + \mathcal{C}_{ij}^{(1)}(X_0) + \mathcal{C}_{ij}^{(2)}(X_0), \quad (8.28)$$

where

$$\mathcal{C}_{ij}^{(1)} = \frac{\lambda \lambda'}{4608\pi} (N+2) \frac{1}{\omega^4} \frac{T}{V} e^{4\omega X_0} \left(\delta_{ij} \vec{\Phi}^2 + 2\Phi_i \Phi_j \right), \quad (8.29)$$

$$\mathcal{C}_{ij}^{(2)} = -\frac{\delta_{ij}}{\pi^2} \lambda^2 \lambda' \left(\frac{N+2}{96} \right)^3 \frac{1}{\omega^6} \left(\frac{T}{V} \right)^2 e^{4\omega X_0} \left[1 - 2\pi \frac{N+8}{(N+2)^2} \frac{1}{\omega^2 T V} \right]. \quad (8.30)$$

Let us write $\varphi_i(X_0)$, Eq. (8.16), in the form

$$\varphi_i(X_0) = \sum_{l=0}^3 \varphi_i^{[l]}(X_0),$$

where $\varphi_i^{[l]}(X_0)$ includes l ζ 's. $\mathcal{C}_{ij}^{(1)}(X_0)$, Eq. (8.29), comes from $\langle \varphi_i^{[1]}(X_0) \varphi_j^{[1]}(X_0) \rangle_{\zeta}$ and $\mathcal{C}_{ij}^{(2)}(X_0)$ comes from $\langle \varphi_i^{[1]}(X_0) \varphi_j^{[3]}(X_0) \rangle_{\zeta}$ and $\langle \varphi_i^{[3]}(X_0) \varphi_j^{[1]}(X_0) \rangle_{\zeta}$. $\mathcal{C}_{ij}^{(2)}(X_0)$ consists of two terms. The term being proportional to $1/V^2$ comes from (the square of) (8.19) while the term being proportional to $1/V^3$ comes from (8.20).

Here we summarize the characteristic features of (8.28) - (8.30).

- Comparing (8.28) - (8.30) with (8.27) we see that (8.28) with (8.29) and (8.30) is reliable in the region,

$$e^{2\omega X_0} < O \left(\text{Min} \left(\frac{\omega^2}{\lambda' \vec{\Phi}^2}, \frac{\omega^4 V}{\lambda \lambda' T}, \frac{\omega^6 V^2}{\lambda \lambda'} \right) \right).$$

- In contrast to (8.27) and $\mathcal{C}_{ij}^{(2)}(X_0)$, $\mathcal{C}_{ij}^{(1)}(X_0)$ is not diagonal in N -dimensional vector space, which means that the correlation between different component of $\vec{\varphi}(X_0)$ exists. $\mathcal{C}_{ii}^{(1)}(X_0)$ (with no summation over i) is positive, while $\mathcal{C}_{ij(\neq i)}^{(1)}(X_0)$ can either be positive or negative depending on $\vec{\Phi}$.
- The $1/V^2$ term in $\mathcal{C}_{ij}^{(2)}(X_0)$ is negative and then enhances (8.27), while the $1/V^3$ term is positive.

3. Four-point ζ -correlation function

Setting $X_0 = Y_0 = U_0 = V_0$ in (8.23), we obtain

$$\begin{aligned} & \langle \varphi_i^{(0)}(X_0) \varphi_j^{(0)}(Y_0) \varphi_k^{(0)}(U_0) \varphi_l^{(0)}(V_0) \rangle_{\zeta c} \\ &= -\frac{N+8}{4\pi} \left(\frac{\lambda}{48} \right)^2 \frac{T}{\omega^6 V^3} [\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}] \mathcal{H}(\omega X_0), \end{aligned} \quad (8.31)$$

where

$$\mathcal{H}(\omega X_0) = \sinh(4\omega X_0) - 8 \sinh(2\omega X_0) + 12\omega X_0.$$

One can easily show that $\mathcal{H}(\omega X_0)$ ($X_0 > 0$) is positive definite, so that the four-point correlation (8.31) is of negative. For $\omega X_0 \ll 1$ and $\omega X_0 \gg 1$, we have

$$\mathcal{H}(\omega X_0) = \begin{cases} \frac{32}{5} (\omega X_0)^5 & (\omega X_0 \ll 1) \\ \frac{1}{2} e^{4\omega X_0} & (\omega X_0 \gg 1). \end{cases}$$

4. Features of the distribution function (8.4)

For the purpose of comparison, instead of (8.4), we take the quasi equilibrium distribution function,

$$N^{(\xi)}(X_0; p_0) \Big|_{\text{ref}} = \frac{1}{e^{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)|}/T} - 1}. \quad (8.32)$$

Same approximations as above yield,

$$\mathcal{I}_\xi^{(\beta)}(X_0) \simeq \frac{T^2}{12} + \frac{iT}{4\pi} \sqrt{|\mathcal{M}_\xi^2(X_0)|}, \quad (8.33)$$

$$\ddot{\vec{\varphi}}(X_0) = \left[\omega^2 - \frac{\lambda}{6} \vec{\varphi}^2(X_0) \right] \vec{\varphi}(X_0) + \vec{\zeta}(X_0) + H \vec{e}, \quad (8.34)$$

$$= -\frac{\partial \mathcal{V}}{\partial \vec{\varphi}(X_0)} + \vec{\zeta}(X_0), \quad (8.35)$$

$$\mathcal{V} \simeq \frac{\pi^2}{90} N T^4 - \frac{\omega^2}{2} \vec{\varphi}^2(X_0) + \frac{\lambda}{4!} (\vec{\varphi}^2(X_0))^2 - H({}^t \vec{e} \cdot \vec{\varphi}(X_0)), \quad (8.36)$$

$$\langle \zeta_i(X_0) \zeta_j(Y_0) \rangle_\zeta \simeq -\frac{\lambda}{24\pi} (N+2) \frac{\omega_0 T}{V} \delta_{ij} \delta(X_0 - Y_0), \quad (8.37)$$

$$\begin{aligned} \langle \zeta_i(X_0) \zeta_j(Y_0) \zeta_k(U_0) \zeta_l(V_0) \rangle_{\zeta c} &\simeq -\frac{\lambda^2}{144\pi} \frac{N+8}{V^3} \frac{T}{\omega_0} [\delta_{ik} \delta_{lj} + \delta_{kj} \delta_{li} + \delta_{ij} \delta_{lk}] \\ &\times \delta(X_0 - Y_0) \delta(X_0 - U_0) \delta(V_0 - Y_0). \end{aligned} \quad (8.38)$$

Let us summarize distinctive features of our results when compared to (8.33) - (8.38).

- Change in λ , $\lambda \rightarrow \lambda'$ [Eq. (8.11)], does not occur in (8.33) - (8.38). We emphasize again that $\lambda' < \lambda$.
- Equation (8.38) is two times (8.20), so that, besides the change $\lambda \rightarrow \lambda'$, (8.23), $1/V^3$ part of $\mathcal{C}_{ij}^{(2)}$ in (8.30), and (8.31) are one-half of the corresponding formulae with the reference distribution (8.32).
- We rewrite the potential $\mathcal{V}$, Eq. (8.14), in the form,

$$\mathcal{V} \simeq \frac{\pi^2}{90} N T^4 - \frac{\omega^2 + \Delta\omega^2}{2} \vec{\varphi}^2(X_0) + \frac{\lambda}{4!} (\vec{\varphi}^2(X_0))^2 - H({}^t \vec{e} \cdot \vec{\varphi}(X_0)). \quad (8.39)$$

Here

$$\begin{aligned} \omega^2 + \Delta\omega^2 &= \omega^2 + 2(N+8) \left(\frac{\lambda}{24\pi} \right)^2 \frac{T}{\omega} \vec{\varphi}^2(X_0) \\ &\simeq -\mu^2 - \frac{\lambda}{72} (N+2) (T + \Delta T)^2 \end{aligned}$$

with

$$\Delta T \simeq -\frac{\lambda}{8\pi^2} \frac{N+8}{N+2} \frac{\vec{\varphi}^2(X_0)}{\omega}. \quad (8.40)$$

Equation (8.39) is to be compared with (8.36). Comparing $\omega^2 + \Delta\omega^2$ with

$$\omega^2 = -\mu^2 - \frac{\lambda}{72} (N+2) T^2, \quad (8.41)$$

we see that, as $\vec{\varphi}(X_0)$ is “rolling down” the potential hill, the temperature “decreases,” and thus the system “cools.”

- We write the distribution function (8.4) in the form,

$$N^{(\xi)}(X_0; p_0) = \frac{1}{e^{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)| - e_\xi \lambda \vec{\varphi}^2(X_0)}/T} - 1} \simeq \frac{1}{e^{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)|}/T_{eff}(X_0)} - 1} \quad (e_\pi = 1/6, e_\sigma = 1/2), \quad (8.42)$$

where

$$\frac{T_{eff}(X_0)}{T} = \frac{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)|}}{\sqrt{p^2 - |\mathcal{M}_\xi^2(X_0)| - e_\xi \lambda \vec{\varphi}^2(X_0)}}. \quad (8.43)$$

Let us compare this with the reference distribution (8.32). We first note that the parts of the stable (oscillator) modes in (8.32),

$$|\mathcal{M}_\xi^2(X_0)| \leq p^2 \leq |\mathcal{M}_\xi^2(X_0)| + e_\xi \lambda \vec{\varphi}^2(X_0), \quad (8.44)$$

are the unstable modes in (8.42) and thus the “number” of unstable modes in the latter are larger than that in the former. For the modes, $|\mathcal{M}_\xi^2(X_0)| + e_\xi \lambda \vec{\varphi}^2(X_0) \geq p^2$, (8.43) shows that the effective temperature increases when compared to the reference distribution (8.32). This temperature increase is interpreted as follows. As seen above, while $\vec{\varphi}(X_0)$ becomes large, the potential $\mathcal{V}$ becomes small and the energy of the condensate fields $\vec{\varphi}(X_0)$ is released and converted to the kinetic energy of the quantum fields. On the other hand, for the unstable modes, $p^2 \leq |\mathcal{M}_\xi^2(X_0)|$, (8.43) shows that the effective temperature decreases when compared to the reference distribution (8.32). These unstable modes yield negative contribution to $Re \mathcal{I}_\xi^{(\beta)}$ in (8.6). All the effects mentioned above compete and $Re \mathcal{I}_\xi^{(\beta)}$ turns out to be smaller than the reference-distribution counterpart.

B. Nonuniform system

In this subsection, we briefly analyze the spatially nonuniform system under the similar situation as in the last subsection. We assume that the initial distribution function is given by (8.1):

$$N^{(\xi)}(X_0 = 0, \mathbf{X}; P) = \frac{1}{e^{|p_0|/T} - 1}. \quad (8.45)$$

The determining equation for $f^{(\xi)}(X; P) = \theta(p_0)N^{(\xi)}(X; P) - \theta(-p_0)(1 + N^{(\xi)}(X; -P))$, Eq. (6.53), may be solved as in (5.42):

$$N^{(\xi)}(X; P) \simeq \mathcal{F}(\underline{\mathbf{X}}; P^2 - \mathcal{M}_\xi^2(X), \underline{\mathbf{p}}), \quad (8.46)$$

where $\underline{\mathbf{X}}$ and $\underline{\mathbf{p}}$ are as in (5.43) with $T_{in} = 0$. Substitution of the initial data (8.45) into (8.46) shows that $\mathcal{F}$ is independent of $\underline{\mathbf{X}}$ and $\underline{\mathbf{p}}$ and $N^{(\xi)}(X; P)$ becomes (8.4):

$$N^{(\xi)}(X; P) = \frac{1}{e^{\sqrt{p^2 + M_\xi^2(X_0=0, \mathbf{X})}/T} - 1}.$$

At the initial time $X_0 = 0$, $|\vec{\varphi}|$ and $|\partial \vec{\varphi} / \partial X_\mu|$ are small, so that (cf. (8.10) and (8.9))

$$M_\xi^2(X_0 = 0, \mathbf{X}) \simeq -\omega^2 \quad (\xi = \pi, \sigma),$$

which is independent of $\mathbf{X}$. In place of (8.19) and (8.20), we have, in respective order,

$$\langle \zeta_i(X) \zeta_j(Y) \rangle_\zeta \simeq -\frac{\lambda}{24\pi} (N+2) T \omega \delta_{ij} \delta^4(X-Y), \quad (8.47)$$

$$\begin{aligned} \langle \zeta_i(X) \zeta_j(Y) \zeta_k(U) \zeta_l(V) \rangle_{\zeta c} &\simeq -\frac{\lambda^2}{288\pi} (N+8) \frac{T}{\omega} [\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}] \\ &\times \delta^4(X-Y) \delta^4(X-U) \delta^4(V-Y). \end{aligned} \quad (8.48)$$

In this subsection, we are only concerned with a zeroth-order solution for $\vec{\varphi}(X)$ (cf. (8.15)),

$$\vec{\varphi}^{(0)}(X) = \vec{\varphi}_{pure}^{(0)}(X) + \int \frac{d^3p}{(2\pi)^3} \frac{e^{i\mathbf{p}\cdot\mathbf{X}}}{\omega'} \int_0^{X_0} dX'_0 \vec{\zeta}(X'_0; \mathbf{p}) \sinh(\omega'(X_0 - X'_0)), \quad (8.49)$$

where

$$\omega' = \sqrt{\omega^2 - p^2}, \quad (8.50)$$

$$\vec{\zeta}(X'_0; \mathbf{p}) = \int d^3X e^{-i\mathbf{p}\cdot\mathbf{X}} \vec{\zeta}(X'_0, \mathbf{X}). \quad (8.51)$$

For, $\omega^2 - p^2 < 0$, ω' should be understood to be $\omega' = i\sqrt{p^2 - \omega^2}$. In (8.49), $\vec{\varphi}_{pure}^{(0)}(X)$ is the solution to the equation

$$(\partial_X^2 - \omega^2) \vec{\varphi}_{pure}^{(0)}(X) = H\vec{e},$$

under a given initial data.

1. Two-point ζ -correlation function

Using (8.47), one can readily compute the two-point ζ -correlation function:

$$\begin{aligned} \langle \varphi_i^{(0)}(X) \varphi_j^{(0)}(Y) \rangle_{\zeta c} &= -\frac{\lambda}{96\pi} \delta_{ij} (N+2) \omega T \int \frac{d^3p}{(2\pi)^3} e^{i\mathbf{p}\cdot(\mathbf{X}-\mathbf{Y})} \frac{1}{\omega^2 - p^2} \\ &\quad \times \left[\frac{\sinh(\sqrt{\omega^2 - p^2}(X_0 + Y_0)) - \sinh(\sqrt{\omega^2 - p^2}(X_0 - Y_0))}{\sqrt{\omega^2 - p^2}} \right. \\ &\quad \left. - 2Y_0 \cosh(\sqrt{\omega^2 - p^2}(X_0 - Y_0)) \right] \\ &= -\frac{\lambda}{192\pi^3} \delta_{ij} (N+2) \frac{\omega T}{|\mathbf{X} - \mathbf{Y}|} \int_0^\infty dp p \sin(p|\mathbf{X} - \mathbf{Y}|) \frac{1}{\omega^2 - p^2} \\ &\quad \times \left[\theta(\omega^2 - p^2) \left\{ \frac{\sinh(\sqrt{\omega^2 - p^2}(X_0 + Y_0)) - \sinh(\sqrt{\omega^2 - p^2}(X_0 - Y_0))}{\sqrt{\omega^2 - p^2}} \right. \right. \\ &\quad \left. \left. - 2Y_0 \cosh(\sqrt{\omega^2 - p^2}(X_0 - Y_0)) \right\} \right. \\ &\quad \left. + \theta(p^2 - \omega^2) \left\{ \frac{\sin(\sqrt{p^2 - \omega^2}(X_0 + Y_0)) - \sin(\sqrt{p^2 - \omega^2}(X_0 - Y_0))}{\sqrt{p^2 - \omega^2}} \right. \right. \\ &\quad \left. \left. - 2Y_0 \cos(\sqrt{p^2 - \omega^2}(X_0 - Y_0)) \right\} \right], \end{aligned} \quad (8.52)$$

where $X_0 \geq Y_0$ has been assumed. Manipulation of (8.52) in Appendix E yields

$$\begin{aligned} \langle \varphi_i^{(0)}(X) \varphi_j^{(0)}(Y) \rangle_{\zeta c} &= -\frac{\lambda}{192\pi^3} (N+2) \delta_{ij} \frac{T}{|\mathbf{X} - \mathbf{Y}|} \\ &\quad \times [\theta(X_0 + Y_0 - |\mathbf{X} - \mathbf{Y}|) \theta(|\mathbf{X} - \mathbf{Y}| - (X_0 - Y_0)) \\ &\quad \times \left\{ \frac{\pi\omega}{2} (X_0 + Y_0) \cos(\omega|\mathbf{X} - \mathbf{Y}|) - \sin(\omega|\mathbf{X} - \mathbf{Y}|) \right. \\ &\quad \left. + \int_0^1 d\xi \frac{\xi}{(1 - \xi^2)^{3/2}} \left[\sin(\omega|\mathbf{X} - \mathbf{Y}|\xi) \cosh(\sqrt{1 - \xi^2}\omega(X_0 + Y_0)) \right. \right. \\ &\quad \left. \left. - \sin(\omega|\mathbf{X} - \mathbf{Y}|) \right] \right\} \\ &\quad + \theta(X_0 - Y_0 - |\mathbf{X} - \mathbf{Y}|) \\ &\quad \times \left\{ \int_0^1 d\xi \frac{\xi \sin(\omega|\mathbf{X} - \mathbf{Y}|\xi)}{(1 - \xi^2)^{3/2}} \sum_{\tau=\pm} \tau \cosh(\sqrt{1 - \xi^2}\omega(X_0 + \tau Y_0)) \right. \\ &\quad \left. - 2\omega Y_0 \int_0^1 d\xi \frac{\xi \sin(\omega|\mathbf{X} - \mathbf{Y}|\xi)}{1 - \xi^2} \sinh(\sqrt{1 - \xi^2}\omega(X_0 - Y_0)) \right\}. \end{aligned} \quad (8.53)$$

First of all, we note that $\langle \varphi_i^{(0)}(X) \varphi_j^{(0)}(Y) \rangle_{\zeta c}$ vanishes for $|\mathbf{X} - \mathbf{Y}| > X_0 + Y_0$, which means that the correlation does not spread with super-light velocity. Several limiting cases are analyzed in Appendix E.

1) $X_0 - Y_0 < |\mathbf{X} - \mathbf{Y}| < X_0 + Y_0$; $\omega|\mathbf{X} - \mathbf{Y}| < \omega(X_0 + Y_0) \ll 1$:

$$\begin{aligned} \langle \varphi_i(X) \varphi_j(Y) \rangle_{\zeta c} \simeq & -\frac{\lambda}{384\pi^2} (N+2) \delta_{ij} \frac{\omega T}{|\mathbf{X} - \mathbf{Y}|} \\ & \times \left[X_0 + Y_0 - |\mathbf{X} - \mathbf{Y}| \left\{ 1 - \frac{\omega^2(X_0 + Y_0)^2}{4} \right\} \right]. \end{aligned} \quad (8.54)$$

This shows that the correlation is negative. For fixed $|\mathbf{X}|$ and $|\mathbf{Y}|$, the correlation is maximum at the smallest $|\mathbf{X} - \mathbf{Y}|$, $|\mathbf{X} - \mathbf{Y}| = \text{Max}(X_0 - Y_0, ||\mathbf{X}| - |\mathbf{Y}||)$ and is minimum at the largest $|\mathbf{X} - \mathbf{Y}|$, $|\mathbf{X} - \mathbf{Y}| = \text{Min}(X_0 + Y_0, |\mathbf{X}| + |\mathbf{Y}|)$. We are dealing with the zeroth-order solution to the evolution equation (the counterpart to (8.12)), i.e., the nonlinear term, $-\lambda' \vec{\varphi}^2 \vec{\varphi}/6$, in the evolution equation has been ignored. The condition that the ignored term is small restrict the domain, where (8.54) is valid, to

$$\lambda \lambda' \frac{X_0 + Y_0 - |\mathbf{X} - \mathbf{Y}|}{|\mathbf{X} - \mathbf{Y}|} < O\left(\frac{\omega}{T}\right).$$

2) $1 \ll \omega(X_0 + Y_0)$ and $\omega|\mathbf{X} - \mathbf{Y}| \ll \sqrt{\omega(X_0 + Y_0)}$:

$$\begin{aligned} \langle \varphi_i(X) \varphi_j(Y) \rangle_{\zeta c} \simeq & -\frac{\sqrt{2\pi}}{768\pi^3} \lambda(N+2) \delta_{ij} \omega T \\ & \times \left[\frac{1}{(\omega(X_0 + Y_0))^{3/2}} \exp\left(\omega \left\{ X_0 + Y_0 - \frac{|\mathbf{X} - \mathbf{Y}|^2}{2(X_0 + Y_0)} \right\}\right) \right. \\ & \times \left(1 + O\left(\frac{1}{\omega(X_0 + Y_0)}\right) \right) \\ & + \sqrt{2\pi} \theta(X_0 + Y_0 - |\mathbf{X} - \mathbf{Y}|) \theta(|\mathbf{X} - \mathbf{Y}| - (X_0 - Y_0)) \\ & \left. \times \frac{X_0 + Y_0}{|\mathbf{X} - \mathbf{Y}|} \cos(\omega|\mathbf{X} - \mathbf{Y}|) \right], \end{aligned} \quad (8.55)$$

which is valid for

$$\lambda \lambda' e^{\omega(X_0 + Y_0)} < O\left(\frac{\omega}{T} \{\omega(X_0 + Y_0)\}^{3/2}\right) \quad \text{and} \quad \lambda \lambda' \frac{X_0 + Y_0}{|\mathbf{X} - \mathbf{Y}|} < O\left(\frac{\omega}{T}\right).$$

Equation (8.55) indicates that, for large $X_0 + Y_0$ with $\omega|\mathbf{X} - \mathbf{Y}| \ll \sqrt{\omega(X_0 + Y_0)}$, the (negative) correlation spreads over the distance,

$$|\mathbf{X} - \mathbf{Y}| \sim \sqrt{\frac{2(X_0 + Y_0)}{\omega}},$$

and asymptotically spreads over whole space region. It should be emphasized, however, that, for too large $X_0 + Y_0$, our whole approximations in this section cease to hold.

3) $|\mathbf{X} - \mathbf{Y}| < X_0 - Y_0$; $\omega(X_0 + Y_0) \ll 1$:

$$\langle \varphi_i(X) \varphi_j(Y) \rangle_{\zeta c} \simeq -\frac{\lambda}{384\pi^2} (N+2) \delta_{ij} \omega T (\omega Y_0)^2.$$

2. Four-point ζ -correlation function

Using (8.49) and (8.48), we obtain

$$\begin{aligned} & \langle \varphi_{i_1}^{(0)}(X_1) \varphi_{i_2}^{(0)}(X_2) \varphi_{i_3}^{(0)}(X_3) \varphi_{i_4}^{(0)}(X_4) \rangle_{\zeta c} \\ & \simeq -\frac{\lambda^2}{288\pi} (N+8) \frac{T}{\omega} [\delta_{i_1 i_2} \delta_{i_3 i_4} + \delta_{i_1 i_3} \delta_{i_2 i_4} + \delta_{i_1 i_4} \delta_{i_2 i_3}] \\ & \times \int_0^{X_0} dX'_0 \int d\mathbf{X}' \prod_{j=1}^4 \int \frac{d^3 p_j}{(2\pi)^3} \frac{e^{i\mathbf{p}_j \cdot (\mathbf{x}_j - \mathbf{x}')}}{\sqrt{\omega^2 - p_j^2}} \sinh\left(\sqrt{\omega^2 - p_j^2} (X_{j0} - X'_0)\right), \end{aligned} \quad (8.56)$$

where $X_{10}, X_{20}, X_{30} \geq X_{40}$ has been assumed. For equal times, $X_{10} = X_{20} = X_{30} = X_{40} (\equiv X_0)$, computation of (8.56) in the limit, $\omega X_0 \gg 1$ and $|\mathbf{X}_i - \mathbf{X}_j| \ll X_0$ ($i, j = 1, \dots, 4$) is relatively simple. Similar calculation as in Appendix E yields

$$\begin{aligned} & \langle \varphi_{i_1}^{(0)}(X_0, \mathbf{X}_1) \varphi_{i_2}^{(0)}(X_0, \mathbf{X}_2) \varphi_{i_3}^{(0)}(X_0, \mathbf{X}_3) \varphi_{i_4}^{(0)}(X_0, \mathbf{X}_4) \rangle_{\zeta c} \\ & \simeq -\frac{\lambda^2}{288\pi} (N+8) [\delta_{i_1 i_2} \delta_{i_3 i_4} + \delta_{i_1 i_3} \delta_{i_2 i_4} + \delta_{i_1 i_4} \delta_{i_2 i_3}] \\ & \times \omega^3 T \frac{e^{4\omega X_0}}{(8\pi\omega X_0)^{9/2}} \exp\left(-\frac{\omega}{8X_0} \sum_{i < j} (\mathbf{X}_i - \mathbf{X}_j)^2\right), \end{aligned} \quad (8.57)$$

which is valid for

$$(\lambda\lambda')^2 e^{4\omega X_0} < O\left(\frac{\omega}{T} (\omega X_0)^{9/2}\right).$$

It should be emphasized that, if we use the reference distribution (8.32), we obtain two times (8.57).

IX. CONCLUDING REMARKS

In this paper, two related subjects are dealt with:

- A) Deduction from first principles of the perturbative framework for dealing with out-of-equilibrium relativistic complex-scalar-field systems.
- B) Deduction from first principles of the perturbative-loop-expansion framework for studying an $O(N)$ linear σ model.

We have assumed the existence of two different spacetime scales, the microscopic and macroscopic. The first small scale characterizes the microscopic correlations and the second large scale is inherent in the relaxation phenomena. Besides this setup, in principle, we have not made use of any further approximations. Let us summarize what has been added and what has been clarified in this paper. Some additional comments are also given.

3. On the issue A)

- (1) The assumption that is necessary for the propagator to take the standard form has been singled out. In most available work, among $2n$ -point initial correlation functions $\mathcal{W}_{2n}$ ($n \geq 1$), Eq. (3.13), $\mathcal{W}_{2n}$ ($n \geq 2$) are dropped and only $\mathcal{W}_2$ is kept into the propagator. This is so done as an approximation or by assuming some specific form for the density matrix. In this paper, we have seen that, to what extent, $\mathcal{W}_{2n}$ ($n \geq 2$) may be discarded. Whenever necessary, one can incorporate them into the perturbative framework (cf. [2]).
- (2) We have emphasized that the perturbative framework is effective only when the inverse size of a spacetime cell is much smaller than the infrared scale in the microscopic sector of the theory. In the present complex-scalar-field theory, this condition is $(L^\mu)^{-1} \lesssim \sqrt{\lambda} \mathcal{P}$ ($\mu = 0, 1, 2, 3$) with λ the coupling constant and $\mathcal{P}$ the typical parameter(s) (of mass dimension) of the system.
- (3) We have proposed mutually equivalent two perturbative schemes. Both of them lead to a generalized relativistic kinetic or Boltzmann equation. Derivation of it has proceeded in a physically transparent manner. In the physical- f scheme, the equation is derived through the process of redefinition of the number-density or relativistic Wigner function, from the initial number-density function to the physically sensible one. Traditional derivation of the equation proceeds [2,13] in a rather abstract way, using the Dyson equation. (For the case of nonrelativistic many-body theory, see, e.g., [39].) As seen in Section V, the Dyson equation is the equation that serves as determining the full propagator in terms of the self-energy part. In other word, the Dyson equation simply serves as an efficient way of resumming the self-energy part. Then, by referring to the full propagator thus obtained, one introduces [2,13,39] a redefined distribution function. It is this function that subjects to the relativistic Boltzmann equation.

- (4) In massless theories, divergence due to infrared and/or mass (or collinear) singularities appear in some amplitudes. This occurs as a result of an interplay of *bare* massless propagators. By performing resummations of the self-energy part for such propagators (cf. (5.11)), one can get rid of this divergence disaster (cf., e.g., [2,40]. For a different calculational scheme, in which no divergence emerges, see [25].

4. On the issue B)

- (1) Through introducing effective masses, which depend on the macroscopic spacetime coordinates, we have derived self-consistent gap equation for them. Together with the generalized relativistic Boltzmann equation (cf. item (3) above) and the equation of motion for the condensate fields, this constitutes the system of coupled equations, which describes how the phase transition proceeds. It should be emphasized, however, that, as for other work of this sort, the present scheme applies only for the systems away from the critical region.
- (2) The negative curvature region of the “potential” has been dealt with by introducing random-force fields. To leading one-loop order, this leads to negative correlation between the condensate fields.
- (3) As in the issue A) above, within the gradient approximation, the propagator consists of two parts, $\hat{\Delta} = \hat{\Delta}^{(0)} + \hat{\Delta}^{(1)}$, where $\hat{\Delta}^{(0)}$ is the dominant part and the “correction part” $\hat{\Delta}^{(1)}$ contains a derivative with respect to the macroscopic spacetime coordinates. In a $(\vec{\pi}, \sigma)$ -space, $\hat{\Delta}^{(0)}$ is diagonal, while $\hat{\Delta}^{(1)}$ is not. This means that, in the level of $\hat{\Delta}^{(1)}$, there occurs mixing between $\vec{\pi}$ and σ sectors. Then, when one takes the part $\hat{\Delta}^{(1)}$ into account, rediagonalization of the propagator in the $(\vec{\pi}, \sigma)$ -space is necessary, which leads to “physical” $\vec{\pi}$ and σ .

Finally, we mention two related works [4,15,30]. In the theories developed there, a “generalized mass (function)” is introduced, which is 2×2 matrix in the “complex-time plane” and matrix elements are complex functions of spacetime. Due to the last fact, the theories provide for a treatment of dissipative effects. The “generalized mass (function)” is determined self consistently through renormalization [4,15] or through some optimization procedure [30]. Through this process, the Boltzmann equation emerges. As applied to the complex-scalar field theory, the part of the hat-Lagrangian density that include the “generalized mass (function)” corresponds to (cf. (4.39) - (4.41) and (5.46))

$$- \sum_{j=1}^2 (-)^{j-1} \phi_j^*(x) M^2(x) \phi_j(x) - \hat{\mathcal{L}}_c$$

(with $\hat{\mathcal{L}}_c$ as in (4.41)) in this paper.

ACKNOWLEDGMENTS

This work was supported in part by the Grant-in-Aide for Scientific Research ((A)(1) (No. 08304024)) of the Ministry of Education, Science and Culture of Japan.

APPENDIX A: MASS-DERIVATIVE FORMULA

Here an argument for necessity of adopting the $|p_0|$ prescription is given. We add $\mathcal{L}_m = -\delta m^2 \phi^\dagger \phi$ to the Lagrangian density (3.3). The effect of $\mathcal{L}_m$ on the propagator $\hat{\Delta}(X; P)$ should be

$$\hat{\Delta}(X; P) \rightarrow \hat{\Delta}(X; P) \Big|_{m^2 \rightarrow m^2 + \delta m^2}. \quad (\text{A1})$$

Let us evaluate the $O(\delta m^2)$ correction, $\delta \hat{\Delta}$, to $\hat{\Delta}$ due to the perturbation $\mathcal{L}_m$:

$$\begin{aligned} \delta \hat{\Delta}(x, y) &= \delta m^2 \int d^4 u \hat{\Delta}(x, u) \hat{\tau}_3 \hat{\Delta}(u, y) \\ &\simeq \delta m^2 \int d^4 u \left[\hat{\Delta}(X, x - u) + \frac{u - y}{2} \cdot \partial_X \hat{\Delta}(X, x - u) \right] \\ &\quad \times \hat{\tau}_3 \left[\hat{\Delta}(X, u - y) + \frac{u - x}{2} \cdot \partial_X \hat{\Delta}(X, u - y) \right], \end{aligned}$$

where $X = (x + y)/2$. Taking the Fourier transform with respect to $x - y$ and using $\hat{A}_+ \hat{\tau}_3 \hat{A}_+ = 0$ and (4.33), we obtain

$$\delta \hat{\Delta}(X; P) \simeq \delta m^2 \hat{\Delta}(X; P) \hat{\tau}_3 \hat{\Delta}(X; P). \quad (\text{A2})$$

It is to be noted that (A2) is valid to the gradient approximation, i.e., the term with first derivative (with respect to X) is absent in (A2).

Substituting (4.28), we finally obtain

$$\delta \hat{\Delta}(X; P) \simeq \delta m^2 \left[\begin{pmatrix} \Delta_R^2 & 0 \\ \Delta_R^2 - \Delta_A^2 & -\Delta_A^2 \end{pmatrix} + f(\Delta_R^2 - \Delta_A^2) \hat{A}_+ \right]. \quad (\text{A3})$$

If we adopt the $|p_0|$ prescription, $f = f(X; P)$ is independent of m^2 (cf. (4.30)), so that $\delta \hat{\Delta}$ in (A3) may be written as

$$\delta \hat{\Delta}(X; P) = \delta m^2 \frac{\partial \hat{\Delta}(X; P)}{\partial m^2}, \quad (\text{A4})$$

which is in accord with (A1). On the other hand, if we take original f (Eq. (4.30)), $f = \theta(p_0)N_+(X; E_p, \hat{\mathbf{p}}) - \theta(-p_0)[1 + N_+(X; E_p, -\hat{\mathbf{p}})]$, it depends on m^2 through E_p and $\delta \hat{\Delta}$ in (A3) cannot be written in the “mass-derivative form” (A4). We have confirmed the consistency between the mass-derivative formula and the $|p_0|$ prescription at least up to the terms with second-order derivative with respect to X .

APPENDIX B: PROPERTIES OF THE SELF-ENERGY PART

B.1. Proof of $\Sigma_{11} + \Sigma_{12} + \Sigma_{21} + \Sigma_{22} = 0$

Let $\Sigma_l(x, y; z_1, \dots, z_n)$ be a contribution from a loop diagram to the self-energy part in configuration space, where x and y are the space-time coordinates of external vertices and $z_1, \dots, z_n$ are those of internal vertices. Here an external vertex means a vertex to which an external leg is attached. The times $x_0, y_0, z_{10}, \dots, z_{n0}$ are on the closed time path $C_1 \oplus C_2$.

Let us first show that, when one of the internal vertices, say z_{i0} , is the largest time,

$$z_{i0} > x_0, y_0, \text{ other } z_0\text{'s}, \quad (\text{B1})$$

Σ_l vanishes [41]. This can be proved as follows. From (3.4) and (3.5), we see that a vertex factor with $z_{i0} \in C_1$ possesses an opposite sign relative to the corresponding vertex factor with $z_{i0} \in C_2$:

$$(iV_4)_1 = -(iV_4)_2, \quad (iV_2)_1 = -(iV_2)_2, \quad (iV'_4)_1 = -(iV'_4)_2. \quad (\text{B2})$$

On the other hand, as can be easily be seen from (2.3), all the propagators in Σ_l are the same for $z_{i0} \in C_1$ and $z_{i0} \in C_2$.

Let $(\Sigma_l)_{ij}$ be $\Sigma_l(x, y; z_1, \dots, z_n)$ with $x_0 \in C_i$ and $y_0 \in C_j$. Similar observation as above shows that

$$\begin{aligned} (\Sigma_l)_{1i} + (\Sigma_l)_{2i} &= 0 \quad (i = 1, 2) \quad \text{for } x_0 > y_0, z_1, \dots, z_n, \\ (\Sigma_l)_{i1} + (\Sigma_l)_{i2} &= 0 \quad (i = 1, 2) \quad \text{for } y_0 > x_0, z_1, \dots, z_n. \end{aligned}$$

Then, we obtain

$$\sum_{i,j=1}^2 (\Sigma_l(x, y; z_1, \dots, z_n))_{ij} = 0. \quad (\text{B3})$$

There are contributions to the self-energy part, $\Sigma_l(x; z_1, \dots, z_n)$ which include only one external vertex. In such cases, $(\Sigma_l)_{12} = (\Sigma_l)_{21} = 0$. Similar argument as above shows that

$$\sum_{i=1}^2 (\Sigma_l(x; z_1, \dots, z_n))_{ii} = 0. \quad (\text{B4})$$

There is an exception to this argument, i.e., a contribution from the lowest-order tadpole diagram, $(\Sigma_l(x))_{ij}$. Explicit computation shows that $\sum_{i=1}^2 (\Sigma_l(x))_{ii} = 0$ for the complex-scalar field theory analyzed in Sections III-V and for the $O(N)$ linear σ model in the symmetric phase (cf. Section VIIA). This is not the case for the broken-symmetric phase (cf. Section VIIB). As a matter of fact, in the region, where the curvature of the “potential” is negative, imaginary part develops in Σ_{11} and Σ_{22} : $Im\Sigma_{11} = Im\Sigma_{22}$. Then $\sum_{i=1}^2 (\Sigma_l(x))_{ii} \neq 0$ but $\sum_{i=1}^2 Re(\Sigma_l(x))_{ii} = 0$. Thus, in studying this region, one should take this fact into account.

Inclusion of the vertices $i\hat{\mathcal{V}}_c(x, y)$ (Eq. (4.42)) that comes from the counter Lagrangian does not invalidate the above argument. The vertex $i\hat{\mathcal{V}}_c(x, y)$ yields a contribution to the self-energy part $\hat{\Sigma}_c(x, y) = -\hat{\mathcal{V}}_c(x, y)$ (see (5.1)). From (4.42) with (3.16), it is obvious that

$$\sum_{i,j=1}^2 (\Sigma_c(x, y))_{ij} = 0.$$

$i\hat{\mathcal{V}}_c$ also appears as an internal vertex of Σ_l 's. Using the form (4.42), we can readily see that the similar argument as above after (B1) holds, so that (B3) and (B4) hold.

This completes the proof.

B.2. Proof of $\Sigma_A(X; P) = [\Sigma_R(X; P)]^*$

From the definition of G 's, Eq. (2.3), it can easily be shown that

$$\begin{aligned} iG_{11}(x, y) &= [iG_{22}(y, x)]^*, \\ iG_{12(21)}(x, y) &= [iG_{12(21)}(y, x)]^*. \end{aligned} \quad (B5)$$

This shows that, by taking a complex conjugate of $iG_{jl}(x, y)$, the index j [l] at the end point x [y] changes; $j = 1$ (respect. 2) $\rightarrow 2$ (respect. 1) [$l = 1$ (respect. 2) $\rightarrow 2$ (respect. 1)]. Let us define

$$\Sigma_l(x, y) \equiv \int_C \left(\prod_{j=1}^2 d^4 z_j \right) \Sigma_l(x, y; z_1, \dots, z_n),$$

where $\Sigma_l(x, y; z_1, \dots, z_n)$ is as in the previous subsection. From (3.4) and (3.5), we obtain

$$i(V_4)_2 = [i(V_4)_1]^*, \quad i(V_2)_2 = [i(V_2)_1]^*, \quad i(V'_4)_2 = [i(V'_4)_1]^*. \quad (B6)$$

Thus, again by taking a complex conjugate of $i(V_4)_j$, etc., the index j changes; $j = 1$ (respect 2) $\rightarrow 2$ (respect 1). For the $O(N)$ linear σ model, similar relations hold. As to the vertex $i\hat{\mathcal{V}}_c(x, y)$, Eq. (4.42), we have $i\hat{\mathcal{V}}_c(x, y) = \{i\hat{\mathcal{V}}_c(y, x)\}^*$, so that

$$\begin{aligned} i(\mathcal{V}_c(x, y))_{11} &= \{i(\mathcal{V}_c(y, x))_{22}\}^*, \\ i(\mathcal{V}_c(x, y))_{12(21)} &= \{i(\mathcal{V}_c(y, x))_{12(21)}\}^*. \end{aligned} \quad (B7)$$

Incidentally, this shows that $(\Sigma_c(x, y))_{ij}$ ($i, j = 1, 2$) in (5.1) satisfies (B7) with $(\mathcal{V}_c)_{ij} \rightarrow (\Sigma_c)_{ij}$.

Using (B5) - (B7), we can show that

$$\begin{aligned} -i(\Sigma_l(x, y))_{11} &= [-i(\Sigma_l(y, x))_{22}]^*, \\ -i(\Sigma_l(x, y))_{12(21)} &= [-i(\Sigma_l(y, x))_{12(21)}]^*. \end{aligned} \quad (B8)$$

Fourier transformation on $x - y$ yields

$$\begin{aligned} (\Sigma_l(X; P))_{11} &= -(\Sigma_l(X; P))_{22}^*, \quad \Sigma_{11}(X; P) = -\Sigma_{22}^*(X; P), \\ (\Sigma_l(X; P))_{12}, \quad (\Sigma_l(X; P))_{21} &: \text{pure imaginary.} \end{aligned} \quad (B9)$$

Using this in (5.4), we obtain $\Sigma_A(X; P) = [\Sigma_R(X; P)]^*$.

APPENDIX C: USEFUL FORMULA

We deal with

$$\int d^4(x-y)e^{iP \cdot (x-y)} \int d^4u d^4v A(x,u)B(u,v)C(v,y). \quad (C1)$$

We wish to express this quantity using the macroscopic coordinate $X = (x+y)/2$ up to the first order with respect to the X dependence. To this end, we expand A as

$$\begin{aligned} A(x,u) &= A((x+u)/2; x-u) \\ &\simeq A(X; x-u) + \frac{u-y}{2} \cdot \partial_X A(X; x-u). \end{aligned}$$

B and C may be expanded similarly. Substituting these into (C1) and Fourier transforming with respect to the relative coordinates, we obtain

$$\text{Eq. (C1)} \simeq ABC + \frac{i}{2} \left[\frac{\partial A}{\partial P^\mu} \frac{\partial B}{\partial X_\mu} C - A \frac{\partial B}{\partial X_\mu} \frac{\partial C}{\partial P^\mu} - \frac{\partial A}{\partial X_\mu} \frac{\partial BC}{\partial P^\mu} + \frac{\partial AB}{\partial P^\mu} \frac{\partial C}{\partial X_\mu} \right], \quad (C2)$$

where $A = A(X; P)$, etc.

The second form we deal with is

$$\int d^4(x-y)e^{iP \cdot (x-y)} \int d^4u d^4v f(x)A(x,u)B(u,v)C(v,y)g(y), \quad (C3)$$

where $f(x)$ and $g(y)$ depend weakly on x . In a similar manner as above, we obtain

$$\begin{aligned} \text{Eq. (C3)} \simeq fgABC &+ \frac{i}{2} \left[fg \left\{ \frac{\partial A}{\partial P^\mu} \frac{\partial B}{\partial X_\mu} C - A \frac{\partial B}{\partial X_\mu} \frac{\partial C}{\partial P^\mu} - \frac{\partial A}{\partial X_\mu} \frac{\partial BC}{\partial P^\mu} + \frac{\partial AB}{\partial P^\mu} \frac{\partial C}{\partial X_\mu} \right\} \right. \\ &\left. + \left(f \overset{\leftrightarrow}{\partial}_{X_\mu} g \right) \frac{\partial ABC}{\partial P^\mu} \right], \end{aligned} \quad (C4)$$

where $f = f(X)$ and $g = g(X)$.

APPENDIX D: RENORMALIZABILITY OF OF THE THEORY BASED ON (5.46)

Let us call the perturbation theory based on (3.3) [(5.46)] the theory I [II]. We first observe that, in theory II, if we include the term $\chi(x)\phi^\dagger\phi \equiv (M^2(x) - m^2)\phi^\dagger\phi$ in $\mathcal{L}_{rc}$ into $\mathcal{L}_0$, theory II reduces to theory I, which is UV-renormalizable. As stated in Section VE, the difference between theory I and theory II lies in the fact that, in theory II, $\chi(x)\phi^\dagger\phi$ is regarded as a perturbative term.

Let

$$\Gamma(m^2, X) = \sum_{l=l_0}^n \Gamma^{(l)}(m^2, X) \quad (D1)$$

be a renormalized and then finite Green function evaluated in theory I. In (D1), X is the center-of-mass coordinates of the “reaction” described by Γ and irrelevant arguments have been dropped. If we are making a traditional perturbative expansion, $\Gamma^{(l)}$ stands for the $O(\lambda^l)$ contribution and, if we are making a loop or δ expansion, $\Gamma^{(l)}$ stands for the l -loop contribution. The X dependence of $\Gamma^{(l)}(m^2, X)$ has come from various sources. Among those, the relevant one to the following argument is the distribution function (in theory I), which we write $f_I(X; P)$. In fact, $\Gamma^{(l)}(m^2, X)$ is written as a sum of integrals over momenta P ’s, whose integrands contain $f_I(X; P)$, $\partial f_I(X; P)/\partial X_\mu$, etc. In order to visualize this dependence, we write (D1) as

$$\Gamma(m^2, \{f_I(X)\}) = \sum_{l=l_0}^n \Gamma^{(l)}(m^2, \{f_I(X)\}) \quad (D2)$$

Substituting

$$m^2 = M^2(X) - \chi(X),$$

into $\Gamma^{(l)}(m, \{f_I(X)\})$, we obtain

$$\Gamma^{(l)}(M^2(X) - \chi(X), \{f_I(X)\}). \quad (\text{D3})$$

As stated in Section VE, in theory II, $\chi(x)$ is one-order higher than $M^2(x)$. More precisely, in traditional perturbative expansion, $\chi(x) = O(\lambda)$ and, in loop expansion, $\chi(x) = O(\delta)$. Then, we write (D3) as

$$\text{Eq. (D3)} = \Gamma^{(l)}(M^2(X), \{f_I(X)\}) + \sum_{l'=1}^{\infty} (-\chi(x))^{l'} \frac{\partial^{l'} \Gamma^{(l)}(M^2(X), \{f_I(X)\})}{\partial (M^2(X))^{l'}}. \quad (\text{D4})$$

Note that $(-\chi)^{l'} \partial^{l'} / \partial (M^2)^{l'}$ is of $(l + l')$ th order.

The distribution functions in theory I and in theory II are different. The UV-divergence issue is foreign to them. We can rewrite $\Gamma^{(l)}(M^2(X), \{f_I(X)\})$ in (D4), with obvious notation, as

$$\begin{aligned} \Gamma^{(l)}(M^2(X), \{f_I(X)\}) &= \Gamma^{(l)}(M^2(X), \{f_{II}(X)\}) \\ &+ \sum_{l'=1}^{\infty} (\{f_I(X)\} - \{f_{II}(X)\})^{l'} \frac{\delta^{l'} \Gamma^{(l)}(M^2(X), \{f_{II}(X)\})}{\delta (\{f_{II}(X)\})^{l'}}. \end{aligned} \quad (\text{D5})$$

Since, the difference $\chi(x) = M^2(x) - m^2$ is $O(\lambda)$ [$O(\delta)$] for a perturbative [loop] expansion, it is obvious that $\{f_I(X)\} - \{f_{II}(X)\} = O(\lambda)$ [$O(\delta)$]. Substituting (D3) - (D5) into (D2), we obtain a new series. Truncating this series at the n th order, we obtain a perturbation series in theory II. It is obvious from the above argument that this series is free from UV divergences.

APPENDIX E: DERIVATIONS OF (8.53) - (8.55)

Here we compute

$$\begin{aligned} \mathcal{F}(X_0, Y_0; r) &\equiv - \int_0^\infty dp \frac{p \sin(pr)}{p^2 - \omega^2} \left[\frac{\sin(\sqrt{p^2 - \omega^2}(X_0 + Y_0)) - \sin(\sqrt{p^2 - \omega^2}(X_0 - Y_0))}{\sqrt{p^2 - \omega^2}} \right. \\ &\quad \left. - 2Y_0 \cos(\sqrt{p^2 - \omega^2}(X_0 - Y_0)) \right] \end{aligned} \quad (\text{E1})$$

$$\begin{aligned} &= \frac{1}{4} \int_0^\infty dp \left[\frac{p}{(p^2 - \omega^2)^{3/2}} \sum_{\rho, \sigma, \tau = \pm} \rho \sigma \tau \exp\left(i \left[\rho pr + \sigma \sqrt{p^2 - \omega^2}(X_0 + \tau Y_0) \right] \right) \right. \\ &\quad \left. - 2iY_0 \frac{p}{p^2 - \omega^2} \sum_{\rho, \sigma = \pm} \rho \exp\left(i \left[\rho pr + \sigma \sqrt{p^2 - \omega^2}(X_0 - Y_0) \right] \right) \right]. \end{aligned} \quad (\text{E2})$$

In the above equations, $X_0 \geq Y_0$ and ω should be understood to be $\omega + i\eta$ ($\eta = 0^+$).

Now we pick out from (E2),

$$f(r, Z_0) \equiv \int_0^\infty dp \frac{p}{(p^2 - \omega^2)^{3/2}} \exp\left(-i \left[pr - \sqrt{p^2 - \omega^2} Z_0 \right] \right). \quad (\text{E3})$$

We note that, for $r > Z_0$,

$$\oint_{C_-} dp \frac{p}{(p^2 - \omega^2)^{3/2}} \exp\left(-i \left[pr - \sqrt{p^2 - \omega^2} Z_0 \right] \right) = 0, \quad (\text{E4})$$

where C_- is the closed contour in a complex- p plane:

$$C_- = 0 \rightarrow R \rightarrow -iR \rightarrow 0 \quad (R \rightarrow \infty).$$

Here the contour segment $(0 \rightarrow R)$ is on the real axis, $(R \rightarrow -iR)$ is on $Re^{-i\theta}$ ($\theta = 0 \rightarrow \pi/2$), and $(-iR \rightarrow 0)$ is on the imaginary axis. The contribution from the contour segment $(R \rightarrow -iR)$ to (E4) vanishes in the limit $R \rightarrow \infty$.

The contribution from $(-iR \rightarrow 0)$ is pure imaginary. Then, $f(r, Z_0)$ in (E3) is pure imaginary. Since (E2) is pure real, when (E3) is used in (E2), (E3) should be canceled by other contribution(s), so that we can regard (E3) as vanishing.

In the complex- p plane, the integrand of (E3) has a branch point at $p = \omega + i\eta$. We take the branch cut to be a straight line $[\omega + i\eta, -\infty + i\eta)$. Then, for $r < Z_0$, we have

$$\oint_{C_+} dp \frac{p}{(p^2 - \omega^2)^{3/2}} \exp\left(-i \left[pr - \sqrt{p^2 - \omega^2} Z_0\right]\right) = 0, \quad (\text{E5})$$

where the closed contour C_+ is defined as

$$\begin{aligned} C_+ = 0 \rightarrow R \rightarrow iR \rightarrow i(\eta + 0^+) \rightarrow i(\eta + 0^+) + \omega - \alpha \\ \rightarrow i(\eta - 0^+) + \omega - \alpha \rightarrow i(\eta - 0^+) \rightarrow 0 \quad (R \rightarrow \infty). \end{aligned}$$

Here, $(R \rightarrow iR)$ is on $Re^{i\theta}$ ($\theta = 0 \rightarrow \pi/2$) and $(i(\eta + 0^+) + \omega - \alpha \rightarrow i(\eta - 0^+) + \omega - \alpha)$ is on a circle of radius α , whose center is at $p = \omega + i\eta$. From (E5), we obtain, in the limit $\eta \rightarrow 0^+$,

$$f(r, Z_0) = -2i \int_0^{\omega - \alpha} dp \frac{pe^{-ipr}}{(\omega^2 - p^2)^{3/2}} \cosh(\sqrt{\omega^2 - p^2} Z_0) + i\sqrt{\frac{2}{\omega\alpha}} e^{-i\omega r} - \pi Z_0 e^{-i\omega r}.$$

Here, as above, a pure-imaginary contribution has been dropped. Taking the limit $\alpha \rightarrow 0^+$, we obtain

$$\begin{aligned} f(r, Z_0) = -i\frac{2}{\omega} \int_0^1 d\xi \frac{\xi}{(1 - \xi^2)^{3/2}} \left[e^{-i\omega r \xi} \cosh(\sqrt{1 - \xi^2} \omega Z_0) - e^{-i\omega r} \right] \\ + i\frac{2}{\omega} e^{-i\omega r} - \pi Z_0 e^{-i\omega r}. \end{aligned}$$

Another type of integral in (E2) may be computed similarly. Up to a pure-imaginary contribution, we obtain

$$\begin{aligned} g(r, Z_0) \equiv \int_0^\infty dp \frac{p}{p^2 - \omega^2} e^{-i[pr - \sqrt{p^2 - \omega^2} Z_0]} \\ = \theta(Z_0 - r) \left[i\pi e^{-i\omega r} - 2 \int_0^1 d\xi \frac{\xi e^{-i\omega r \xi}}{1 - \xi^2} \sinh\left(\sqrt{1 - \xi^2} \omega Z_0\right) \right]. \end{aligned}$$

With these preliminaries, we can now compute (E2):

$$\begin{aligned} \mathcal{F}(X_0, Y_0; r) = \theta(X_0 + Y_0 - r) \theta(r - (X_0 - Y_0)) \frac{1}{\omega} \left\{ \frac{\pi\omega}{2} (X_0 + Y_0) \cos(\omega r) - \sin(\omega r) \right. \\ \left. + \int_0^1 d\xi \frac{\xi}{(1 - \xi^2)^{3/2}} \left[\sin(\omega r \xi) \cosh\left(\sqrt{1 - \xi^2} \omega (X_0 + Y_0)\right) - \sin(\omega r) \right] \right\} \\ + \theta(X_0 - Y_0 - r) \frac{1}{\omega} \left\{ \int_0^1 d\xi \frac{\xi \sin(\omega r \xi)}{(1 - \xi^2)^{3/2}} \sum_{\tau=\pm} \tau \cosh\left(\sqrt{1 - \xi^2} \omega (X_0 + \tau Y_0)\right) \right. \\ \left. - 2\omega Y_0 \int_0^1 d\xi \frac{\xi \sin(\omega r \xi)}{1 - \xi^2} \sinh\left(\sqrt{1 - \xi^2} \omega (X_0 - Y_0)\right) \right\}. \quad (\text{E6}) \end{aligned}$$

It is straightforward to obtain the approximate form for $\mathcal{F}(X_0, Y_0; r)$ in several limits for the arguments:

1. $X_0 - Y_0 < r < X_0 + Y_0$; $\omega r < \omega(X_0 + Y_0) \ll 1$:

$$\mathcal{F}(X_0, Y_0; r) \simeq \frac{\pi}{2} \left[X_0 + Y_0 - r \left\{ 1 - \frac{\omega^2 (X_0 + Y_0)^2}{4} \right\} \right].$$

2. $X_0 - Y_0 < r < X_0 + Y_0$; $1 \ll \omega(X_0 + Y_0)$ and $\omega r \ll \sqrt{\omega(X_0 + Y_0)}$:

$$\begin{aligned} \mathcal{F}(X_0, Y_0; r) \simeq \frac{\sqrt{2\pi}}{4} \frac{r}{(\omega(X_0 + Y_0))^{3/2}} e^{\omega(X_0 + Y_0)} \left[h(\omega(X_0 + Y_0)) - \frac{1}{2} \frac{\omega r^2}{X_0 + Y_0} \right] \\ + \frac{\pi}{2} (X_0 + Y_0) \cos \omega r, \end{aligned}$$

where

$$h(x) = 1 + \frac{21}{8} \frac{1}{x} + \dots$$

3. $r < X_0 - Y_0$; $\omega(X_0 + Y_0) \ll 1$:

$$\mathcal{F}(X_0, Y_0; r) \simeq \frac{\pi}{2} (\omega Y_0)^2 r.$$

4. $r < X_0 - Y_0$; $\omega(X_0 - Y_0) \ll 1 \ll \omega(X_0 + Y_0)$:

$$\mathcal{F}(X_0, Y_0; r) \simeq \frac{\sqrt{2\pi}}{4} \frac{r}{(\omega(X_0 + Y_0))^{3/2}} e^{\omega(X_0 + Y_0)} \left[h(\omega(X_0 + Y_0)) - \frac{1}{2} \frac{\omega r^2}{X_0 + Y_0} \right].$$

- [1] J. Schwinger, *J. Math. Phys.* **2** (1961), 407;
L. V. Keldysh, *Zh. Eksp. Teor. Fiz.* **47** (1964), 1515 (*Sov. Phys. JETP* **20** (1965), 1018);
R. A. Craig, *J. Math. Phys.* **9** (1968), 605;
J. Rammer and H. Smith, *Rev. Mod. Phys.* **58** (1986), 323;
- [2] K.-C. Chou, Z.-B. Su, B.-L. Hao, and L. Yu, *Phys. Rep.* **118** (1985), 1.
- [3] N. P. Landsman and Ch. G. van Weert, *Phys. Rep.* **145** (1987), 141.
- [4] H. Umezawa, “Advanced Field Theory — Micro, Macro, and Thermal Physics,” AIP, New York, 1993.
- [5] Proceedings of Quark Matter 1997, Tsukuba, Japan, to appear.
- [6] Proceedings of the fifth Workshop on Thermal Field Theories and Their Applications, Regensburg, Germany, 1998, to appear.
- [7] M. Gell-Mann and M. Lévy, *Nuovo Cim.* **16** (1960), 705.
- [8] A. Aoki et al., *Nucl. Phys. (Proc. Suppl.)* **63** (1998), 397.
- [9] R. D. Pisarski, *Phys. Rev. D* **52** (1995), R3773;
S. Chiku and T. Hatsuda, *Phys. Rev. D* **57** (1998), 6.
- [10] S. Chiku and T. Hatsuda, *Phys. Rev. D* **58** (1998), 076001.
- [11] K. Rajagopal and F. Wilczek, *Nucl. Phys. B* **404** (1993), 577;
S. Gavin, A. Gocksch, and R. D. Pisarski, *Phys. Rev. Lett.* **72** (1994), 2143;
S. Gavin and B. Müller, *Phys. Lett. B* **329** (1994), 486;
Z. Huang and X.-N. Wang, *Phys. Rev. D* **49** (1994), R4335;
J.-P. Blaizot and a. Krzywicki, *Phys. Rev. D* **50** (1994), 442;
D. Boyanovsky, H. J. de Vega, and R. Holman, *Phys. Rev. D* **51** (1995), 734;
F. Cooper, Y. Kluger, E. Mottola, and J. P. Paz, *Phys. Rev. D* **51** (1995), 2377;
A. Bialas, W. Czyz, and M. Gmyrek, *Phys. Rev. D* **51** (1995), 3739;
M. Asakawa, Z. Huang, and X.-N. Wang, *Phys. Rev. Lett.* **74** (1995), 3126;
D. Boyanovsky, M. D’Attanasio, H. J. de Vega, and P. Holman, *Phys. Rev. D* **54** (1996), 1748;
G. Amelino-Camelia, J. D. Bjorken, and S. E. Larsson, *Phys. Rev. D* **56** (1997), 6942;
H. Yabu, K. Nozawa, and T. Suzuki, *Phys. Rev. D* **57** (1998), 1687;
D. Boyanovsky, H. J. de Vega, R. Holman, S. P. Kumar, and R. D. Pisarski, *Phys. Rev. D* **57** (1998), 3653;
D. Boyanovsky, F. Cooper, H. J. de Vega, and P. Sodano, *Phys. Rev. D* **58** (1998), 025007;
See also, L. P. Csernai and I. N. Mishustin, *Phys. Rev. Lett.* **74** (1995), 5005;
J. Randrup, *Phys. Rev. D* **55** (1997), 1188;
M. Ishihara, M. Maruyama, and F. Takagi, *Phys. Rev. C* **57** (1998), 1440;
D. Boyanovsky, H. J. de Vega, R. Holman, S. P. Kumar, and R. D. Pisarski, hep-ph/9802370, February 1998;
A. Berera, M. Gleiser, and R. O. Ramos, hep-ph/9803394, March 1998;
E. R. Takano Natti, C.-Y. Lin, A. F. R. de Toledo Piza, and P. L. Natti, hep-th/9805060, May 1998.
- [12] J. D. Bjorken, *Acta Phys. Polon. B* **28** (1997), 2773.
- [13] E. Calzetta and B. L. Hu, *Phys. Rev. D* **37** (1988), 2878.
- [14] M. Le Bellac, “Thermal Field Theory,” Cambridge University Press, Cambridge, 1996.
- [15] Y. Yamanaka, H. Umezawa, K. Nakamura and T. Arimitsu, *Int. J. Mod. Phys. A* **9** (1994), 1153;
Y. Yamanaka and K. Nakamura, *Mod. Phys. Lett. A* **9** (1994), 2879;
H. Chu and H. Umezawa, *Int. J. Mod. Phys. A* **9** (1994), 1703; 2363. Earlier work is quoted in these papers.

- [16] C. Itzykson and J.-B. Zuber, “Quantum Field Theory,” McGraw-Hill, New York, 1980.
- [17] T. Muta, “Foundations of Quantum Chromodynamics,” World Scientific, Singapore, 1987.
- [18] R. Fausser and H. H. Walter, *Nucl. Phys. A* **584** (1995), 604.
- [19] R. D. Pisarski, *Phys. Rev. Lett.* **63** (1989), 1129;
E. Braaten and R. D. Pisarski, *Nucl. Phys. B* **337** (1990), 569;
J. Frenkel and J. C. Taylor, *Nucl. Phys. B* **334** (1990), 199.
- [20] M. E. Carrington, H. Defu, and M. H. Thoma, to be published in *Eur. phys. J. C*, (hep-ph/9708363, August 1997); in [5], (hep-ph/9808359, August, 1998).
- [21] A. Niégawa, *Phys. Lett. B* **247** (1990), 351;
N. Ashida, H. Nakkagawa, A. Niégawa, and H. Yokota, *Ann. Phys. (N.Y.)* **215** (1992), 315 [E: **230** (1994), 161]; *Phys. Rev. D* **45** (1992), 2066;
Asida, N., *Int. J. Mod. Phys. A* **8** (1993), 1729;
P. V. Landshoff, *Phys. Lett. B* **386** (1996), 291;
A. Niégawa, *Phys. Rev. D* **57** (1998), 1379;
See also, H. A. Weldon, *Phys. Rev. D* **28** (1983), 2007.
- [22] H. Matsumoto, I. Ojima, and H. Umezawa, *Ann. Phys. (N.Y.)* **152** (1984), 348;
A. J. Niemi and G. W. Semenoff, *Nucl. Phys. B* **230** [FS10] (1984), 181.
- [23] M. A. van Eijck and Ch. van Weert, *Phys. Lett. B* **278** (1992), 305;
P. Aurenche and T. Becherawy, *Nucl. Phys. B* **379** (1992), 259.
- [24] A. Niégawa, *Phys. Rev. D* **40** (1989), 1199;
R. J. Furnstahl and B. D. Serot, *Phys. Rev. C* **44** (1991), 2141;
F. Gelis, *Z. Phys. C* **70** (1996), 321;
H. Mabilat, *Z. Phys. C* **75** (1997), 155.
- [25] A. Niégawa, *Phys. Lett. B* **416** (1998), 137.
- [26] P. Zhuang and U. Heinz, *Phys. Rev. D* **57** (1998), 6525;
S. Ochs and U. Heinz, *Ann. Phys. (N.Y.)* **266** (1998), 351.
- [27] B. W. Lee, *Nucl. Phys. B* **9** (1969), 649;
T. Kugo, *Prog. Theor. Phys.* **57** (1977), 593.
- [28] S. Weinberg, *Phys. Rev. D* **9** (1974), 3357;
L. Dolan and R. Jackiw, *Phys. Rev. D* **9** (1974), 3320;
D. A. Kirzhnits and A. D. Linde, *Ann. Phys. (N.Y.)* **101** (1976), 195.
- [29] P. Fendley, *Phys. Lett. B* **196** (1987), 175.
- [30] I. D. Lawrie, in [6], (hep-ph/9809469 September, 1998). Earlier work is quoted therein.
- [31] R. Jackiw, *Phys. Rev. D* **9** (1974), 1686.
- [32] Y. Fujimoto, R. Grigjanis, and R. Kobes, *Prog. Theor. Phys.* **73** (1985), 434. Earlier work is quoted therein.
- [33] T. S. Evans, *Z. Phys. C* **36** (1987), 153.
- [34] J. W. Negele and H. Orland, “Quantum Many-Particle Systems,” Addison-Wesley, New York, 1988.
- [35] P. M. Stevenson, *Phys. Rev. D* **23** (1981) 2916.
- [36] G. Grunberg, *Phys. Lett.* **95B** (1980), 70.
- [37] E. J. Weinberg and A. Wu, *Phys. Rev. D* **36** (1987), 2474.
- [38] M. Morikawa, *Prog. Theor. Phys.* **93** (1995), 685;
C. Greiner and M. Müller, *Phys. Rev. D* **55** (1997), 1026;
A. Berera, M. Gleiser, and R. O. Ramos, in [11].
- [39] J. Rammer and H. Smith, *Rev. Mod. Phys.* **58** (1986), 323.
- [40] T. Altherr, *Phys. Lett. B* **341** (1995), 325.
- [41] R. Kobes, *Phys. Rev. D* **43** (1991), 1269.